

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MANIN'S PROOF OF THE MORDELL CONJECTURE OVER
FUNCTION FIELDS
Autor: Coleman, Robert F.
Kapitel: 2. Lang-Siegel towers
DOI: <https://doi.org/10.5169/seals-57915>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

in $X'(S)$. Suppose X is an Abelian scheme over S and R is the subgroup of $X(S)$ consisting of constant sections of X/S . Let $s \in X(S)$. Then the set $s + R$ is a set of bounded height.

LEMMA 3.1.1 (Manin). *Suppose E is a finite dimensional K vector subspace of $K(C)$. Then the set*

$$T = \{s \in C(S) : \exists k \neq 0 \in E \text{ such that } s*k = 0\}$$

has bounded height.

Proof. Without loss of generality we may increase E to suppose that the rational map $g: C \rightarrow \mathbf{P}_K(E)$ given on points by $x \rightarrow (e \in E \rightarrow e(x))$ is birational onto its image (note: g is actually a morphism on the complement of the polar locus of E). It follows that g induces an embedding of the generic fiber of C/S into $\mathbf{P}_{K(S)}(E \otimes K(S))$. Let h denote the logarithmic height with respect to this embedding. It follows that if $s \in C(S)$, $g \circ s$ is constant or $g \circ s$ has degree one. In the former case $h(s)$ is zero and the degree of the Zariski closure of $g \circ s(S)$ in $\mathbf{P}(E)$ in the latter.

Now if $s \in T$, and $g \circ s$ is not constant, it follows that the Zariski closure of $g \circ s(S)$ is a component of a hyperplane section of the Zariski closure of $g(C)$. Hence, $h(s)$ is less than or equal to the degree of the Zariski closure of $g(C)$. This proves the lemma. \square

The key property about heights we will need is:

THEOREM 3.1.2. *Suppose $C \rightarrow S$ is as in the above theorem. If $C(S)$ contains an infinite set of bounded height then C is a constant family.*

(See Corollary 2.2, Chapter 8 of [L-FD].)

Hence all we need prove is that the elements of $C(S)$ have bounded height.

2. LANG-SIEGEL TOWERS

Suppose the genus of C is at least 1. Suppose T is an infinite subset of $C(S)$.

PROPOSITION 3.2.1. *There exists a projective system of curves*

$(\{C_n\}, \{h_{m,n}\}), m, n \in \mathbf{Z}_{>0}$ and $n \leq m$, over K such that

- (i) $C_1 = C$,
- (ii) $h_{m,n}: C_m \rightarrow C_n$ is étale,

- (iii) $(h_{m,1})^{-1}(T) \cap C_m(S)$ is infinite,
- (iv) There exists a finite covering $S_{m,n}$ of S such that the fiber product of $h_{m,n}$ with $S_{m,n}$ is Galois, Abelian and of positive degree.

Let J denote the Jacobian scheme of C over S . Let $a: C \rightarrow J$ be an Albanese morphism. Let p be a prime. Let \bar{T} denote the closure of $a(T)$ in $J(S) \otimes \mathbf{Z}_p$. Since $a(T)$ is infinite it follow from the Mordell-Weil Theorem that there exists a $t \in \bar{T} - a(T)$. Let $t_n \in T$ such that $t - a(t_n) \in p^n J(S)$. Let C_n denote the normalization of the fiber-product of C and J via the map $H_n: x \rightarrow p^n x + t_n$ and $h_{n,1}$ the natural map from C_n to C . It follows that C_n is defined over S and since $H_m(J(S)) \supseteq \{t_n: m \mid n\}$ that $h_{n,1}(C_m(S))$ contains an infinite subset of T .

All that remains is to exhibit the maps $h_{m,n}$. Clearly, $t_m - t_n = p^n r_{m,n}$ for some $r_{m,n} \in J(S)$. Let $H_{m,n}$ denote the map $x: p^{m-n}x + r_{m,n}$. Then $H_{m,k} = H_{n,k} \circ H_{m,n}$. It follows that $H_{m,n}$ pulls back to a morphism $h_{m,n}: C_m \rightarrow C_n$. It is easy to see that this morphism becomes Abelian after adjoining the p^{m-n} -torsion points on J . This proves the proposition. \square

Remark. One can also prove the above proposition with the condition $n \leq m$ replaced by $n \mid m$.

3. COROLLARIES OF THE THEOREM OF THE KERNEL

LEMMA 3.3.1. *Suppose $g: X' \rightarrow X$ is a morphism of smooth proper schemes with geometrically connected fibers over S . Then if $\mu \in PF(X'/S)$ and $s, t \in X(S)$, $(g^*\mu)(s, t) = \mu(g \circ s, g \circ t)$.*

Proof. This follows easily from Lemma 1.3.2. \square

Suppose J is the Jacobian of C over S and g is an Albanese morphism, then since $g^*: H_{DR}^1(J/S) \rightarrow H_{DR}^1(C/S)$ is an isomorphism $g^*: PF(J/S) \rightarrow PF(C/S)$ is an isomorphism.

LEMMA 3.3.2. *Let μ be a fixed Picard-Fuchs differential equation on C/S . Then $\{\mu(s, t): s, t \in C(S)\}$ lies in a finite dimensional subspace of $K[S]$ over K .*

Proof. Suppose $\tilde{\mu} \in PF(J/S)$ such that $g^*\tilde{\mu} = \mu$. The lemma follows from the Mordell-Weil theorem which together with the Theorem of the kernel implies that $J(S)$ modulo the kernel of the homomorphism $s \rightarrow \tilde{\mu}(e, s)$ is a finitely generated Abelian group. \square