

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MANIN'S PROOF OF THE MORDELL CONJECTURE OVER FUNCTION FIELDS
Autor: Coleman, Robert F.
Kapitel: 3. COROLLARIES OF THE THEOREM OF THE KERNEL
DOI: <https://doi.org/10.5169/seals-57915>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

- (iii) $(h_{m,1})^{-1}(T) \cap C_m(S)$ is infinite,
- (iv) There exists a finite covering $S_{m,n}$ of S such that the fiber product of $h_{m,n}$ with $S_{m,n}$ is Galois, Abelian and of positive degree.

Let J denote the Jacobian scheme of C over S . Let $a:C \rightarrow J$ be an Albanese morphism. Let p be a prime. Let \bar{T} denote the closure of $a(T)$ in $J(S) \otimes \mathbf{Z}_p$. Since $a(T)$ is infinite it follows from the Mordell-Weil Theorem that there exists a $t \in \bar{T} - a(T)$. Let $t_n \in T$ such that $t - a(t_n) \in p^n J(S)$. Let C_n denote the normalization of the fiber-product of C and J via the map $H_n:x \mapsto p^n x + t_n$ and $h_{n,1}$ the natural map from C_n to C . It follows that C_n is defined over S and since $H_m(J(S)) \supseteq \{t_n : m \mid n\}$ that $h_{n,1}(C_m(S))$ contains an infinite subset of T .

All that remains is to exhibit the maps $h_{m,n}$. Clearly, $t_m - t_n = p^n r_{m,n}$ for some $r_{m,n} \in J(S)$. Let $H_{m,n}$ denote the map $x \mapsto p^{m-n}x + r_{m,n}$. Then $H_{m,k} = H_{n,k} \circ H_{m,n}$. It follows that $H_{m,n}$ pulls back to a morphism $h_{m,n}:C_m \rightarrow C_n$. It is easy to see that this morphism becomes Abelian after adjoining the p^{m-n} -torsion points on J . This proves the proposition. \square

Remark. One can also prove the above proposition with the condition $n \leq m$ replaced by $n \mid m$.

3. COROLLARIES OF THE THEOREM OF THE KERNEL

LEMMA 3.3.1. Suppose $g:X' \rightarrow X$ is a morphism of smooth proper schemes with geometrically connected fibers over S . Then if $\mu \in PF(X'/S)$ and $s, t \in X(S)$, $(g^*\mu)(s, t) = \mu(g \circ s, g \circ t)$.

Proof. This follows easily from Lemma 1.3.2. \square

Suppose J is the Jacobian of C over S and g is an Albanese morphism, then since $g^*:H_{DR}^1(J/S) \rightarrow H_{DR}^1(C/S)$ is an isomorphism $g^*:PF(J/S) \rightarrow PF(C/S)$ is an isomorphism.

LEMMA 3.3.2. Let μ be a fixed Picard-Fuchs differential equation on C/S . Then $\{\mu(s, t) : s, t \in C(S)\}$ lies in a finite dimensional subspace of $K[S]$ over K .

Proof. Suppose $\tilde{\mu} \in PF(J/S)$ such that $g^*\tilde{\mu} = \mu$. The lemma follows from the Mordell-Weil theorem which together with the Theorem of the kernel implies that $J(S)$ modulo the kernel of the homomorphism $s \mapsto \tilde{\mu}(e, s)$ is a finitely generated Abelian group. \square

LEMMA 3.3.3. Suppose A is an Abelian scheme over S such that $[W_{A/S}] = H^1_{DR}(A/S)$ and $g:C \rightarrow A$ is a non-constant morphism over S . Fix $s \in C(S)$. Then the set $T = \{t \in C(S) : (g^*\mu)(s, t) = 0 \text{ for all } \mu \in PF(A/S)\}$ is of bounded height.

Proof. Let A' denote the smallest Abelian subscheme of A over S containing $g(C)$. Since the map $g^*: PF(A/S) \rightarrow PF(A'/S)$ is surjective and $[W_{A/S}] = H^1_{DR}(A/S)$, it follows from Proposition 2.1.2 that $g(T)$ is contained in a translation of the group of constant sections of A'/S . Hence, $g(T)$ is a set of bounded height. Finally, since $C \rightarrow g(C)$ is a finite morphism, it follows that T is a set of bounded height. \square

In particular,

COROLLARY 3.3.4. Suppose A is an Abelian scheme over S such that $\kappa_{A/S}$ is an isomorphism and $g:C \rightarrow A$ is a non-constant morphism over S . Fix $s \in C(S)$. Then the set $\{t \in C(S) : (g^*\mu_\omega)(s, t) = 0 \text{ for all } \omega \in \omega_{A/S}\}$ is of bounded height.

4. PROOF OF MORDELL'S CONJECTURE

PROPOSITION 3.4.1. Suppose the kernel of the $\kappa_{C/S}$ has rank at least 2 over $K[S]$, then the points of $C(S)$ have bounded height.

Proof. Suppose $C(S)$ contains points of arbitrarily large height. Fix $s \in C(S)$. By shrinking S , if necessary, we may suppose that there exists a function $z \in K[S]$ such that $\Omega_S^1 = K[S]dz$ and there exists a finite covering \mathcal{L} of C by affine opens U and functions $v_U \in \mathcal{O}_C(U)$ such that $s \in U(S)$, and $\Omega_C^1(U)$ is spanned by dz and dv_U . We may also suppose that $s^*v_U = 0$ by replacing v_U with $v_U - (s \circ f)^*v_U$ if necessary. For $U \in \mathcal{L}, u \in \mathcal{O}_C(U)$ we define $\partial_{U,z}u$ and $\partial_{U,v}u$ by the equation

$$du = \partial_{U,z}udz + \partial_{U,v}udv_U.$$

Then $\partial_{U,z}$ is a lifting of $\partial = \partial/\partial z$. We set $\mu(t) = \mu(s, t)$ for

$$\mu \in PF = PF(C/S)$$

and $t \in C(S)$.

Let ω_1 and ω_2 be two independent elements in the kernel of $\kappa_{C/S}$. It follows that there exist ω'_1 and $\omega'_2 \in \omega_{C/S}$ such that

$$\partial[\omega'_i] = [\omega'_i].$$