

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 36 (1990)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TOPOLOGICAL SERIES OF ISOLATED PLANE CURVE SINGULARITIES  
**Autor:** Schrauwen, Robert  
**Anhang:** Appendix  
**DOI:** <https://doi.org/10.5169/seals-57905>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 06.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

interesting observation is, that in general the monomials of  $\varphi$  themselves will have a smaller order in  $t$  than  $\varphi$ .

#### 6.7. THE CASE THAT $f$ IS ARBITRARY.

If  $f = f_1^{m_1} \cdots f_r^{m_r} g$  with  $f_i$  irreducible,  $m_i \geq 2$  and  $g$  reduced, we still have that  $f + \varepsilon\varphi$  has the diagram of  $f$  with the multiple arrows replaced. We know exactly which replacements are possible (see section 3.8). To find out what is the type of  $f + \varepsilon\varphi$ , it again suffices to investigate linking behaviour. Some possibilities that only become apparent when  $f_i$  and  $f + \varepsilon\varphi$  are drawn in one diagram (that is the diagram of their product), have to be opted out by considering linking with cables which are known to be correct, using such valuations as  $\nu^{(2)}$ .

Although the tests become increasingly difficult, this gives a way to generalize theorem 6.5.

#### 6.8. IOMDIN TYPE SERIES.

We end with a remark on series of the form  $f + \varepsilon l^k$ , where  $l$  is a linear form not tangent to any branch of  $f$  and  $k \geq k_0$ , the largest polar ratio of  $f$ . These series have been studied by Iomdin and Lê, see [Lê], not only in the curve case but for general dimensions. Siersma [Si] has given a formula for the  $\Delta_*$  of these series. In the curve case this is just a special case of our results. Notice that:

$$\begin{aligned} \nu_i(l) &= d_i k \quad \text{where} \quad d_i = e_l(\Sigma_i) = \Sigma_i \cdot l, \\ \nu_i^{(2)}(l) &= 2d_i k. \end{aligned}$$

We would like to stress again that these Iomdin type series are generally much coarser than our topological series: they are single indexed and for example the Milnor number increases with steps of  $d = d_1 + \cdots + d_r$  within the series.

## APPENDIX










In this appendix the EN-diagrams of the series of plane curve singularities listed in [AGV] are drawn.

The first part consists of the *exceptional families*  $E$ ,  $W$  and  $Z$ .

The second part contains the *infinite series*  $A$ ,  $D$ ,  $J$ ,  $W$ ,  $W^\#$ ,  $X$ ,  $Y$  and  $Z$ . All variants are given. In the tables, we have that:

- (a)  $\mu$  = the Milnor number;  
 (b)  $N_0$  and the graph constant  $c$  are as in theorem 3.4;  
 (c)  $\Delta_*^\infty$  is the  $\Delta_*$  of the non-isolated singularity, the  $\Delta_*$  of an element of the series can be obtained by multiplying with  $t^{\Delta_* + c} - (-1)^\Delta$ .

Name	Formula	$\mu$	EN-diagram
$E_{6k}$	$y^3 + x^{3k+1}$	$6k$	
$E_{6k+1}$	$y^3 + x^{2k+1}y$	$6k + 1$	
$E_{6k+2}$	$y^3 + x^{3k+2}$	$6k + 2$	
$W_{12k}$	$y^4 + x^{4k+1}$	$12k$	
$W_{12k+1}$	$y^4 + yx^{3k+2}$	$12k + 1$	
$W_{12k+5}$	$y^4 + yx^{3k+2}$	$12k + 5$	
$W_{12k+6}$	$y^4 + yx^{3k+3}$	$12k + 6$	
$Z_{6k+11}$	$x(y^3 + yx^{2k+3} + x^{3k+4})$	$6k + 11$	
$Z_{6k+12}$	$x(y^3 + yx^{2k+3} + x^{3k+5})$	$6k + 12$	
$Z_{6k+13}$	$x(y^3 + yx^{2k+4} + x^{3k+5})$	$6k + 13$	

Name	Formula	$\mu$	EN-diagram	
$A_\infty$	$y^2$	0		$\Delta_*^\infty = \frac{1}{t^2 - 1}$
$A_0$	$y$	0		$p \geq 2, N = p + 1,$ $N_0 = 1, c = 0$
$A_1$	$y^2 + x^2$	1		
$A_p$	$y^2 + x^{p+1}$	$p$		
$D_\infty$	$xy^2$	1		$\Delta_*^\infty = 1$
$D_4$	$xy^2 + x^3$	4		$p \geq 5, N = p - 2,$ $N_0 = 2, c = 1$
$D_p$	$xy^3 + x^{p-1}$	$p$		
$J_{k,\infty}$	$y^3 + x^k y^2$	$3k - 2$		$\Delta_*^\infty = \frac{t^{3k} - 1}{t^3 - 1}$
$J_{k,0}$	$y^3 + x^k y + x^{3k}$	$6k - 2$		$k \geq 2, p \geq 1, c = k$ $N = p + 2k, N_0 = 2k$
$J_{k,p}$	$y^3 + x^k y^2 + x^{3k+p}$	$6k - 2 + p$		
$W_{k,\infty}$	$y^4 + y^2 x^{2k+1}$	$8k + 1$		$\Delta_*^\infty = \frac{t^{8k+4} - 1}{t^4 - 1}$
$W_{k,0}$	$y^4 + y^2 x^{2k+1} + x^{4k+2}$	$12k + 3$		$k \geq 1, p \geq 1,$ $N = p + 2k + 1$
$W_{k,p}$	$y^4 + y^2 x^{2k+1} + x^{4k+2+p}$	$12k + 3 + p$		

Name	Formula	$\mu$	EN-diagram	
$W_{k,\infty}^\#$	$(y^2 + x^{2k+1})^2$	$4k$		$\Delta_*^\infty = \frac{t^{4k+2} + 1}{t^4 - 1}$
$W_{k,2q-1}^\#$	$(y^2 + x^{2k+1})^2 + yx^{3k+1+q}$	$12k+2q+2$		$k \geq 1, q \geq 1,$ $c = 0$
$W_{k,2q}^\#$	$(y^2 + x^{2k+1})^2 + y^2x^{2k+1+q}$	$12k+2q+3$		$N = 8k + 2q + 3$ $N' = 8k + 2q + 4$
$X_\infty$	$y^4 + x^2y^2$	$5$		$\Delta_*^\infty = t^4 - 1$
$X_9$	$y^4 + x^2y^2 + x^4$	$9$		$p \geq 10, N = p - 7,$ $N_0 = 2, c = 2$
$X_p$	$y^4 + x^2y^2 + x^{4+p-9}$	$p$		
$X_{h,\infty}$	$y^4 + x^hy^3 + x^{2h}y^2$	$8h-3$		$\Delta_*^\infty = \frac{(t^{4h} - 1)^2}{t^4 - 1}$
$X_{h,0}$	$y^4 + x^hy^3 + x^{2h}y^2 + x^{3h}y$	$12h-3$		$h \geq 2, p \geq 1,$ $N = p + 2h,$ $N_0 = 2h, c = 2h$
$X_{h,p}$	$y^4 + x^hy^3 + x^{2h}y^2 + x^{4h+p}$	$12h-3+p$		
$Y_{\infty,\infty}$	$x^2y^2$	$4$		$\Delta_*^{\infty,\infty} = 1$
$Y_{r,\infty}$	$y^{4+r} + x^2y^2$	$r+5$		$r, s \geq 1,$ $c_1 = c_2 = 2$
$Y_{r,s}$	$y^{4+r} + x^2y^2 + x^{4+s}$	$9+r+s$		

Name	Formula	$\mu$	EN-diagram	
$Y_{\infty, \infty}^h$		$4h-2$		$h \geq 2, r, s \geq 1$
$Y_{r, s}^h$	See [AGV], p. 248	$12h+r+s-3$		
$Z_{k, \infty}$	$xy^3 + x^{k+2}y^2$	$3k+5$		$\Delta_*^\infty = t^{3k+4} - 1$
$Z_{k, 0}$	$xy^3 + x^{k+2}y^2 + x^{3k+4}$	$6k+9$		$k, p \geq 1, c = k + 2$ $N = p + 2k + 2,$ $N_0 = 2k + 2$
$Z_{k, p}$	$xy^3 + x^{k+2}y^2 + x^{3k+4+p}$	$6k+9+p$		
$Z_{k, \infty}^h$		$8h+3k-3$		$\Delta_*^\infty = \frac{(t^{4h}-1)(t^{4h+3k}-1)}{t^4-1}$
$Z_{k, 0}^h$	See [AGV], p. 249	$12h+6k-3$		$h \geq 2, k, p \geq 1,$ $N = p + 2h + 2k,$ $N_0 = 2h + 2k,$ $c = 2h + k$
$Z_{k, p}^h$	See [AGV], p. 249	$12h+6k-3+p$		

## REFERENCES

- [A'C] A'CAMPO, N. La fonction zêta d'une monodromie. *Comment. Math. Helv.* 50 (2) (1975), 233-248.
- [AGV] ARNOL'D, V.I., S.M. GUSEIN-ZADE and A.N. VARCHENKO. *Singularities of Differentiable Maps I*. Monographs in Mathematics 82, Birkhäuser, 1985.
- [BK] BRIESKORN, E. and H. KNÖRRER. *Ebene Algebraische Kurven*. Birkhäuser, 1981.
- [EN] EISENBUD, D. and W.D. NEUMANN. *Three-Dimensional Link Theory and Invariants of Plane Curve Singularities*. Annals of Mathematics Study 110, Princeton U.P., 1985.
- [Lê] LÊ, D. T. Ensembles analytiques complexes avec lieu singulier de dimension un (d'après I. N. Yomdin). *Séminaire sur les singularités*, Publ. Math. Univ. Paris VII (1980), 87-95.