Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 36 (1990)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE INVERSIVE DIFFERENTIAL GEOMETRY OF PLANE CURVES

Autor: Cairns, G. / Sharpe, R. W.

Kapitel: §5. A GENERALIZED FOUR VERTEX THEOREM

DOI: https://doi.org/10.5169/seals-57907

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 16.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Combining this with 4.5 yields

$$\mathbf{n}(\tau(f)) - \tau(\mathbf{n}(f)) = [\mathbf{n}, \tau](f) = \kappa_1\tau(f) + \mu_1\mathbf{n}(f) = \kappa_1\tau(f) + \mathbf{n}\tau(f)$$

which gives 4.6 as required.

§5. A GENERALIZED FOUR VERTEX THEOREM

The curves of constant curvature in the round 2-sphere S^2, the upper half plane H^2 (hyperbolic space), and the Euclidean plane $\mathbf{R}^2$ are just the circles. Moreover, the stereographic projection $p: S^2 \to \mathbf{R}^2$ and the inclusion $i: H^2 \to \mathbf{R}^2$ both preserve these circles. Thus theorem 4.2 says that our form

$$\omega = \sqrt{|\kappa'(s)|}\, ds$$

along a curve γ in $\mathbf{R}^2$ pulls back via p or i to the form

$$\omega = \sqrt{|\kappa_g'(s)|}\, ds$$

along the corresponding curve γ', where here $\kappa_g(s)$ and s refer to the geodesic curvature and arc-length of γ' in the metric for S^2 or H^2. Thus we obtain the four vertex theorem for S^2 and H^2. It follows that the four vertex theorem holds for all complete simply connected Riemannian surfaces of constant curvature. Finally if γ is a null-homotopic smooth simple closed curve on an arbitrary complete Riemannian surface M of constant curvature, then γ lifts one-to-one to a smooth simple closed curve with the same number of vertices on the simply connected universal cover of M. Once again it follows that the number of vertices is at least four.

Remark 5.1. Interestingly, simple closed homotopically non-trivial curves in the real projective plane always have at least three vertices [17]. Note that in non-orientable surfaces the number of honest vertices of a closed curve need not necessarily be even, since here geodesic curvature is only defined up to a sign.

§6. NORMAL FORM AND INVERSIVE CURVATURE

Let p be a non-vertex point of an oriented curve γ. Since the subgroup of Euclidean motions in G acts transitively on the points of $\mathbf{R}^2$ and the unit tangent vectors at these points, we may assume that the point $p \in \gamma$ which