

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 37 (1991)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE AVERAGE BEHAVIOUR OF THE LARGEST DIVISOR OF  $n$  PRIME TO A FIXED INTEGER  $k$   
**Autor:** PÉTERMANN, Y.-F. S.  
**Kapitel:** 2. The case  $(k) = 2$   
**DOI:** <https://doi.org/10.5169/seals-58739>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 18.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$$(1.16) \quad D_h(\delta) = D_{0,h}(\delta) \quad (\delta = \beta \text{ or } \gamma).$$

In view of (0.14) this implies that

$$(1.17) \quad \begin{aligned} 1 - D_h(S_k - \varepsilon) &\geq D_h\left(S_k - \frac{3\varepsilon}{4}\right) - D_h(S_k - \varepsilon) \\ &\geq D_h(\gamma) - D_h(\beta) = D_{0,h}(\gamma) - D_{0,h}(\beta) \geq \frac{1}{L} \cdot \frac{\varepsilon}{5} \cdot \frac{\sigma(k)}{k}. \quad \square \end{aligned}$$

*Remark.* I studied in [P2] an error term associated with the  $k$ -th Jordan totient function (for  $k \geq 2$ ), that can be expressed in terms of the function

$$(1.18) \quad g_k(x) := - \sum_{n=1}^{\infty} \frac{\mu(n)}{n^k} \psi\left(\frac{x}{n}\right),$$

where  $\mu$  denotes the Moebius function, and I proved by a direct method that

$$(1.19) \quad \liminf_{x \rightarrow \infty} g_k(x) = - \limsup_{x \rightarrow \infty} g_k(x).$$

This can also be obtained by an argument similar to the above proof.

## 2. THE CASE $\omega(k) = 2$

In this section we obtain an estimate more general than (0.10) of Theorem 2.

**THEOREM 2'.** *Let  $k = pq$  where  $p < q$  and  $p$  and  $q$  are prime numbers, and let  $d = q - ps$  with  $1 \leq d \leq p - 1$  be the remainder of the Euclidean division of  $q$  by  $p$ . Then we have*

$$(2.1) \quad S_k \geq \frac{k}{\sigma(k)} + \frac{1}{(p+1)} - \frac{pd}{(p+1)(q+1)} + \frac{(p+1)(p-2)(q-1)}{p^2q}.$$

The right side of (2.1) is easily seen to exceed  $k/\sigma(k)$  for any  $p$  and  $q$ . And in the special case where  $p = 2$  it reduces to  $\left(q - \frac{1}{3}\right)/(q+1)$ .

*Proof.* Let  $N$  be a positive integer. We define, modulo  $p^N q^N$ , the integer  $x = x_N$  by the system of congruences

$$(2.2) \quad \begin{cases} x \equiv -1(p^N) \\ x \equiv -d-1(q^N). \end{cases}$$

We have, for  $1 \leq i \leq N$  and  $1 \leq j \leq N$ ,

$$(2.3) \quad x \equiv s_{i,j} q^j - d - 1 (p^i q^j) \quad \text{where} \quad \begin{cases} s_{1,1} = 1 \\ 1 \leq s_{i,j} \leq p^i - 1, \end{cases}$$

whence

$$(2.4) \quad \begin{aligned} H(k, x) &\geq \frac{1}{2} + \sum_{i=1}^N \frac{(1-p)}{p^i} \left( -\frac{1}{2} + \frac{1}{p^i} \right) + \sum_{j=1}^N \frac{(1-q)}{q^j} \left( -\frac{1}{2} + \frac{d+1}{q^j} \right) \\ &+ \frac{(p-1)(q-1)}{pq} \left( \frac{1}{2} - \frac{q-d-1}{pq} \right) \\ &+ \sum_{\substack{1 \leq i, j \leq N \\ (i,j) \neq (1,1)}} \frac{(p-1)(q-1)}{p^i q^j} \left( \frac{1}{2} - \frac{(p^i-1)q^j - d - 1}{p^i q^j} \right) + o_N(1). \end{aligned}$$

The right side of (2.4) tends to the right side of (2.1) as  $N \rightarrow \infty$ , and the theorem is proved in virtue of (0.15).  $\square$

### PROOF OF THEOREM 3

The function  $f_r$  defined in (0.11) satisfies, provided  $r \geq 3$ ,

$$(3.1) \quad f_r(p_2, \dots, p_r) < f_{r-1}(p_2, \dots, p_{r-1}) \leq p_2,$$

and thus the condition

$$(3.2) \quad f_r(p_2, \dots, p_r) \geq x$$

implies, for any  $x$ , that

$$(3.3) \quad p_2 \begin{cases} > x & \text{if } r \geq 3, \\ \geq x & \text{if } r = 2. \end{cases}$$

Also note that, since

$$(3.4) \quad \sum_{n=1}^{\infty} \frac{\gamma_k(n)}{n} = \prod_{p|k} \left( 1 + (1-p) \sum_{i \geq 1} \frac{1}{p^i} \right) = 0,$$

we have in fact

$$(3.5) \quad H(k, x) = - \sum_{n \geq 1} \frac{\gamma_k(n)}{n} \left\{ \frac{x}{n} \right\}.$$