

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 37 (1991)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE EVALUATION OF SELBERG CHARACTER SUMS  
**Autor:** Evans, Ronald J.  
**Kapitel:** §1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-58741>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## THE EVALUATION OF SELBERG CHARACTER SUMS

by Ronald J. EVANS

ABSTRACT. The evaluations of Selberg character sums conjectured on p. 207 of *Enseignement Math.* 27 (1981) are proved.

### § 1. INTRODUCTION

Many of the classical special functions over  $\mathbf{C}$  have character sum analogs over finite fields. For example, the Gauss and Jacobi sums defined in (1.1) are analogs of the gamma and beta integrals

$$\Gamma(a) = \int_0^\infty e^{-x} x^a \frac{dx}{x}, \quad \beta(a, b) = \int_0^1 x^a (1-x)^b \frac{dx}{x(1-x)}.$$

Some identities for character sums over finite fields seem more difficult to prove than their classical counterparts; compare, e.g., the Hasse-Davenport product formula for Gauss sums [7, (7)] with the Gauss multiplication formula for gamma functions. The identities for  $n$ -dimensional Selberg character sums given in Theorems 1.1, 1.1a provide further examples. Their counterparts are the well known  $n$ -dimensional Selberg integral extensions of the gamma and beta integral formulas.

The case  $n = 3$  of the Selberg character sum identity in Theorem 1.1 has been used to evaluate a sum connected with the root system  $G_2$  [8]. The case  $n = 2$  is equivalent to an analog of Dixon's summation formula [11, (2.1.5)] involving hypergeometric  ${}_3F_2$  character sums over finite fields. We remark that hypergeometric character sums have been used, e.g., in the computation of the number of points on hypersurfaces [13], [12], in proving congruences for Apery numbers [14], and in graph theory [6], [9].

Let  $GF(q)$  be a finite field of  $q$  elements, where  $q$  is a power of an odd prime. Fix a multiplicative character  $\tau: GF(q)^* \rightarrow \mathbf{C}^*$  of order  $q - 1$  and a nontrivial additive character  $\psi: GF(q) \rightarrow \mathbf{C}^*$ . Extend  $\tau$  by defining  $\tau(0) = 0$ . Let  $\phi = \tau^{(q-1)/2}$  be the quadratic character on  $GF(q)$ . For all integers  $a, b$ , define the Gauss sums  $G(a)$  and Jacobi sums  $J(a, b)$  by

$$(1.1) \quad G(a) = \sum_{\xi \in GF(q)^*} \tau(\xi)^a \psi(\xi), \quad J(a, b) = \sum_{1 \neq \xi \in GF(q)^*} \tau(\xi)^a \tau(1 - \xi)^b.$$

For integers  $n \geq 0$  and  $a, b, c > 0$ , define the Selberg character sums

$$(1.2) \quad S_n(a, b, c) = \sum_E \tau((-1)^{an} E(0)^a E(1)^b \Delta_E^c) \phi(\Delta_E),$$

$$(1.2a) \quad S_n(a, c) = \sum_E \psi(e_{n-1}) \tau(E(0)^a \Delta_E^c) \phi(\Delta_E),$$

$$(1.2b) \quad S_n(c) = \sum_E \psi(e_{n-1}^2 / 2 - e_{n-2}) \tau(\Delta_E)^c \phi(\Delta_E),$$

where each sum is over all monic polynomials

$$(1.3) \quad E = E(x) = x^n + e_{n-1} x^{n-1} + e_{n-2} x^{n-2} + \cdots + e_0$$

of degree  $n$  over  $GF(q)$ , and where  $\Delta_E$  denotes the discriminant of  $E$  (with the convention that  $\Delta_E = 1$  when  $\deg(E) \leq 1$ ). Define the following products:

$$(1.4) \quad P_n(a, b, c) = \prod_{j=0}^{n-1} \frac{G(a+jc) G(b+jc) G(c+jc) \bar{G}(a+b+(n-1+j)c)}{qG(c)},$$

$$(1.4a) \quad P_n(a, c) = \prod_{j=0}^{n-1} \frac{G(a+jc) G(c+jc)}{G(c)},$$

$$(1.4b) \quad P_n(c) = \prod_{j=0}^{n-1} \frac{G(c+jc) \phi(2) G((q-1)/2)}{G(c)},$$

where  $\bar{G}$  denotes the complex conjugate of  $G$ .

The object of this paper is to prove Theorems 1.1, 1.1a, and 1.1b below. These results, analogs of  $n$ -dimensional integral formulas of Selberg [3, (1.1), (1.3), (1.2)], [2], verify conjectures made in 1981 [7, (29), (29a), (29b)]. The decisive breakthrough came in 1990 when Anderson [1] proved a somewhat weakened form of Theorem 1.1. The proofs here are based on modifications of the method in [1]. The modifications are designed to handle complications arising from “imprimitive”  $L$ -functions (see §2).

**THEOREM 1.1.** *For all integers  $n, a, b, c > 0$ , if none of*

$$a + b + (n-1+j)c \quad (0 \leq j \leq n-1)$$

*are divisible by  $q-1$ , then  $S_n(a, b, c) = P_n(a, b, c)$ .*

**THEOREM 1.1a.** *For all integers  $n, a, c > 0$ ,  $S_n(a, c) = P_n(a, c)$ .*

**THEOREM 1.1b.** *For all integers  $n, c > 0$ ,  $S_n(c) = P_n(c)$ .*

Given a monic polynomial  $E$  over  $GF(q)$ , define  $\sigma(E) = 0$  if  $E$  is not squarefree,  $\sigma(E) = 1$  if  $E = 1$ , and otherwise let  $\sigma(E)$  denote the sign of the permutation of the zeros of  $E$  effected by the  $q^{\text{th}}$  power automorphism of  $GF(q)$ . For odd  $q$ ,  $\sigma(E) = \phi(\Delta_E)$ . If  $\phi(\Delta_E)$  is replaced by  $\sigma(E)$  in the definitions (1.2), (1.2a) of  $S_n(a, b, c)$ ,  $S_n(a, c)$ , then Theorems 1.1 and 1.1a remain valid without the stipulation “ $q$  odd”; the proofs for even  $q$  are virtually the same. This observation is due to Serre; see [1].

The following result is equivalent to Theorem 1.1, as was shown in [10, p. 116].

**THEOREM 1.2.** *For integers  $n, a, b, c > 0$ , if none of  $a + jc$  ( $0 \leq j \leq n - 1$ ) are divisible by  $q - 1$ , or if none of  $b + jc$  ( $0 \leq j \leq n - 1$ ) are divisible by  $q - 1$ , or if none of  $a + b + (n - 1 + j)c$  ( $0 \leq j \leq n - 1$ ) are divisible by  $q - 1$ , then  $S_n(a, b, c) = P_n(a, b, c)$ .*

Theorems 1.3 and 1.4 below, analogs of more recent Selberg integral formulas (see [4]), were stated as conjectures in [5]. They are consequences of Theorems 1.1a and 1.1b, respectively, as is shown in [5, Theorems 2.2 and 2.5].

**THEOREM 1.3.** *For all integers  $n, a, b, c > 0$ ,*

$$\sum_E \tau(E(0)^a (1 + e_{n-1})^b \Delta_E^c) \phi(\Delta_E) = \begin{cases} \frac{G(-b - na - n(n-1)c)}{G(-b)} P_n(a, c) , & \text{if } b \not\equiv 0 \pmod{q-1} \\ \frac{\tau(-1)^{an} G(b)}{G(b + na + n(n-1)c)} P_n(a, c) , & \text{if } b + na + n(n-1)c \\ & \not\equiv 0 \pmod{q-1} , \end{cases}$$

where the sum is over all polynomials  $E$  of degree  $n$  given by (1.3).

**THEOREM 1.4.** *For  $w \in GF(q)^*$  and all integers  $n, b, c > 0$  with  $b \not\equiv 0 \pmod{q-1}$ ,*

$$\sum_E \tau((w + e_{n-1}^2/2 - e_{n-2})^b \Delta_E^c) \phi(\Delta_E) = \tau(w)^{b+n(q-1)/2+cn(n-1)/2} \frac{G(-b - cn(n-1)/2 - n(q-1)/2)}{G(-b)} P_n(c) ,$$

where the sum is over all polynomials  $E$  of degree  $n$  given by (1.3).

*Acknowledgement.* We are very grateful to G. W. Anderson for helpful correspondence on  $L$ -functions and character sums.

## § 2. $L$ -FUNCTIONS

Throughout this section,  $V$  denotes a *monic* polynomial over  $GF(q)$ , and  $v$  ranges over the distinct monic irreducible factors of  $V$  over  $GF(q)$ . Write

$$(2.1) \quad V = \prod_{v|V} v^{\text{ord}_v V}, \quad F = F_V = \prod_{v|V} v.$$

If no exponent  $\text{ord}_v V$  in (2.1) is divisible by  $q - 1$ , then  $V$  is said to be *primitive*. Note that  $V = 1$  is primitive. For any monic polynomial

$$(2.2) \quad W = W(x) = x^n + w_{n-1}x^{n-1} + w_{n-2}x^{n-2} + \cdots + w_0$$

over  $GF(q)$ , set

$$(2.3) \quad \alpha(W) = w_{n-1}, \quad \beta(W) = w_{n-1}^2/2 - w_{n-2}.$$

Define the  $L$ -functions

$$(2.4) \quad L(t, V) = \sum_W \tau(R(V, W)) t^{\deg W},$$

$$(2.4a) \quad L_1(t, V) = \sum_W \psi(\alpha(W)) \tau(R(V, W)) t^{\deg W},$$

$$(2.4b) \quad L_2(t, V) = \sum_W \psi(\beta(W)) \tau(R(V, W)) t^{\deg W},$$

where in each sum,  $W$  ranges over all monic polynomials over  $GF(q)$ , and  $R(V, W)$  is the resultant of  $V$  and  $W$ . It is easily checked that

$$(2.5) \quad \begin{aligned} L(t, 1) &= (1 - qt)^{-1}, & L_1(t, 1) &= 1, \\ L_2(t, 1) &= 1 + \phi(2)G((q-1)/2)t. \end{aligned}$$

Since the summands in (2.4), (2.4a), (2.4b) are multiplicative in  $W$ , each of the  $L$ -functions has an Euler product expansion. Thus we have the following result.

LEMMA 2.1. *Write  $V = GH$  where  $G$  and  $H$  are monic, relatively prime polynomials over  $GF(q)$  with  $G$  primitive and  $H$  a  $(q-1)$ th power. Then*

$$(2.6) \quad L(t, V) = L(t, G) \prod_{v|H} (1 - \tau(R(G, v))t^{\deg v}),$$

$$(2.6a) \quad L_1(t, V) = L_1(t, G) \prod_{v|H} (1 - \psi(\alpha(v))\tau(R(G, v))t^{\deg v}),$$