

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 37 (1991)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE DIFFEOMORPHISM GROUPS OF CERTAIN ALGEBRAIC SURFACES
Autor: Ebeling, Wolfgang / Okonek, Christian
Kapitel: 3. NON-REALIZABLE ISOMETRIES
DOI: <https://doi.org/10.5169/seals-58742>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Let X be any simply connected algebraic surface with a big monodromy group. If $p_g(X) \equiv 0 \pmod{2}$ then the degree of $\Phi_l(X)$ is odd. If $p_g(X) \equiv 1 \pmod{2}$ and $k_X^2 \equiv 1 \pmod{2}$ then the degree of $\Phi_{k_X, \alpha, P}(X)$ is odd. So k_X divides $\Phi_l(X)$ or $\Phi_{k_X, \alpha, P}(X)$ in these cases.

Remark. Theorem 4 and its corollary remain true for polynomials $\Phi_{c, \alpha, P}(X)$ if $c \in H^2(X, \mathbf{Z})$ is a class with $\bar{c} \neq 0$ such that $\overline{f^*(c)} = \bar{c}$ for all $f \in \psi(\text{Diff}_+(X))$. The question which elements of $H^2(X, \mathbf{Z})$ or $H^2(X, \mathbf{Z}/2)$ have this invariance property will be treated in §4.

3. NON-REALIZABLE ISOMETRIES

We shall show that for a simply connected algebraic surface with odd geometric genus, -1 is not induced by an orientation preserving diffeomorphism. For K3 surfaces this was shown by Donaldson in the proof of [D, Proposition 6.2]. There he proves the nontriviality of a certain polynomial $\Phi_{c, \alpha, P}(X)$ for a K3 surface X . With Zuo's nontriviality result (Theorem 3) we are able to generalize this as follows.

THEOREM 6. *If X is a simply connected algebraic surface with $p_g(X) \equiv 1 \pmod{2}$ then $-1 \notin \psi(\text{Diff}_+(X))$.*

Proof. Suppose that there is an orientation preserving diffeomorphism $f: X \rightarrow X$ such that $f^* = -1$. Let $c \in H^{1,1}(X, \mathbf{Z})$ be a class with $\bar{c} \neq 0$, and choose a principal $SO(3)$ -bundle P with $w_2(P) = \bar{c}$ such that $\Phi_{c, \alpha, P}(X)$ is nontrivial. This is possible according to Theorem 3. Then

$$f^* \Phi_{c, \alpha, P}(X) = (-1)^d \Phi_{c, \alpha, P}(X),$$

since $\Phi_{c, \alpha, P}(X)$ is a polynomial of degree d on L .

On the other hand, by §2(c)

$$f^* \Phi_{c, \alpha, P}(X) = \Phi_{f^*c, f^*\alpha, f^*P}(X).$$

We have $f^*c = -c$ and $f^*\alpha = -\alpha$ because $f^* = -1$ and the dimension of α is odd. Since f is orientation preserving and $f^* = -1$ we find $f^*p_1(P) = p_1(P)$ and $f^*w_2(P) = w_2(P)$, so that the bundle f^*P is isomorphic to P . Therefore

$$f^* \Phi_{c, \alpha, P}(X) = \Phi_{-c, -\alpha, P}(X) .$$

Applying §2 (a) and (b) with $a = -c$ we get

$$f^* \Phi_{c, \alpha, P}(X) = -\Phi_{-c, \alpha, P}(X) = -(-1)^{c^2} \Phi_{c, \alpha, P}(X) .$$

By assumption $\Phi_{c, \alpha, P}(X) \neq 0$, so we must have

$$(-1)^{c^2+1} = (-1)^d .$$

Now $d = 4c_2 - c^2 - 3(1 + p_g(X))$ implies that $(-1)^{p_g(X)} = 1$, i.e. $p_g(X) \equiv 0 \pmod{2}$. This proves Theorem 6.

4. DIFFEOMORPHISM GROUPS OF SOME ALGEBRAIC SURFACES

In §1 we saw that the image of ψ contains the group $O'_k(L) \cdot \{\sigma_*, \text{id}\}$ in many cases. In §2 we showed that under certain conditions $\{\pm k_X\}$ is invariant under $\psi(\text{Diff}_+(X))$. Finally we proved in the previous section that for algebraic surfaces of odd geometric genus -1 is not induced by an orientation preserving diffeomorphism. It turns out that these facts suffice to determine the image of ψ .

PROPOSITION 7. *Let X be a simply connected algebraic surface which satisfies the following conditions:*

- (i) $O'_k(L) \cdot \{\sigma_*, \text{id}\} \subset \psi(\text{Diff}_+(X))$,
- (ii) $\{\pm k_X\}$ is invariant under $\psi(\text{Diff}_+(X))$,
- (iii) $-1 \notin \psi(\text{Diff}_+(X))$.

Then

$$\psi(\text{Diff}_+(X)) = O'_k(L) \cdot \{\sigma_*, \text{id}\} .$$

Proof. Let $g = -\sigma_*$. Then $g \in O_k(L)$, but $g \notin \psi(\text{Diff}_+(X))$, since $-1 \notin \psi(\text{Diff}_+(X))$. Hence by (i), $g \notin O'_k(L)$. Therefore

$$O_k(L) = O'_k(L) \cdot \{g, \text{id}\} .$$

Now let $h \in \psi(\text{Diff}_+(X))$. By (ii) either $h(k) = k$ or $h(k) = -k$. In the first case $h \in O_k(L)$. Moreover, $h \in O'_k(L)$ since otherwise $h = gh_0$ for some $h_0 \in O'_k(L)$ which would imply $g \in \psi(\text{Diff}_+(X))$, a contradiction. In the second case we have $h' = h\sigma_* \in O_k(L)$. By the same argument as before we see that $h' \in O'_k(L)$. Hence $h = h'\sigma_* \in O'_k(L) \cdot \{\sigma_*, \text{id}\}$. This proves Proposition 7.

Putting everything together we get the main result of our paper.