

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	37 (1991)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ON THE DIFFEOMORPHISM GROUPS OF CERTAIN ALGEBRAIC SURFACES
Autor:	Ebeling, Wolfgang / Okonek, Christian
Kapitel:	4. Diffeomorphism groups of some algebraic surfaces
DOI:	https://doi.org/10.5169/seals-58742

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$f^* \Phi_{c, \alpha, P}(X) = \Phi_{-c, -\alpha, P}(X) .$$

Applying §2(a) and (b) with $a = -c$ we get

$$f^* \Phi_{c, \alpha, P}(X) = -\Phi_{-c, \alpha, P}(X) = -(-1)^{c^2} \Phi_{c, \alpha, P}(X) .$$

By assumption $\Phi_{c, \alpha, P}(X) \not\equiv 0$, so we must have

$$(-1)^{c^2+1} = (-1)^d .$$

Now $d = 4c_2 - c^2 - 3(1 + p_g(X))$ implies that $(-1)^{p_g(X)} = 1$, i.e. $p_g(X) \equiv 0 \pmod{2}$. This proves Theorem 6.

4. DIFFEOMORPHISM GROUPS OF SOME ALGEBRAIC SURFACES

In §1 we saw that the image of ψ contains the group $O'_k(L) \cdot \{\sigma_*, \text{id}\}$ in many cases. In §2 we showed that under certain conditions $\{\pm k_X\}$ is invariant under $\psi(\text{Diff}_+(X))$. Finally we proved in the previous section that for algebraic surfaces of odd geometric genus -1 is not induced by an orientation preserving diffeomorphism. It turns out that these facts suffice to determine the image of ψ .

PROPOSITION 7. *Let X be a simply connected algebraic surface which satisfies the following conditions:*

- (i) $O'_k(L) \cdot \{\sigma_*, \text{id}\} \subset \psi(\text{Diff}_+(X))$,
- (ii) $\{\pm k_X\}$ is invariant under $\psi(\text{Diff}_+(X))$,
- (iii) $-1 \notin \psi(\text{Diff}_+(X))$.

Then

$$\psi(\text{Diff}_+(X)) = O'_k(L) \cdot \{\sigma_*, \text{id}\} .$$

Proof. Let $g = -\sigma_*$. Then $g \in O_k(L)$, but $g \notin \psi(\text{Diff}_+(X))$, since $-1 \notin \psi(\text{Diff}_+(X))$. Hence by (i), $g \notin O'_k(L)$. Therefore

$$O_k(L) = O'_k(L) \cdot \{g, \text{id}\} .$$

Now let $h \in \psi(\text{Diff}_+(X))$. By (ii) either $h(k) = k$ or $h(k) = -k$. In the first case $h \in O_k(L)$. Moreover, $h \in O'_k(L)$ since otherwise $h = gh_0$ for some $h_0 \in O'_k(L)$ which would imply $g \in \psi(\text{Diff}_+(X))$, a contradiction. In the second case we have $h' = h\sigma_* \in O_k(L)$. By the same argument as before we see that $h' \in O'_k(L)$. Hence $h = h'\sigma_* \in O'_k(L) \cdot \{\sigma_*, \text{id}\}$. This proves Proposition 7.

Putting everything together we get the main result of our paper.

THEOREM 8. *Let X be a simply connected algebraic surface with $p_g(X) \equiv 1 \pmod{2}$ and $k_X^2 \equiv 1 \pmod{2}$. If X is either*

- (i) *a complete intersection, or*
- (ii) *a Moishezon or Salvetti surface,*

then

$$\psi(\text{Diff}_+(X)) = O'_k(L) \cdot \{\sigma_*, \text{id}\}.$$

Example. Consider a complete intersection surface X of multidegree (d_1, \dots, d_r) . Using the formulas of [E4], we can translate the conditions of Theorem 8 into numerical conditions on the degrees d_i . The condition $k_X^2 \equiv 1 \pmod{2}$ is equivalent to

$$(1) \quad d_i \equiv 1 \pmod{2} \quad \text{for } i = 1, \dots, r.$$

Write $d_i = 2e_i + 1$ ($i = 1, \dots, r$). Then $p_g(X) \equiv 1 \pmod{2}$ is equivalent to the following two conditions

$$(2) \quad \sum_{i < j} e_i e_j \equiv 1 \pmod{2},$$

$$(3) \quad \text{either } 3 \mid d_j \text{ for some } j, 1 \leq j \leq r, \quad \text{or} \quad 3 \mid \sum_{i=1}^r e_i(e_i + 1).$$

In particular there is an infinite sequence of complete intersection surfaces satisfying the conditions of Theorem 8, e.g. the surfaces with $(d_1, d_2) = (3, 3 + 4m)$, $m \in \mathbf{Z}$, $m \geq 0$.

We leave it to the reader to formulate similar conditions for the case (ii) of Theorem 8.

We shall give two further applications of the results of the first three sections.

Let X be a surface as in Theorem 1 and denote the symmetric bilinear form corresponding to q_X by $\langle \cdot, \cdot \rangle$. Define $L' := \ker k_X = k_X^\perp \subset L$, and let $\Delta \subset L'$ be the set of vanishing cycles of X (cf. [EO]). The pair (L', Δ) is then a vanishing lattice in the sense of [E1, Definition (2.1)]. This means the following. If $\delta \in \Delta$, then $\langle \delta, \delta \rangle = -2$ and one has an associated reflection s_δ defined by

$$s_\delta(x) = x + \langle x, \delta \rangle \delta$$

for all $x \in L'$. Let Γ_Δ denote the subgroup of $O(L')$ generated by these reflections s_δ , $\delta \in \Delta$. Then (L', Δ) satisfies the following conditions:

- (i) $\langle \delta, \delta \rangle = -2$ for all $\delta \in \Delta$.
- (ii) Δ generates L' .
- (iii) Δ is a Γ_Δ -orbit.
- (iv) Unless $\text{rank } L' = 1$, there exist $\delta_1, \delta_2 \in \Delta$ with $\langle \delta_1, \delta_2 \rangle = 1$.

As in Wall's paper [W] we can derive from a statement about $\psi(\text{Diff}_+(X))$ a statement about the possibility of representing homology classes by embedded 2-spheres.

THEOREM 9. *Let X be an algebraic surface, and let $x \in H_2(X, \mathbf{Z})$ be a class with $q_X(x) = -2$. If x is represented by a differentiably embedded 2-sphere, then $\bar{x} \in \bar{k}_X^\perp$. Conversely, if X is a surface as in Theorem 1, if $x \in k_X^\perp$, and if there exists a class $y \in k_X^\perp$ with $\langle x, y \rangle = 1$, then x can be represented by a differentiably embedded 2-sphere.*

Proof. Let $x \in H_2(X, \mathbf{Z})$ be a class with $q_X(x) = -2$. Suppose that x is represented by a differentiably embedded 2-sphere S . Let $j: S \rightarrow X$ be the embedding. The normal bundle N_S of S in X can be regarded as a $U(1)$ -bundle. Therefore the first Chern class $c_1(N_S)$ of the normal bundle is defined. Let $\xi \in H^2(X, \mathbf{Z})$ be the Poincaré dual of x . Then by [H, Theorem 4.8.1] $c_1(N_S) = j^*\xi$. If T_X and T_S denote the tangent bundles of X and S respectively, then we have $j^*\bar{k}_X = w_2(j^*T_X) = w_2(T_S) + w_2(N_S)$. This implies $\langle \bar{k}_X, \bar{\xi} \rangle = \overline{\chi(S)} + \bar{\xi}^2 = 0$ where $\chi(S)$ denotes the Euler characteristic of S . It follows that $\bar{x} \in \bar{k}_X^\perp$.

Conversely, let X be a surface as in Theorem 1. Then (L', Δ) is a complete vanishing lattice in the sense of [E1, Definition (2.2)]. This follows for complete intersection surfaces from [B, E3], for Moishezon surfaces from [M], and for Salvetti surfaces from [S] (see also [EO]). By [E1, Proposition (2.5)] we conclude that

$$\Delta = \{v \in L' \mid q_X(v) = -2 \quad \text{and} \quad \langle v, L' \rangle = \mathbf{Z}\}.$$

Therefore, if $x \in L'$ and if there exists a $y \in L'$ with $\langle x, y \rangle = 1$, then $x \in \Delta$, i.e. x is a vanishing cycle. But vanishing cycles are certainly represented by spheres. This proves Theorem 9.

Remark. We have even proved more, namely that every x satisfying the latter conditions of Theorem 9 is a vanishing cycle.

Our second application concerns a question which was posed by E. Brieskorn. First we show:

PROPOSITION 10. *Let X be an algebraic surface as in Theorem 1. Let $x \in \text{Hom}(L, \mathbf{Z})$. If $\{\pm x\}$ is invariant under $\psi(\text{Diff}_+(X))$, then $x = \lambda k_X$ for some $\lambda \in \mathbf{Q}$, unless $k_X^2 = 0, k_X \neq 0$.*

Proof. Let $U \subset L'_Q = L' \otimes \mathbf{Q}$ be a subspace of L'_Q which is invariant under $\psi(\text{Diff}_+(X))$. We show that either $U = L'_Q$ or U is contained in $(L'_Q)^\perp = \{v \in L'_Q \mid \langle v, w \rangle = 0 \text{ for all } w \in L'_Q\}$. Let $\Delta \subset L'$ be the set of vanishing cycles. If $\delta \in \Delta$ is not orthogonal to U then there exists a $y \in U$ with $\langle \delta, y \rangle \neq 0$. From $s_\delta(y) = y + \langle y, \delta \rangle \delta \in U$ it follows that $\delta \in U$. Since Γ_Δ acts transitively on Δ , we obtain $\Delta \subset U$. But Δ generates L' so that we must have $L'_Q = U$.

Now let $x \in \text{Hom}(L, \mathbf{Z})$ be invariant up to sign under $\Gamma_\Delta \subset \psi(\text{Diff}_+(X))$. If $k_X = 0$ then $L = L'$ and $(L')^\perp = \{0\}$. Hence it follows from what we have just shown that $x = 0$. If $k_X^2 \neq 0$ then we can write $x = \lambda k_X + x_0$ where $\lambda \in \mathbf{Q}$ and $x_0 \in \text{Hom}(L', \mathbf{Z}) \otimes \mathbf{Q}$. Now k_X is invariant under Γ_Δ , so we see that $\{\pm x_0\}$ is invariant under Γ_Δ . Since $(L')^\perp = \{0\}$, it follows that $x_0 = 0$. This proves Proposition 10.

Now $(\bar{L}', \bar{\Delta})$, where $\bar{\Delta} = \{\bar{\delta} \mid \delta \in \Delta\}$, is also a vanishing lattice. So we can derive the following proposition by the same arguments.

PROPOSITION 11. *Let X be an algebraic surface as in Theorem 1. Write $k_X = n_X \kappa_X$ for some primitive element $\kappa_X \in \text{Hom}(L, \mathbf{Z})$ and some non-negative integer n_X . If $x \in \text{Hom}(L, \mathbf{Z})$ is an element with $\overline{g(x)} = \bar{x}$ for all $g \in \psi(\text{Diff}_+(X))$, then $\bar{x} \in \{0, \overline{\kappa_X}\}$, unless $\kappa_X^2 = 0, \kappa_X \neq 0$.*

Consider two simply connected algebraic surfaces X and X' with corresponding lattices L_X and $L_{X'}$. Let $h: X \rightarrow X'$ be an orientation preserving homeomorphism between X and X' . Then $h_*: L_X \rightarrow L_{X'}$ is an isometry. For a subgroup $G \subset O(L_{X'})$ we define $G^{h*} := h_*^{-1} G h_*$. Note that we have $k_X^2 = k_{X'}^2$, since k_X^2 can be expressed in terms of the rank and signature of L_X .

THEOREM 12. *Let X, X' be algebraic surfaces as in Theorem 8, and suppose that $k_X^2 = k_{X'}^2 \neq 0$. Let $h: X \rightarrow X'$ be an orientation preserving homeomorphism. Then $\psi(\text{Diff}_+(X))$ and $\psi(\text{Diff}_+(X'))^{h*}$ are conjugate subgroups in $O(L_X)$ if and only if the divisibilities of k_X and $k_{X'}$ in integral cohomology are equal.*

Proof. Suppose that $\psi(\text{Diff}_+(X))$ and $\psi(\text{Diff}_+(X'))^{h*}$ are conjugate by $g \in O(L_X)$, i.e. $\psi(\text{Diff}_+(X)) = (\psi(\text{Diff}_+(X'))^{h*})^g$. Let $f \in \psi(\text{Diff}_+(X'))$. Then

$$(h_*g)^{-1}f(h_*g)(k_X) = \pm k_X,$$

since $\{\pm k_X\}$ is invariant under $\psi(\text{Diff}_+(X))$. Therefore

$$f((h_*g)(k_X)) = \pm (h_*g)(k_X).$$

This holds for every $f \in \psi(\text{Diff}_+(X'))$, so that $\{\pm (h_*g)(k_X)\}$ is invariant under $\psi(\text{Diff}_+(X'))$. We conclude from Proposition 10 that $(h_*g)(k_X) = \lambda k_X$, for some $\lambda \in \mathbf{Q}$. Now

$$k_{X'}^2 = k_X^2 = ((h_*g)(k_X))^2 = \lambda^2 k_{X'}^2,$$

hence $\lambda = \pm 1$, and it follows that the divisibilities of k_X and $k_{X'}$ in integral cohomology are equal.

Conversely, let $k_X = n_X \kappa_X$ and $k_{X'} = n_{X'} \kappa_{X'}$ for some primitive elements $\kappa_X \in H^2(X, \mathbf{Z})$ and $\kappa_{X'} \in H^2(X', \mathbf{Z})$ and some nonnegative integers n_X and $n_{X'}$ respectively, and assume that n_X and $n_{X'}$ are equal. Let λ_X and $\lambda_{X'}$ be the Poincaré duals of κ_X and $\kappa_{X'}$, and let K and K' be the one-dimensional sublattices of L_X spanned by λ_X and $h_*^{-1}(\lambda_{X'})$ respectively. Let $g: K \rightarrow K'$ be the homomorphism defined by $g(\lambda_X) = h_*^{-1}(\lambda_{X'})$. Then $\langle g(\lambda_X), g(\lambda_X) \rangle = \langle \lambda_X, \lambda_X \rangle$, hence g is an isometry. Now $b_2^+(X) \geq 3$, so that by a generalization of Witt's theorem [N, § 1.14, in particular 1.14.4, § 1.16] g can be extended to an isometry $g \in O(L_X)$. One can easily verify that $\psi(\text{Diff}_+(X))$ and $\psi(\text{Diff}_+(X'))^{h*}$ are conjugate by g . This proves Theorem 12.

COROLLARY 13. *Let X be an algebraic surface as in Theorem 8. If an element $h \in O'(L)$ normalizes $\psi(\text{Diff}_+(X))$, then h is contained in $\psi(\text{Diff}_+(X))$.*

Proof. This follows from Theorem 12, because

$$\psi(\text{Diff}_+(X)) = O'_k(L) \cdot \{\sigma_*, \text{id}\}.$$

Remark. Since -1 is not contained in $\psi(\text{Diff}_+(X))$ but in the normalizer $\text{Norm}(\psi(\text{Diff}_+(X)))$ of $\psi(\text{Diff}_+(X))$ we obtain from Corollary 13:

$$\text{Norm}(\psi(\text{Diff}_+(X))) / \psi(\text{Diff}_+(X)) \cong \mathbf{Z}/2 \cong \{\pm \text{id}\}.$$

Example. Let X be a complete intersection in \mathbf{P}^6 of multidegree $(7, 7, 5, 3)$. Let X' be the Salvetti surface $Y_4(10, 10, 6, 5; 5, 5, 3, 5)$. Both surfaces have $k^2 = 165375$, $p_g = 24499$, but the divisibilities of the canonical classes are 15 and 21 respectively. Therefore the surfaces are homeomorphic, but not diffeomorphic. Both surfaces satisfy the assumptions of Theorem 8.

We conclude from Theorem 12 that the subgroups of $O(L)$ corresponding to $\psi(\text{Diff}_+(X))$ and $\psi(\text{Diff}_+(X'))$ are not conjugate. This example was found by a computer search.

REFERENCES

- [BPV] BARTH, W., C. PETERS and A. VAN DE VEN. *Compact complex surfaces*. Berlin, Heidelberg, New York, Tokyo: Springer 1984.
- [B] BEAUVILLE, A. Le groupe de monodromie des familles universelles d'hyper-surfaces et d'intersections complètes. In: Grauert, H. (ed.) *Complex analysis and algebraic geometry*. Proc. Conf. Göttingen 1985 (*Lect. Notes Math. Vol. 1194*, pp. 8-18). Berlin, Heidelberg, New York: Springer 1986.
- [D] DONALDSON, S.K. Polynomial invariants for smooth four-manifolds. *Topology* 29 (1990), 257-315.
- [E1] EBELING, W. An arithmetic characterisation of the symmetric monodromy groups of singularities. *Invent. math.* 77 (1984), 85-99.
- [E2] —— The monodromy groups of isolated singularities of complete intersections. *Lect. Notes Math. Vol. 1293*. Berlin, Heidelberg, New York, Tokyo: Springer 1987.
- [E3] —— Vanishing lattices and monodromy groups of isolated complete intersection singularities. *Invent. math.* 90 (1987), 653-668.
- [E4] —— An example of two homeomorphic, nondiffeomorphic complete intersection surfaces. *Invent. math.* 99 (1990), 651-654.
- [EO] EBELING, W. and Ch. OKONEK. Donaldson invariants, monodromy groups, and singularities. *International Journal of Mathematics* 1 (1990), 233-250.
- [F] FREEDMAN, M. The topology of four-dimensional manifolds. *J. Diff. Geom.* 17 (1982), 357-454.
- [FMM] FRIEDMAN, R., B. MOISHEZON and J. W. MORGAN. On the C^∞ invariance of the canonical classes of certain algebraic surfaces. *Bull. Amer. Math. Soc. (N.S.)* 17 (1987), 283-286.
- [FM] FRIEDMAN, R. and J. W. MORGAN. On the diffeomorphism types of certain algebraic surfaces. I. *J. Diff. Geom.* 27 (1988), 297-369.
- [H] HIRZEBRUCH, F. *Topological methods in algebraic geometry*. Third enlarged edition. New York: Springer 1966.
- [HH] HIRZEBRUCH, F. und H. HOPF. Felder von Flächenelementen in 4-dimensionalen Mannigfaltigkeiten. *Math. Ann.* 136 (1958), 156-172.
- [Kr] KRECK, M. Isotopy classes of diffeomorphisms of $(k-1)$ -connected almost-parallelizable $2k$ -manifolds. In: Dupont, J. L., Madsen, I. H. (ed.) *Algebraic Topology Aarhus 1978* (*Lect. Notes Math., Vol. 763*, pp. 643-663). Berlin, Heidelberg, New York: Springer 1979.
- [M] MOISHEZON, B. Analogs of Lefschetz theorems for linear systems with isolated singularities. *J. Diff. Geom.* 31 (1990), 47-72.