

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 37 (1991)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AUTOMORPHIC SPECTRA ON THE TREE OF PGL_2
Autor: Efrat, Isaac
Kapitel: 1. The tree of $\mathrm{PGL}_2(K)$
DOI: <https://doi.org/10.5169/seals-58728>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)


Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. The above describes a decomposition $L^2(F) = R \oplus E$ (Theorem 5.1), made explicit in Theorem 5.3 (compare [L]). In particular, the spectrum of T on $L^2(F)$ is



1. THE TREE OF $PGL_2(K)$

The material in this section is adapted from Serre [S] and Weil [W]. An O -lattice in K^2 is a set

$$L = \{ \alpha v_1 + \beta v_2 \mid \alpha, \beta \in O \}$$

with v_1, v_2 a basis for K^2 . We can associate to L the matrix $(v_1, v_2) \in GL_2(K)$ and different choices of bases v_1, v_2 will give cosets in $GL_2(K)/GL_2(O)$. Two lattices L and L' are said to be equivalent if $L' = aL$ for some $a \in K^\times$. We thus have a natural correspondence between equivalence classes of O -lattices in K^2 and points in X .

We define a graph structure on X . Let Λ and Λ' be two equivalence classes of lattices. We say that Λ and Λ' are adjacent if there exist representatives $L \in \Lambda, L' \in \Lambda'$ such that

$$(1) \quad L' \subset L \quad \text{and} \quad L/L' \cong k.$$

THEOREM 1.1 ([S]). *The graph whose set of vertices is X and whose edges are the pairs (Λ, Λ') satisfying (1) is the (infinite) $(q+1)$ -regular tree.*

We seek a more explicit realization of X . Let B be the Borel subgroup of G_K consisting of the matrices whose bottom row is $(0, 1)$. Then the Iwasawa decomposition is

$$G_K = BG_O,$$

but it is not difficult to see that in fact any coset in X has a representative of the form

$$\begin{pmatrix} t^n & x \\ 0 & 1 \end{pmatrix}$$

with $x \in K$ and a uniquely determined $n \in \mathbf{Z}$.

In these coordinates we can write down the $q + 1$ vertices of X that are adjacent to a typical vertex $\begin{pmatrix} t^n & x \\ 0 & 1 \end{pmatrix}$. They are


$$\begin{pmatrix} t^{n+1} & x \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} t^{n-1} & \xi t^n + x \\ 0 & 1 \end{pmatrix}, \quad \xi \in k.$$

The group G_K acts on the tree X as a group of automorphisms. We can therefore define a graph structure on the quotient F for the action of Γ on X .

THEOREM 1.2 ([S], [W]). *The quotient graph $F = \Gamma \backslash X$ is given by (the cosets of)*

$$\begin{pmatrix} t^n & 0 \\ 0 & 1 \end{pmatrix} \quad n \geq 0,$$

so that F is the tree



In fact, the vertex $\begin{pmatrix} t^n & x \\ 0 & 1 \end{pmatrix}$ corresponds to n , and so if $n \geq 1$, its neighbor $\begin{pmatrix} t^{n+1} & x \\ 0 & 1 \end{pmatrix}$ corresponds to $n + 1$ while the other q neighbors are represented by $n - 1$. If $n = 0$, all neighbors correspond to 1.

2. THE OPERATOR T

Let μ be the Haar measure on G_K normalized so that $\mu(G_O) = q(q - 1)$. We compute the measure of F induced from μ . Since

$$F = \Gamma \backslash X = \Gamma \backslash G_K / G_O$$

we have

$$\Gamma \backslash G_K = \cup_{s \in F} sG_O,$$

where

$$sG_O = \{\Gamma su \mid u \in G_O\} \subset \Gamma \backslash G_K.$$

The point measure at s will be the measure of sG_O in the quotient space $\Gamma \backslash G_K$. Now we have a correspondence