

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 37 (1991)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AUTOMORPHIC SPECTRA ON THE TREE OF PGL_2
Autor: Efrat, Isaac
Kapitel: 3. Eigenfunctions
DOI: <https://doi.org/10.5169/seals-58728>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$\begin{aligned}
&= f(1)g(0) + \sum_{n=0}^{\infty} qf(n)g(n+1)q^{-(n+1)} + \sum_{n=2}^{\infty} f(n)g(n-1)q^{-(n-1)} \\
&= f(1)g(0) + f(0)g(1) + \sum_{n=1}^{\infty} f(n)g(n+1)q^{-n} \\
&\quad + q \sum_{n=1}^{\infty} f(n)g(n-1)q^{-n} - f(1)g(0) \\
&= f(0)g(1) + \sum_{n=0}^{\infty} (f(n)qg(n-1) + f(n)g(n+1))q^{-n} = \langle f, T\bar{g} \rangle .
\end{aligned}$$

3. EIGENFUNCTIONS

An automorphic eigenfunction of T on X with eigenvalue λ is a function on F that satisfies

$$\begin{aligned}
\lambda f(0) &= (q+1)f(1) , \\
\lambda f(n) &= qf(n-1) + f(n+1) , \quad n \geq 1 .
\end{aligned}$$

If we write $u(n) = \begin{pmatrix} f(n+1) \\ f(n) \end{pmatrix}$ and normalize $u(0) = \begin{pmatrix} \lambda \\ q+1 \end{pmatrix}$, we obtain the recursion

$$u(n) = A^n u(0)$$

with

$$A = \begin{pmatrix} \lambda & -q \\ 1 & 0 \end{pmatrix} .$$

Let $x_1, x_2 = \frac{1}{2}(\lambda \pm \sqrt{\lambda^2 - 4q})$ be the characteristic roots of A and assume that $x_1 \neq x_2$, i.e., that $\lambda \neq \pm 2\sqrt{q}$. Solving the recursion we get

PROPOSITION 3.1. *The eigenfunctions on F with eigenvalue λ are the multiples of the function*

$$(4) \quad f_{\lambda}(n) = \begin{cases} \frac{1}{x_1 - x_2} (\lambda(x_1^n - x_2^n) - q(q+1)(x_1^{n-1} - x_2^{n-1})) , & \text{if } n \geq 1 \\ q+1 & \text{if } n = 0 . \end{cases}$$

Example 3.2. If $\lambda = q + 1$ then $x_1 = q, x_2 = 1$ and

$$f_{q+1}(n) \equiv q + 1 ,$$

generating the space of constant functions. If $\lambda = -(q + 1)$, then

$$f_{-(q+1)}(n) = (-1)^n(q + 1) .$$

Example 3.3. When $\lambda = 2\sqrt{q}$ we can solve directly to get

$$f_{2\sqrt{q}}(n) = (q + 1 - (q - 1)n)q^{\frac{n}{2}},$$

and similarly

$$f_{-2\sqrt{q}}(n) = (-1)^n(q + 1 - (q - 1)n)q^{\frac{n}{2}} .$$

Remark 3.4. Since our tree is bipartite, we expect f_λ to be related to $f_{-\lambda}$ by a factor of $(-1)^n$ (compare [B, §8]). This can be seen from (4).

PROPOSITION 3.5. *The only eigenvalues λ with $|\lambda| > 2\sqrt{q}$ for which f_λ is in $L^2(F)$ are $\lambda = \pm(q + 1)$.*

Proof. Recalling (2) we see that if $f_\lambda \in L^2(F)$ then

$$f_\lambda(n) = o(q^{\frac{n}{2}}) \quad \text{as} \quad n \rightarrow \infty .$$

Now

$$(x_1 - x_2)f_\lambda(n) = x_1^{n-1}(\lambda x_1 - q(q + 1)) - x_2^{n-1}(\lambda x_2 - q(q + 1)) .$$

Assuming with no loss of generality that $|x_1| > \sqrt{q}, |x_2| = \frac{q}{|x_1|} < \sqrt{q}$, then $x_2^{n-1}(\lambda x_2 - q(q + 1)) = o(q^{\frac{n}{2}})$, so that we must have

$$\lambda x_1 - q(q + 1) = 0 ,$$

i.e., $\lambda = \pm(q + 1)$. Conversely, f_{q+1} and $f_{-(q+1)}$ are clearly in $L^2(F)$.

We turn our attention to λ with $|\lambda| < 2\sqrt{q}$. Then $x_2 = \bar{x}_1, |x_1| = \sqrt{q}$ and we let $x_1 = \sqrt{q}e^{i\theta}$. Then $\lambda = 2\sqrt{q}\cos\theta, 0 < \theta < \pi$. We renormalize and define

$$\tilde{f}_\theta = \frac{x_1 - x_2}{2\sqrt{q}} f_{2\sqrt{q}\cos\theta} .$$

Then, for $n \geq 1$,

$$(5) \quad \begin{aligned} \tilde{f}_\theta(n) &= q^{\frac{n}{2}} i((q+1)\sin\theta\cos(n\theta) - (q-1)\cos\theta\sin(n\theta)) \\ &= q^{\frac{n}{2}} i(\sin((n+1)\theta) - q\sin((n-1)\theta)), \end{aligned}$$

and

$$\tilde{f}_\theta(0) = (q+1)i\sin\theta.$$

PROPOSITION 3.6. *The functions \tilde{f}_θ , $0 < \theta < \pi$, are not in $L^2(F)$.*

Proof. It is sufficient to show that

$$(q+1)\sin\theta\cos(n\theta) - (q-1)\cos\theta\sin(n\theta) \not\rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This is the dot product of the two vectors

$$v_1 = ((q+1)\sin\theta, -(q-1)\cos\theta) \quad \text{and} \quad v_2 = (\cos(n\theta), \sin(n\theta)).$$

Since v_2 is not a constant function of n , we see that cosine of the angle between v_1 and v_2 is bounded away from 0 for arbitrarily large n .

Combining Propositions 3.5 and 3.6 we conclude

COROLLARY 3.7. *The discrete spectrum of T consists of the numbers $\pm(q+1)$, whose corresponding eigenfunctions (given in Example 3.1) span two one-dimensional eigenspaces of $L^2(F)$.*

Unlike the typical f_λ with $|\lambda| > 2\sqrt{q}$, those with $|\lambda| < 2\sqrt{q}$ satisfy

$$f_\lambda = O(q^{\frac{n}{2}}).$$

Our goal now is to show that these are approximate eigenfunctions that can be used to completely decompose $L^2(F)$.

4. CONTINUOUS SPECTRA

We wish to embed $L^2([0, \pi])$ with an appropriate measure into $L^2(F)$. To this end, let $\psi \in L^2([0, \pi])$ and $\tilde{f}_\theta(n)$ be extended as odd functions of $\theta \in [-\pi, \pi]$, and define

$$F_\psi(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \psi(\theta) \tilde{f}_\theta(n) d\theta.$$