Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 37 (1991)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AUTOMORPHIC SPECTRA ON THE TREE OF PGL_2
Autor: Efrat, Isaac
Kapitel: 4. CONTINUOUS SPECTRA
DOI: https://doi.org/10.5169/seals-58728

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 13.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Then, for $n \geqslant 1$,

$$\begin{aligned} \tilde{f}_\theta(n) &= q^{\frac{n}{2}} i((q+1)\sin\theta\cos(n\theta) - (q-1)\cos\theta\sin(n\theta)) \\ &= q^{\frac{n}{2}} i(\sin((n+1)\theta) - q\sin((n-1)\theta)) , \end{aligned} \tag{5}$$

and

$$\tilde{f}_\theta(0) = (q+1)\, i \sin\theta .$$

PROPOSITION 3.6. *The functions* $\tilde{f}_\theta, 0 < \theta < \pi$, *are not in* $L^2(F)$.

Proof. It is sufficient to show that

$$(q+1)\sin\theta\cos(n\theta) - (q-1)\cos\theta\sin(n\theta) \not\to 0 \quad \text{as} \quad n \to \infty .$$

This is the dot product of the two vectors

$$v_1 = ((q+1)\sin\theta, -(q-1)\cos\theta) \quad \text{and} \quad v_2 = (\cos(n\theta), \sin(n\theta)) .$$

Since v_2 is not a constant function of n, we see that cosine of the angle between v_1 and v_2 is bounded away from 0 for arbitrarily large n.

Combining Propositions 3.5 and 3.6 we conclude

COROLLARY 3.7. *The discrete spectrum of* T *consists of the numbers* $\pm(q+1)$, *whose corresponding eigenfunctions (given in Example 3.1) span two one-dimensional eigenspaces of* $L^2(F)$.

Unlike the typical f_λ with $|\lambda| > 2\sqrt{q}$, those with $|\lambda| < 2\sqrt{q}$ satisfy

$$f_\lambda = O(q^{\frac{n}{2}}) .$$

Our goal now is to show that these are approximate eigenfunctions that can be used to completely decompose $L^2(F)$.

4. CONTINUOUS SPECTRA

We wish to embed $L^2([0,\pi])$ with an appropriate measure into $L^2(F)$. To this end, let $\psi \in L^2([0,\pi])$ and $\tilde{f}_\theta(n)$ be extended as odd functions of $\theta \in [-\pi, \pi]$, and define

$$F_\psi(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \psi(\theta) \tilde{f}_\theta(n)\, d\theta .$$

THEOREM 4.1. *F_ψ is in $L^2(F)$ and we have the Plancherel formula*

$$< F_{\psi_1}, F_{\psi_2} > = < \psi_1, \psi_2 > , \tag{6}$$

where the inner product on the right is

$$\frac{1}{2\pi} \int_0^\pi \psi_1(\theta)\bar{\psi}_2(\theta)\left((q-1)^2 + 4q\sin^2\theta\right)d\theta .$$

Proof. We first note that

$$\frac{1}{2\pi} \int_\pi^\pi \psi(\theta)\left(i\sin((n+1)\theta)) - qi\sin((n-1)\theta)\right)d\theta = q\hat{\psi}(n-1) - \hat{\psi}(n+1) ,$$

where $\hat{\psi}(n)$ is the n-th Fourier coefficient of ψ. Therefore

$$< F_{\psi_1}, \bar{F}_{\psi_2} > = \frac{1}{q+1}(q+1)^2\hat{\psi}_1(1)\hat{\psi}_2(1)$$

$$+ \sum_{n=1}^{\infty} (q^2\hat{\psi}_1(n-1)\hat{\psi}_2(n-1) - q\hat{\psi}_1(n+1)\hat{\psi}_2(n-1)$$

$$- q\hat{\psi}_1(n-1)\hat{\psi}_2(n+1) + \hat{\psi}_1(n+1)\hat{\psi}_2(n+1))$$

$$= (q+1)\hat{\psi}_1(1)\hat{\psi}_2(1) + q^2\sum_{n=0}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n)$$

$$- q\left(\sum_{n=2}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n-2) + \sum_{n=0}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n+2)\right) + \sum_{n=2}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n) .$$

Now

$$\hat{\psi}(n-2) + \hat{\psi}(n+2) = 2\hat{\psi}(n) - 4(\psi\sin^2)^\wedge(n) .$$

Therefore we have

$$(q+1)\hat{\psi}_1(1)\hat{\psi}_2(1) + q^2\sum_{n=1}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n)$$

$$- q\left(2\sum_{n=1}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n) - 4\sum_{n=1}^{\infty} \hat{\psi}_1(n)(\psi_2\sin^2)^\wedge(n)\right)$$

$$- q\left(-\hat{\psi}_1(1)\hat{\psi}_2(-1) + \hat{\psi}_1(0)\hat{\psi}_2(2)\right) + \sum_{n=1}^{\infty} \hat{\psi}_1(n)\hat{\psi}_2(n) - \hat{\psi}_1(1)\hat{\psi}_2(1) .$$

Recalling Parseval's formula

$$\sum_{n=1}^{\infty} \hat{\psi}_1(n)\overline{\hat{\psi}_2(n)} = \frac{1}{2\pi}\int_0^\pi \psi_1(\theta)\hat{\psi}_2(\theta)d\theta$$

we get

$$-\frac{1}{2\pi}\int_0^{\pi}\psi_1(\theta)\psi_2(\theta)\left((q-1)^2+4q\sin^2(\theta)\right)d\theta\ ,$$

and since $\bar{F}_{\psi_2} = -F_{\bar{\psi}_2}$ we obtain (6).

5. Spectral decomposition

Let E be the space of functions F_ψ with $\psi \in L^2([0,\pi])$. It follows from §4 that E is a subspace of $L^2(F)$, invariant with respect to T. Further, let R be the two dimensional subspace generated by the discrete spectrum according to Corollary 3.7.

THEOREM 5.1. *We have a direct sum decomposition into invariant subspaces*

$$L^2(F) = R \oplus E\ .$$

Proof. The two spaces are easily seen to be orthogonal. We show that $E^\perp = R$. Let $g \in L^2(F)$ such that $< g, F_\psi > = 0$ for all ψ, i.e.,

$$0 = \frac{1}{q+1}g(0)\frac{1}{2\pi}\int_{-\pi}^{\pi} i(q+1)\sin\theta\psi(\theta)d\theta$$

$$+\sum_{n=1}^{\infty} g(n)\frac{1}{2\pi}\int_{-\pi}^{\pi}\psi(\theta)i(\sin((n+1)\theta)-q\sin((n-1)\theta))d\theta q^{-\frac{n}{2}}$$

$$= g(0)\hat{\psi}(1)+\sum_{n=1}^{\infty} g(n)(\hat{\psi}(n+1)-q\hat{\psi}(n-1))q^{-\frac{n}{2}}\ .$$

Therefore

$$g(0)\hat{\psi}(1)+\sum_{n=1}^{\infty} g(n)\hat{\psi}(n+1)q^{-\frac{n}{2}} = \sum_{n=0}^{\infty} g(n+1)\hat{\psi}(n)q^{-\frac{n-1}{2}}\ ,$$

or (as $\hat{\psi}(0) = 0$)

$$\sum_{n=1}^{\infty} g(n-1)\hat{\psi}(n)q^{-\frac{n-1}{2}} = \sum_{n=1}^{\infty} g(n+1)\hat{\psi}(n)q^{-\frac{n-1}{2}}\ .$$

Since $\psi \in L^2([0,\pi])$ and $g \in L^2(F)$, this can be viewed as an equality of inner products in the space l^2 of square integrable sequences. Now as ψ varies over