Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TOEPLITZ SEQUENCES, PAPERFOLDING, TOWERS OF HANOI

AND PROGRESSION-FREE SEQUENCES OF INTEGERS

Autor: Allouche, Jean-Paul / Bacher, Roland

Kapitel: 1. Toeplitz sequences

DOI: https://doi.org/10.5169/seals-59494

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 04.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

TOEPLITZ SEQUENCES, PAPERFOLDING, TOWERS OF HANOI AND PROGRESSION-FREE SEQUENCES OF INTEGERS

by Jean-Paul ALLOUCHE and Roland BACHER

ABSTRACT. What is the relationship between folding a piece of paper, moving disks in the classical tower of Hanoi algorithm and searching for minimal sequences of integers having no p terms in arithmetic progression? Our aim is to show how the Toeplitz sequences introduced by Jacobs and Keane in [15] allow us to give (inter alia) a unified description of the preceding problems. We give moreover some connections between Toeplitz sequences and q-automatic sequences.

1. Toeplitz sequences

In [15], (see also [21]), Jacobs and Keane defined the notion of Toeplitz sequence: they wanted to construct "explicit" sequences giving rise to strictly ergodic systems. They proved moreover that the unique invariant measure attached to such a sequence has a discrete rational spectrum. Roughly speaking a Toeplitz sequence is obtained by successive insertions of periodic sequences into the "holes" of a given periodic sequence, (a precise definition is given below). This construction was inspired by a device used by Toeplitz [28] for building explicitly almost periodic real functions. The method of Jacobs and Keane has since been used by many people working in ergodic theory (see for instance [29], [16] and [25], see also [14] and its impressive bibliography). We now give the definition of a Toeplitz sequence (compare with [15], [16], [14] and [29]):

Let $\Gamma = \{a_1, \dots, a_r, \omega\}$ be an alphabet (finite set) with a "marked" letter ("hole") ω . If $B = (B(k))_{k \ge 0}$ is a sequence with values in Γ , we define a transformation $T_B : \Gamma^N \to \Gamma^N$ as follows: for any sequence $C = (C(k))_{k \ge 0}$ with values in Γ , let $h_0 < h_1 < \cdots$ be the increasing sequence (which might

be finite or even empty) of those integers h for which $C(h) = \omega$. Then one defines

$$T_BC(j) = C(j)$$
 if $C(j) \neq \omega$,
 $T_BC(h_k) = B(k)$ for every k .

Suppose we are now given a sequence of periodic sequences B_0 , B_1 , \cdots , B_k , \cdots with values in Γ , and such that the zeroth value of each B_j is not equal to ω . Writing T_j instead of T_{B_j} , we then define a sequence of periodic sequences as follows:

$$A_0 = B_0$$
 $A_1 = T_1(A_0)$
 $A_2 = T_2(A_1) = T_2(T_1(A_0))$
...
 $A_{k+1} = T_{k+1}(A_k) = T_{k+1}(T_k(\cdots(T_1(A_0))\cdots))$.

As k goes to infinity the sequence A_k tends to a limit A with values in $\Gamma - \{\omega\}$ (the existence of this limit, for the topology of simple convergence, is left to the reader): such a sequence is called a *Toeplitz sequence*. An alternative (equivalent) definition of a Toeplitz sequence is given in [29]:

A is a Toeplitz sequence if and only if one has

$$\forall n \in \mathbb{N} \quad \exists p \in \mathbb{N}^* \quad \forall n' \equiv n \pmod{p} \quad A(n') = A(n)$$
.

In what follows we first suppose that the set Γ is not necessarily a finite set; second, we restrict ourselves to the case where the sequence B_0, B_1, \cdots has the following form: there exist a periodic sequence B with values in Γ such that $B(0) \neq \omega$ and a function f from Γ to Γ with $f^{-1}(\omega) = {\omega}$, such that

$$\forall k \geqslant 0 \quad B_k = f^{(k)}(B) ,$$

where $f^{(k)}$ is the k^{th} iterate of the function f and $f^{(k)}(B)$ is the termwise image of the sequence B under $f^{(k)}$; the resulting Toeplitz sequence

$$A = \lim_{k \to \infty} T_k \big(\cdots T_2 \big(T_1(B) \big) \cdots \big)$$

(where
$$T_k = T_{B_k} = T_{f^{(k)}(B)}$$
)

will be called the *Toeplitz transform* of (B, f) and denoted by Tt(B, f).

Example. Let B be the sequence $(0\omega 1\omega)^{\infty}$, and let f be defined on $\{0, 1, \omega\}$ by f(0) = 1, f(1) = 0, $f(\omega) = \omega$, then one has: $B_0 = (0\omega 1\omega 0\omega 1\omega \cdots) (= B),$ $B_1 = (1\omega 0\omega 1\omega 0\omega \cdots),$ $B_2 = B_0,$ \cdots $A_0 = (0\omega 1\omega 0\omega 1\omega \cdots),$ $A_1 = (011\omega 001\omega \cdots),$ $A_2 = (0110001\omega \cdots),$

Note that if Γ is finite, and f one-to-one, such a sequence Tt(B, f) can also be obtained by replacing B by a sequence of greater period and f by id. We now give four examples of Toeplitz transforms in (apparently) unrelated domains.

2. Paperfolding sequences and Toeplitz transforms

In [23] and [22] Prodinger and Urbanek study the Toeplitz transform of $((0\omega 1\omega)^{\infty}, id)$ and of $((0\omega 1\omega 1\omega 0\omega)^{\infty}, id)$. They prove that these sequences do not have arbitrarily long squares (a sequence A contains a square of length 2k if there exists an index j such that A(j+n) = A(j+n+k) for every n between 0 and k-1). Dekking already noticed in [10] that the first sequence is nothing but the regular paperfolding sequence (see [9], [18], [20], [17]), which is obtained by repeatedly folding a piece of paper, and we obtained in [1] the same result as Prodinger and Urbanek for the general paperfolding sequences. Let us give here two simple examples:

PROPOSITION. Let B be the sequence $B = (0\omega 1\omega)^{\infty}$ and let f be defined by f(0) = 1, f(1) = 0 and $f(\omega) = \omega$. Then

the sequence Tt(B, id) is the regular paperfolding sequence, the sequence Tt(B, f) is the alternate paperfolding sequence.