Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TOEPLITZ SEQUENCES, PAPERFOLDING, TOWERS OF HANOI AND PROGRESSION-FREE SEQUENCES OF INTEGERS

Autor: Allouche, Jean-Paul / Bacher, Roland

Kapitel: 2. Paperfolding sequences and Toeplitz transforms

DOI: https://doi.org/10.5169/seals-59494

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Example. Let B be the sequence $(0\omega 1\omega)^\infty$, and let f be defined on $\{0, 1, \omega\}$ by $f(0) = 1$, $f(1) = 0$, $f(\omega) = \omega$, then one has:

$$\begin{aligned}
B_0 &= (0\omega 1\omega 0\omega 1\omega \cdots)\ (= B)\ ,\\
B_1 &= (1\omega 0\omega 1\omega 0\omega \cdots)\ ,\\
B_2 &= B_0\ ,\\
&\cdots\\
A_0 &= (0\omega 1\omega 0\omega 1\omega \cdots)\ ,\\
A_1 &= (011\omega 001\omega \cdots)\ ,\\
A_2 &= (0110001\omega \cdots)\ ,\\
&\cdots .
\end{aligned}$$

Note that if Γ is finite, and f one-to-one, such a sequence $Tt(B,f)$ can also be obtained by replacing B by a sequence of greater period and f by id.

We now give four examples of Toeplitz transforms in (apparently) unrelated domains.

2. Paperfolding sequences and Toeplitz transforms

In [23] and [22] Prodinger and Urbanek study the Toeplitz transform of $((0\omega 1\omega)^\infty, id)$ and of $((0\omega 1\omega 1\omega 0\omega)^\infty, id)$. They prove that these sequences do not have arbitrarily long squares (a sequence A contains a square of length $2k$ if there exists an index j such that $A(j+n) = A(j+n+k)$ for every n between 0 and $k-1$). Dekking already noticed in [10] that the first sequence is nothing but the regular paperfolding sequence (see [9], [18], [20], [17]), which is obtained by repeatedly folding a piece of paper, and we obtained in [1] the same result as Prodinger and Urbanek for the general paperfolding sequences. Let us give here two simple examples:

Proposition. *Let B be the sequence $B = (0\omega 1\omega)^\infty$ and let f be defined by $f(0) = 1$, $f(1) = 0$ and $f(\omega) = \omega$. Then*

the sequence $Tt(B, id)$ is the regular paperfolding sequence,

the sequence $Tt(B, f)$ is the alternate paperfolding sequence.

Proof. It follows from instance from [18] (after replacing 1's by 0's and -1's by 1's) that the regular paperfolding sequence R and the alternate paperfolding sequence A are given by

$$R(2^k(2m+1)-1) = \frac{(1-(-1)^m)}{2} \quad \forall k, m \geqslant 0 .$$

$$A(2^k(2m+1)-1) = \frac{(1-(-1)^{k+m})}{2} \quad \forall k, m \geqslant 0 .$$

Let U and V be the sequences defined by

$$U = Tt((0\omega 1\omega)^\infty, id) ,$$
$$V = Tt((0\omega 1\omega)^\infty, f) .$$

A straightforward computation gives

$$U(2n) = \frac{(1-(-1)^n)}{2} \quad \forall n \geqslant 0 ,$$

$$U(2n+1) = U(n) \quad \forall n \geqslant 0 .$$

Hence

$$U(2^k(2m+1)-1) = U(2(2^{k-1}(2m+1)-1)+1)$$
$$= U(2^{k-1}(2m+1)-1) = \cdots = U(2m) = \frac{(1-(-1)^m)}{2} .$$

This proves that $U = R$.

In the same way one has

$$V(2n) = \frac{(1-(-1)^n)}{2} ,$$

$$V(2n+1) = 1 - V(n) .$$

Hence

$$V(2^k(2m+1)-1) = V(2(2^{k-1}(2m+1)-1)+1) = 1 - V(2^{k-1}(2m+1)-1)$$

$$= \cdots = \left\{ \begin{array}{ll} V(2m) & \text{if } k \text{ is even,} \\ 1 - V(2m) & \text{if } k \text{ is odd,} \end{array} \right\} = \frac{(1-(-1)^{k+m})}{2} ,$$

and finally $V = A$.