Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: RATIONALITY OF PIECEWISE LINEAR FOLIATIONS AND HOMOLOGY OF THE GROUP OF PIECEWISE LINEAR HOMEOMORPHISMS

Autor: Tsuboi, Takashi

Kapitel: §2. Discontinuous invariants

DOI: https://doi.org/10.5169/seals-59495

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 17.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$Y \times Z \to \Omega X \times \Omega X \xrightarrow{*} \Omega X .$$

Then this represents $u * v \in H_{p+q}(\Omega X; \mathbf{Z})$. On the other hand, the composition

$$Y \times Z \to \Omega X \times \Omega X \to PX \times \Omega X \to X \times \Omega X$$

bounds $SY \times Z \to PX \times \Omega X \to X \times \Omega X$, which represents $s(u) \otimes v$. Hence $s(u) \otimes v$ and $u * v$ are related under ∂^{p+1}.

§2. Discontinuous invariants

First we review the definition by Morita ([10]) of discontinuous invariants arising from the Godbillon-Vey invariant for codimension one foliations.

Let $\mathscr{F}$ be a codimension one foliation of a closed oriented $3k$-dimensional manifold M. Then the Godbillon-Vey class $gv(\mathscr{F}) \in H^3(M; \mathbf{R})$ is defined ([6]). Let $\{x_1, \ldots, x_n\}$ be a basis of $H^3(M; \mathbf{Q})$. Then $gv(\mathscr{F})$ is written as

$$gv(\mathscr{F}) = a_1 x_1 + \ldots + a_n x_n ,$$

where $a_1, \ldots, a_n \in \mathbf{R}$. The discontinuous invariant GV_k is defined by

$$GV_k(\mathscr{F}) = \sum_{i_1 < \ldots < i_k} (x_{i_1} \cup \ldots \cup x_{i_k})[M]\ a_{i_1} \wedge_{\mathbf{Q}} \ldots \wedge_{\mathbf{Q}} a_{i_k} \in \mathbf{R}^{\wedge k} = \overbrace{\mathbf{R} \wedge_{\mathbf{Q}} \ldots \wedge_{\mathbf{Q}} \mathbf{R}}^{k} ,$$

where $[M] \in H_{3k}(M; \mathbf{Z})$ is the fundamental class. Morita showed that GV_k is natural, GV_k depends only on the foliated cobordism class of $\mathscr{F}$, and hence there is a universal map $GV_k \colon H_{3k}(B\Gamma_1; \mathbf{Z}) \to \mathbf{R}^{\wedge k}$ ([10]).

The same argument applies to transversely piecewise linear foliations and the discrete Godbillon-Vey class defined in [5] and [3]. Then the following theorem is obtained from the description by Greenberg ([7]) of the classifying space for them and Lemma (1.1).

THEOREM (2.1). *Let $\mathscr{F}$ be a codimension one transversely orientable transversely piecewise linear foliation of a closed oriented $3k$-dimensional manifold $M (k \geqslant 2)$. Then $GV_k(\mathscr{F}) = 0$.*

Proof. The weak homotopy type of the classifying space $B\bar{\Gamma}_1^{PL}$ for codimension one transversely oriented transversely piecewise linear foliations is known by Greenberg ([7]). This classifying space $B\bar{\Gamma}_1^{PL}$ has the weak homotopy type of the join $B\mathbf{R}^\delta * B\mathbf{R}^\delta$ of two copies of $B\mathbf{R}^\delta = K(\mathbf{R}, 1)$. Let

gv denote the discrete Godbillon-Vey class defined as a 3-dimensional cohomology class of this classifying space ([5], [3]).

$$gv \in H^3(B\bar{\Gamma}_1^{PL}; \mathbf{R}) .$$

By Lemma (1.1), the higher discontinuous invariants GV_k are trivial in this classifying space $B\bar{\Gamma}_1^{PL}$. Hence by the naturality of GV_k, $GV_k(\mathscr{F}) = 0$.

COROLLARY (2.2). *Let $\mathscr{F}$ be a codimension one transversely piecewise linear foliation of $S^3 \times S^3$. $GV(\mathscr{F}) = (a, b) \in H^3(S^3 \times S^3, \mathbf{R})$ satisfies $a/b \in \mathbf{Q} \cup \{\infty\}$.*

Proof. $0 = GV_2(\mathscr{F}) = a \wedge_{\mathbf{Q}} b$. Hence $a/b \in \mathbf{Q} \cup \{\infty\}$.

Remark. Morita translated the question of rationality into that of graded commutativity of $*$-product defined on the homology of the group of diffeomorphisms of $\mathbf{R}$ with compact support ([10]). In the later sections, we calculate the homology of the group $PL_c(\mathbf{R})$ of piecewise linear homeomorphisms of $\mathbf{R}$ with compact support as well as the $*$-product structure. We see that the $*$-product is certainly not graded commutative, which insures the rationality. The argument on the rationality of transversely piecewise linear foliations uses the fact that the Godbillon-Vey invariant localizes on transversely discrete sets and this argument cannot be generalized for smooth foliations for the moment. See how the class $C^4_{(1,1,1,1)}$ exists in §3. We also see that the Whitehead product of elements of $\pi_n(B\bar{\Gamma}_1^{PL})$ which are not zero in homology is usually nontrivial and has infinite order.

Remark. The Hurewicz map

$$\pi_n(B\bar{\Gamma}_1^{PL}) \to H_n(B\bar{\Gamma}_1^{PL}; \mathbf{Z})$$

is surjective. To see this, note first that by Greenberg ([7]),

$$H_n(B\bar{\Gamma}_1^{PL}; \mathbf{Z}) \cong \sum_{i=1}^{n-1} \mathbf{R}^{\wedge i} \otimes_{\mathbf{Q}} \mathbf{R}^{\wedge n-1-i} .$$

An element $(a_1 \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} a_i) \otimes_{\mathbf{Q}} (b_{i+1} \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} b_{n-1}) \in \mathbf{R}^{\wedge i} \otimes_{\mathbf{Q}} \mathbf{R}^{\wedge n-1-i}$ is represented by the following foliation of $T^i * T^{n-1-i}$. Consider the foliated $\mathbf{R}$-product with noncompact support over T^{n-1} such that the holonomy $h: \pi_1(T^{n-1}) \to PL(\mathbf{R})$ is given by

$$h(e_j)(x) = e^{a_i}x \text{ for } x < 0 \text{ and } h(e_j)(x) = x \text{ for } x > 0 \text{ if } j = 1, \dots, i$$
$$h(e_j)(x) = x \text{ for } x < 0 \text{ and } h(e_j)(x) = e^{b_j}x \text{ for } x > 0 \text{ if } j = i+1, \dots, n-1 .$$

This foliation restricted to $T^{n-1} \times [-1, 1]$ induces a foliation of $T^i * T^{n-1-i}$ which is

$$T^{n-1} \times [-1, 1]/(T^i \times T^{n-1-i} \times \{-1\} \sim T^i \times \{-1\}, \\ T^i \times T^{n-1-i} \times \{1\} \sim T^{n-1-i} \times \{1\}) .$$

Note that there is a degree one map from the suspension of T^{n-1} to $T^i * T^{n-1-i}$. Since we can embed $T^{n-1} \times [-1, 1]$ in S^n, we have a degree one map from S^n to the suspension of T^{n-1}, hence to $T^i * T^{n-1-i}$. Thus Hurewicz map is surjective.

§3. Homology of the group of piecewise linear homeomorphisms

Let $PL_c(\mathbf{R})$ denote the group of piecewise linear homeomorphisms of $\mathbf{R}$ with compact support. Let $\mu: PL_c(\mathbf{R}) \times PL_c(\mathbf{R}) \to PL_c(\mathbf{R})$ be the composition of two isomorphisms $PL_c(\mathbf{R}) \cong PL_c((-\infty, 0))$ and $PL_c(\mathbf{R}) \cong PL_c((0, \infty))$, and the inclusion

$$PL_c((-\infty, 0)) \times PL_c((0, \infty)) \to PL_c(\mathbf{R}) .$$

Then μ induces a product $*$ on the homology of $BPL_c(\mathbf{R})^\delta$ ([10]).

The homology of the group $PL_c(\mathbf{R})$ of piecewise linear homeomorphisms of $\mathbf{R}$ with compact support is described as follows. For positive integers i and j, put

$$V^{i,j} = \mathbf{R}^{\wedge i} \otimes_{\mathbf{Q}} \mathbf{R}^{\wedge j} \\ = \overbrace{(\mathbf{R} \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} \mathbf{R})}^{i} \otimes_{\mathbf{Q}} \overbrace{(\mathbf{R} \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} \mathbf{R})}^{j} .$$

THEOREM (3.1).

$$H_m(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \cong \sum V^{k_1^-, k_1^+} \otimes_{\mathbf{Q}} \dots \otimes_{\mathbf{Q}} V^{k_s^-, k_s^+} ,$$

where the sum is taken over even number of positive integers

$$(k_1^-, k_1^+, \dots, k_s^-, k_s^+)$$

such that $k_1^- + k_1^+ + \dots + k_s^- + k_s^+ = m$. *Moreover, the $*$-product*

$$*: H_i(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \times H_j(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \to H_{i+j}(BPL_c(\mathbf{R})^\delta; \mathbf{Z})$$

coincides with the tensor product.