

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 38 (1992)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: RATIONALITY OF PIECEWISE LINEAR FOLIATIONS AND
HOMOLOGY OF THE GROUP OF PIECEWISE LINEAR
HOMEOMORPHISMS
Autor: Tsuboi, Takashi
Kapitel: §3. Homology of the group of piecewise linear homeomorphisms
DOI: <https://doi.org/10.5169/seals-59495>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

This foliation restricted to $T^{n-1} \times [-1, 1]$ induces a foliation of $T^i * T^{n-1-i}$ which is

$$T^{n-1} \times [-1, 1] / (T^i \times T^{n-1-i} \times \{-1\} \sim T^i \times \{-1\}, \\ T^i \times T^{n-1-i} \times \{1\} \sim T^{n-1-i} \times \{1\}).$$

Note that there is a degree one map from the suspension of T^{n-1} to $T^i * T^{n-1-i}$. Since we can embed $T^{n-1} \times [-1, 1]$ in S^n , we have a degree one map from S^n to the suspension of T^{n-1} , hence to $T^i * T^{n-1-i}$. Thus Hurewicz map is surjective.

§3. HOMOLOGY OF THE GROUP OF PIECEWISE LINEAR HOMEOMORPHISMS

Let $PL_c(\mathbf{R})$ denote the group of piecewise linear homeomorphisms of \mathbf{R} with compact support. Let $\mu: PL_c(\mathbf{R}) \times PL_c(\mathbf{R}) \rightarrow PL_c(\mathbf{R})$ be the composition of two isomorphisms $PL_c(\mathbf{R}) \cong PL_c((-\infty, 0))$ and $PL_c(\mathbf{R}) \cong PL_c((0, \infty))$, and the inclusion

$$PL_c((-\infty, 0)) \times PL_c((0, \infty)) \rightarrow PL_c(\mathbf{R}).$$

Then μ induces a product $*$ on the homology of $BPL_c(\mathbf{R})^\delta$ ([10]).

The homology of the group $PL_c(\mathbf{R})$ of piecewise linear homeomorphisms of \mathbf{R} with compact support is described as follows. For positive integers i and j , put

$$V^{i,j} = \mathbf{R}^{\wedge i} \otimes_{\mathbf{Q}} \mathbf{R}^{\wedge j} \\ = \underbrace{(\mathbf{R} \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} \mathbf{R})}_i \otimes_{\mathbf{Q}} \underbrace{(\mathbf{R} \wedge_{\mathbf{Q}} \dots \wedge_{\mathbf{Q}} \mathbf{R})}_j.$$

THEOREM (3.1).

$$H_m(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \cong \sum V^{k_1^-, k_1^+} \otimes_{\mathbf{Q}} \dots \otimes_{\mathbf{Q}} V^{k_s^-, k_s^+},$$

where the sum is taken over even number of positive integers

$$(k_1^-, k_1^+, \dots, k_s^-, k_s^+)$$

such that $k_1^- + k_1^+ + \dots + k_s^- + k_s^+ = m$. Moreover, the $*$ -product

$$*: H_i(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \times H_j(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \rightarrow H_{i+j}(BPL_c(\mathbf{R})^\delta; \mathbf{Z})$$

coincides with the tensor product.

For small dimensions, this theorem says that

$$\begin{aligned} H_1(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) &\cong 0, \\ H_2(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) &\cong \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R}, \\ H_3(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) &\cong (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}) \otimes_{\mathbf{Q}} \mathbf{R} \oplus \mathbf{R} \otimes_{\mathbf{Q}} (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}), \text{ and} \\ H_4(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) &\cong (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}) \otimes_{\mathbf{Q}} \mathbf{R} \oplus (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}) \otimes_{\mathbf{Q}} (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}) \\ &\quad \oplus \mathbf{R} \otimes_{\mathbf{Q}} (\mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R} \wedge_{\mathbf{Q}} \mathbf{R}) \oplus \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R}. \end{aligned}$$

The first homology group is 0 is equivalent to that $PL_c(\mathbf{R})$ is perfect and this is due to Epstein ([1]). The second and third homologies are given explicitly by Greenberg ([7]). The summand $\mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R} \otimes_{\mathbf{Q}} \mathbf{R}$ of $H_4(BPL_c(\mathbf{R})^\delta; \mathbf{Z})$ is the image of the $*$ -product on $H_2(BPL_c(\mathbf{R})^\delta; \mathbf{Z})$ and since the $*$ -product coincides with the tensor product, the $*$ -product is not graded commutative. This implies that the Whitehead product

$$\pi_3(B\bar{\Gamma}_1^{PL}) \times \pi_3(B\bar{\Gamma}_1^{PL}) \rightarrow \pi_5(B\bar{\Gamma}_1^{PL})$$

is highly nontrivial and this is the obstruction to construct a foliation on $S^3 \times S^3$ with given Godbillon-Vey class. In this way, as in mentioned in §2, this is related to the rationality (see [10]).

Theorem (3.1) is also obtained as an application of the description by Greenberg ([7]) of the classifying space $B\bar{\Gamma}_1^{PL}$. As we mentioned, his result says that this classifying space is weakly homotopy equivalent to the join $B\mathbf{R}^\delta * B\mathbf{R}^\delta$. To show Theorem (3.1), we use the isomorphism

$$H_*(BPL_c(\mathbf{R})^\delta; \mathbf{Z}) \cong H_*(\Omega B\bar{\Gamma}_1^{PL}; \mathbf{Z})$$

due to Mather ([9]) adapted for the PL case by Ghys-Sergiescu ([5]) and Greenberg ([7]) using a result of Segal ([11]), and the homology spectral sequence associated to the fibration

$$\Omega B\bar{\Gamma}_1^{PL} \rightarrow PB\bar{\Gamma}_1^{PL} \rightarrow B\bar{\Gamma}_1^{PL}.$$

Since $B\bar{\Gamma}_1^{PL}$ is simply connected, the E^2 term of this spectral sequence is as follows.

$$E_{p+1,q}^2 = H_{p+1}(B\bar{\Gamma}_1^{PL}; \mathbf{Z}) \otimes_{\mathbf{Q}} H_q(\Omega B\bar{\Gamma}_1^{PL}; \mathbf{Z}).$$

Note that $H_*(B\bar{\Gamma}_1^{PL}; \mathbf{Z})$ is torsion free. From this, Greenberg obtained the second and the third homologies ([7]). To show our theorem, we show that, for $p \geq 0$,

$$\begin{aligned} E_{p+1,q}^2 &= Z_{p+1,q}^2 = \dots = Z_{p+1,q}^p \quad \text{and} \\ Z_{p+1,q}^{p+1} &= Z_{p+1,q}^\infty = 0 = B_{p+1,q}^\infty = \dots = B_{p+1,q}^2. \end{aligned}$$

This is equivalent to that the differentials induce an isomorphism

$$\sum_{p+q=m, p \geq 0} H_{p+1}(B\bar{\Gamma}_1^{PL}; \mathbf{Z}) \otimes_{\mathbf{Q}} H_q(\Omega B\bar{\Gamma}_1^{PL}; \mathbf{Z}) \rightarrow H_{p+q}(\Omega B\bar{\Gamma}_1^{PL}; \mathbf{Z}).$$

To show this we define the cohomology classes of $BPL_c(\mathbf{R})^\delta$ which detect the images of generators of $H_{p+1}(B\bar{\Gamma}_1^{PL}; \mathbf{Z}) \otimes_{\mathbf{Q}} H_q(\Omega B\bar{\Gamma}_1^{PL}; \mathbf{Z})$.

§4. CONSTRUCTION OF COCYCLES OF THE GROUP $PL_c(\mathbf{R})$

Tensor determinants. We define a determinant of an $(n \times n)$ real matrix which takes values in the tensor product over \mathbf{Q} of n copies of \mathbf{R} . For $(a_{ij})_{i,j=1,\dots,n}$, we put

$$\det^{\otimes_{\mathbf{Q}}}(a_{ij}) = \sum_{\sigma} \text{sign}(\sigma) a_{\sigma(1)1} \otimes_{\mathbf{Q}} \dots \otimes_{\mathbf{Q}} a_{\sigma(n)n}.$$

For example,

$$\det^{\otimes_{\mathbf{Q}}}\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = a_{11} \otimes_{\mathbf{Q}} a_{22} - a_{21} \otimes_{\mathbf{Q}} a_{12}.$$

We have the usual multilinearity but we do not have the usual alternativity. For example,

$$\det^{\otimes_{\mathbf{Q}}}\begin{pmatrix} a & b \\ a & b \end{pmatrix} = 0 \quad \text{but} \quad \det^{\otimes_{\mathbf{Q}}}\begin{pmatrix} a & a \\ b & b \end{pmatrix} = a \otimes_{\mathbf{Q}} b - b \otimes_{\mathbf{Q}} a = a \wedge_{\mathbf{Q}} b.$$

The latter is not necessarily zero. In general, if we change the rows then this determinant changes sign, however, there are no simple laws for changing columns. It is worth noticing that we have the usual formula of developing with respect to the first or the last column.

$$\begin{aligned} \det^{\otimes_{\mathbf{Q}}}(a_{ij}) &= \sum_{i+1}^n (-1)^{i+1} a_{i1} \otimes_{\mathbf{Q}} \det^{\otimes_{\mathbf{Q}}}(A_{i1}) \\ &= \sum_{i+1}^n (-1)^{i+n} \det^{\otimes_{\mathbf{Q}}}(A_{in}) \otimes_{\mathbf{Q}} a_{in}, \end{aligned}$$

where A_{ij} is the matrix (a_{ij}) with the i -th row and the j -th column deleted.

Cocycles of Lipschitz homeomorphism groups. We review the construction of cocycles of certain Lipschitz homeomorphism groups of the real line or the circle (see [13]). Let \mathcal{S} be the space of functions with compact support