

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 38 (1992)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BARKER SEQUENCES AND DIFFERENCE SETS
Autor: Eliahou, Shalom / Kervaire, Michel
Kapitel: 3. Barker sequences
DOI: <https://doi.org/10.5169/seals-59496>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

In the fourth column of Table II, we have indicated the known existing cyclic difference sets or the relevant prime power exhibiting non-existence by the semi-primitivity theorem of Section 1. The values of the parameter n left out by these two classes are $n = 7, 25, 28, 37, 43, 44, 49, 52, 61, 67, 72, 75, 76, 86, 97, 99$ and 100 . We have reached a non-existence conclusion in these cases by using the multiplier theorem of Section 1. The required calculations being quite lengthy, it is impossible to expose them all. Instead, Section 4 contains some typical examples of application of this theorem.

3. BARKER SEQUENCES

Recall that a Barker sequence is a binary sequence $A = (a_1, \dots, a_l)$ such that the aperiodic correlations $c_j(A) = \sum_{i=1}^{l-j} a_i a_{i+j}$ belong to $\{-1, 0, 1\}$ for all $j = 1, \dots, l-1$.

The set of Barker sequences of a given length is preserved by the following transformations:

$$A \mapsto \alpha A, \text{ where } (\alpha A)_i = -a_i$$

$$A \mapsto \beta A, \text{ where } (\beta A)_i = (-1)^i a_i$$

$$A \mapsto \gamma A, \text{ where } (\gamma A)_i = a_{l-i+1},$$

with $l = \text{length}(A)$.

The group of transformations of Barker sequences generated by α, β and γ is the elementary abelian 2-group $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/2\mathbf{Z}$ of rank 3 if l is odd, and is the non-abelian dihedral 2-group of order 8 with presentation

$$D_8 = \langle \alpha, \beta, \gamma : \alpha^2 = \beta^2 = \gamma^2 = 1, \alpha\beta = \beta\alpha, \alpha\gamma = \gamma\alpha, \gamma\beta\gamma = \alpha\beta \rangle$$

for l even. Note that in this case, D_8 is also generated by $\rho = \beta\gamma$ and γ with presentation

$$D_8 = \langle \rho, \gamma : \rho^4 = \gamma^2 = 1, \gamma\rho\gamma = \rho^{-1} \rangle.$$

Case of odd length. The complete list of Barker sequences of odd length was established by R. Turyn and J. Storer, [ST] and reads as follows (in lengths ≥ 3):

$$(1, 1, -1)$$

$$(1, 1, 1, -1, 1)$$

$$(1, 1, 1, -1, -1, 1, -1)$$

$$(1, 1, 1, -1, -1, -1, 1, -1, -1, 1, -1)$$

$$(1, 1, 1, 1, 1, -1, -1, 1, 1, -1, 1, -1, 1).$$

The list is complete up to the transformations α , β and γ given above. The orbit of each Barker sequence in the above Turyn-Storer list under this transformation group consists of 4 sequences.

Case of even length. The situation here is completely different. The only known examples are

$$(1, 1) \quad \text{and} \quad (1, 1, 1, -1),$$

again up to modifications by the above transformations α , β and γ . Note that the sequence $(1, 1, 1, -1)$ gives rise to 8 sequences under this transformation group.

It is widely believed that these are the only Barker sequences of even length. We will show that this is true up to length 1 898 884.

We know from Section 1 that a Barker sequence of even length (≥ 4) is also a periodic Barker sequence with correlation $\gamma = 0$, and we know from Section 2 that the length l must be of the form $l = 4N^2$ with N odd, if $l \geq 4$. We also know from Section 2 that if N is an odd integer with a prime factor p such that p is self-conjugate modulo N , then there is no (periodic) Barker sequence of length $4N^2$. In other words, N is excluded if, for p as above, there is some positive integer f such that $p^f \equiv -1 \pmod{N'}$, where N' is the largest divisor of N which is relatively prime to p . An immediate consequence is that N cannot be a prime or a prime power. R. Turyn used the above theorem to show that, if there exists a (periodic) Barker sequence of length $l = 4N^2$ with $N > 1$, then necessarily $N \geq 55$. With the following result of [EKS], this bound can be improved to $N \geq 689$, but only for true (i.e. aperiodic) Barker sequences.

THEOREM. *Let l be an even integer having a prime factor $p \equiv 3 \pmod{4}$. Then there is no Barker sequence of length l .*

For the proof, we will need the following

LEMMA. *Let $f(z), g(z) \in \mathbb{F}_p[z, z^{-1}]$ be non-zero elements satisfying*

$$f(z) f(z^{-1}) + g(z) g(z^{-1}) = 0.$$

Then either $p = 2$ or $p \equiv 1 \pmod{4}$.

Proof. Since $\mathbb{F}_p[z, z^{-1}]$ is a unique factorization domain, we may suppose that $f(z), g(z)$ are coprime, by clearing any common factor. But then, the equation implies that $f(z)$ divides $g(z^{-1})$. We may thus write

$$g(z^{-1}) = h(z) f(z), \quad g(z) = h(z^{-1}) f(z^{-1})$$

for some $h(z) \in \mathbf{F}_p[z, z^{-1}]$. Substituting these expressions for $g(z)$ and $g(z^{-1})$ and clearing the common factor $f(z)f(z^{-1})$ in the resulting equation, we obtain

$$1 + h(z)h(z^{-1}) = 0 .$$

Letting $z = 1$, this gives $-1 = h(1)^2$ in \mathbf{F}_p , and therefore p is not congruent to $3 \pmod{4}$. \square

Proof of the Theorem. Let $A = (a_1, \dots, a_l)$ be a Barker sequence of even length l , and consider the two polynomials

$$F(z) = \sum_{i=1}^l a_i z^{i-1} \quad \text{and} \quad G(z) = F(-z) = \sum_{i=1}^l (-1)^{i-1} a_i z^{i-1} .$$

CLAIM: Then, (F, G) is a Golay pair, i.e.

$$F(z)F(z^{-1}) + G(z)G(z^{-1}) = 2l \quad \text{in } \mathbf{Z}[z, z^{-1}] .$$

Indeed, the constant term of $F(z)F(z^{-1}) + G(z)G(z^{-1})$ is equal to $2 \sum a_i^2 = 2l$. On the other hand, for $j > 0$, the coefficient of $z^j + z^{-j}$ in $F(z)F(z^{-1}) + G(z)G(z^{-1})$ is equal to

$$\sum_{i=1}^{l-j} (a_i a_{i+j} + (-1)^j a_i a_{i+j}) ,$$

which is zero if j is odd, and is equal to $2c_j(A)$ if j is even. But $c_j(A) = 0$ if j is even and positive, since $c_j(A)$ belongs to $\{-1, 0, 1\}$ by hypothesis, and $c_j \equiv j \pmod{2}$. Therefore, $F(z)F(z^{-1}) + G(z)G(z^{-1}) = 2l$ in $\mathbf{Z}[z, z^{-1}]$, as claimed.

Reducing the above equation modulo p , we obtain two non-zero elements $f(z), g(z)$ in $\mathbf{F}_p[z, z^{-1}]$ satisfying

$$f(z)f(z^{-1}) + g(z)g(z^{-1}) = 0 .$$

By the lemma above, we conclude that p cannot be congruent to $3 \pmod{4}$. \square

APPLICATION. There is no Barker sequence of length $l = 4N^2$, if $1 < N < 689$. In particular, there is no Barker sequence of even length greater than 4 and less than 1 898 884.

Of course, it suffices to consider only those $N < 689$ which are odd, are not a prime or a prime power, and have no factor congruent to $3 \pmod{4}$. Since the square root of 689 is smaller than 26, every such N must have a prime factor equal to 5, 13 or 17.

The remaining candidates are listed below, together with an indication in parenthesis showing that each one (except 505) is excluded by Theorem 2 in Section 2: if N has a prime factor p such that $p^f \equiv -1 \pmod{N'}$, where N' is the largest divisor of N relatively prime to p , then there is no (periodic) Barker sequence of length $4N^2$.

REMAINING CANDIDATES (excluded by Theorem 2, except $N = 505$.)

| N | | N | |
|----------------------|--------------------------------|----------------------|------------------------------|
| $65 = 5 \cdot 13$ | $(5^2 \equiv -1 \pmod{13})$ | $425 = 5^2 \cdot 17$ | $(5^8 \equiv -1 \pmod{17})$ |
| $85 = 5 \cdot 17$ | $(17^2 \equiv -1 \pmod{5})$ | $445 = 5 \cdot 89$ | $(89 \equiv -1 \pmod{5})$ |
| $145 = 5 \cdot 29$ | $(29 \equiv -1 \pmod{5})$ | $481 = 13 \cdot 37$ | $(37^6 \equiv -1 \pmod{13})$ |
| $185 = 5 \cdot 37$ | $(37^2 \equiv -1 \pmod{5})$ | $485 = 5 \cdot 97$ | $(97^2 \equiv -1 \pmod{5})$ |
| $205 = 5 \cdot 41$ | $(5^{10} \equiv -1 \pmod{41})$ | $493 = 17 \cdot 29$ | $(17^2 \equiv -1 \pmod{29})$ |
| $221 = 13 \cdot 17$ | $(13^2 \equiv -1 \pmod{17})$ | $505 = 5 \cdot 101$ | |
| $265 = 5 \cdot 53$ | $(53^2 \equiv -1 \pmod{5})$ | $533 = 13 \cdot 43$ | $(43^3 \equiv -1 \pmod{13})$ |
| $305 = 5 \cdot 61$ | $(5^{15} \equiv -1 \pmod{61})$ | $545 = 5 \cdot 109$ | $(109 \equiv -1 \pmod{5})$ |
| $325 = 5^2 \cdot 13$ | $(5^2 \equiv -1 \pmod{13})$ | $565 = 5 \cdot 113$ | $(113^2 \equiv -1 \pmod{5})$ |
| $365 = 5 \cdot 73$ | $(73^2 \equiv -1 \pmod{5})$ | $629 = 17 \cdot 37$ | $(37^8 \equiv -1 \pmod{17})$ |
| $377 = 13 \cdot 29$ | $(13^7 \equiv -1 \pmod{29})$ | $685 = 5 \cdot 137$ | $(137^2 \equiv -1 \pmod{5})$ |

The case $N = 505 = 5 \cdot 101$ cannot be excluded by Theorem 2, because $101 \equiv 1 \pmod{5}$ and $5^{25} \equiv 1 \pmod{101}$. However, 505 can still be excluded by Turyn's Inequality, as observed in [JL]: choosing $p = 101$ and $w = 2 \cdot 101^2$, so that p is trivially semi-primitive modulo w , we would have

$$p \leq \frac{v}{w} = 2 \cdot 5^2 = 50,$$

a contradiction to the assumed existence of a Barker sequence of length $4 \cdot 505^2$.

The first open case is thus $N = 689 = 13 \cdot 53$. We have $53 \equiv 1 \pmod{13}$ and $13^{13} \equiv 1 \pmod{53}$, so that neither 53 is semi-primitive mod 13, nor 13 is semi-primitive mod 53. The next open case is $N = 793 = 13 \cdot 61$.

4. THE USE OF THE MULTIPLIER THEOREM

In this section we give the details of some (typical) non-existence proofs needed to establish the tables, using the multiplier theorem.

Recall that if D is a cyclic difference set with parameters (v, k, λ) , and if $n = k - \lambda$ is greater than λ , then the group of multipliers of D contains the intersection M in $(\mathbf{Z}/v\mathbf{Z})^*$ of the subgroups generated by l_1, \dots, l_r , where l_1, \dots, l_r are the prime factors of n .