

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 38 (1992)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE CATEGORY OF NILMANIFOLDS  
**Autor:** Oprea, John  
**Kapitel:** §4. Category of nilmanifolds  
**DOI:** <https://doi.org/10.5169/seals-59480>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$$\begin{array}{ccccc}
S^1 & \rightarrow & M & = & K(\pi, 1) \\
& & \downarrow & & \\
S^1 & \rightarrow & M_{n-1} & \xrightarrow{\tau_n} & CP(\infty) \\
& & \downarrow & & \\
& & \vdots & & \\
& & \downarrow & & \\
S^1 & \rightarrow & M_1 & \xrightarrow{\tau_2} & CP(\infty) \\
& & \downarrow & & \\
& & * & \xrightarrow{\tau_1} & CP(\infty) .
\end{array}$$

We can assume (by finite dimensionality) that each  $\tau_i$  has image in a finite  $CP(n)$ , so thus may be approximated by a smooth map. Hence, each  $M_j$  is a compact manifold with

$$\dim(M_j) = \dim(M_{j-1}) + 1 .$$

Thus,  $\dim(M) = \text{rank}(\pi) = n$ .

#### §4. CATEGORY OF NILMANIFOLDS

The decomposition of  $M = K(\pi, 1)$  into a tower of principal  $S^1$ -bundles is, in fact, the Postnikov decomposition of  $M$  with  $k$ -invariants the  $\tau_i$ . By the fundamental theorem of rational homotopy theory, the minimal model has the form,

$$\Lambda(M) = (\Lambda(x_1, \dots, x_n), d) , \quad \deg(x_i) = 1$$

with  $dx_i = \tau_i$ , where  $\tau_i$  is a cocycle representing the class  $\tau_i \in H^2(M_{i-1}; \mathbf{Z})$ . Note that  $\Lambda(M)$  is an exterior algebra because all generators are in degree 1. Therefore, since  $\dim M = n$ , the only possibility for a cocycle representing the fundamental class is  $x_1 \cdots x_n$ . Hence,  $e_0(M) = n$  and this immediately implies,

*Proof of Theorem 1.*  $n = e_0(M) \leq \text{cat}_0(M) \leq \text{cat}(M) \leq \dim M = n$ .  $\square$

*Example.* Consider the 3-dimensional Heisenberg group  $U_3(\mathbf{R})$  and mod out by  $U_3(\mathbf{Z})$ . The resulting  $M$  is a 3-manifold obtained as a principal bundle,

$$S^1 \rightarrow M \rightarrow T^2$$

with classifying element (over the rationals)  $xy \in H^2(T^2; \mathbf{Q})$ , where  $x$  and  $y$  are one-dimensional generators. The minimal model of  $M$  is then given by

$$\Lambda(M) = \Lambda(x, y, z) \quad \deg(x) = \deg(y) = \deg(z) = 1$$

with  $dx = 0 = dy$  and  $dz = xy$ . Additive generators for cohomology are then,

$$H^1: x, y$$

$$H^2: xz, yz \text{ (Massey products!)}$$

$$H^3: zyx .$$

Note that  $\text{cup}(M) = 2$ , but  $\text{cat}(M) = 3$ .

In some sense then, the proof of Theorem 1 is simply an observation that the techniques of rational homotopy theory work particularly well for nilmanifolds.

PROBLEM. If  $\pi$  is not nilpotent, then a  $K(\pi, 1)$  is not a nilpotent space, so the minimal model does not describe a "rational type". Is it possible, however, that enough information about a  $K(\pi, 1)$  is present in the model to determine its category (in the compact case say)?

## §5. HIGHER DEGREE ANALOGUES

An analogue of the minimal model of a nilmanifold is one of the form,

$$(\Lambda(x_1, \cdots x_n), d) , \quad \text{degree}(x_i) = \text{odd} .$$

Such an algebra is known to satisfy rational Poincaré duality (see [5]) and to have formal top dimension  $\sum_i \text{deg}(x_i)$ . But, plainly, the same argument as before applies to show that the "only" element in this exterior algebra which can reach the stated dimension is  $x_1 \cdots x_n$ . Hence (since this is the longest product in  $\Lambda$ ), the fundamental class is maximally represented by a product of length  $n$  and

LEMMA.  $e_0(\Lambda) = n$ .

Now, we may consider  $\Lambda$  as built up by adjoining odd generators one at a time (with decomposable differential). Let  $\Lambda Z$  be a minimal cdga and  $y$  of odd degree. Then

PROPOSITION. (See Theorem 4.7 and Lemma 6.6 of [3].)

$$\text{cat}_0(\Lambda Z \otimes \Lambda y) \leq \text{cat}_0(\Lambda Z) + 1 .$$