

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 38 (1992)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** COMPLEX GROWTH SERIES OF COXETER SYSTEMS  
**Autor:** Paris, Luis  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-59485>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## COMPLEX GROWTH SERIES OF COXETER SYSTEMS

by Luis PARIS<sup>1)</sup>

### 1. INTRODUCTION

Let  $(W, S)$  be a *Coxeter system* (see [1] for the definitions). Throughout this paper the generating set  $S$  of  $W$  is assumed to be finite.  $S$  determines a length on  $W$  called *word length*. It is defined by

$$l(w) = l_S(w) = \min \{r \mid w = s_1 \dots s_r, s_i \in S\},$$

for  $w \in W$ . The *growth series* of  $W$  with respect to  $S$  is the formal series

$$W_S(t) = \sum_{w \in W} t^{l(w)}.$$

For a subset  $X \subseteq S$ , we denote by  $W_X$  the subgroup of  $W$  generated by  $X$ ; the system  $(W_X, X)$  is still a Coxeter system.

With a Coxeter system  $(W, S)$  one can associate a simplicial complex  $\Sigma(W, S)$ , called the *Coxeter complex*. This was introduced by Tits in [5] and is an essential ingredient of the theory of buildings (see [2] and [6]).

In this paper we introduce a new formal series  $W_S(t_1, t_2)$ , in two variables, which will be called the *complex growth series* of  $(W, S)$ , and is determined from the complex  $\Sigma(W, S)$ . More precisely,

$$W_S(t_1, t_2) = \sum_F t_1^{d(C_0, F)} t_2^{\text{codim}(F)},$$

where the sum is over all the faces  $F$  of  $\Sigma(W, S)$  (here we assume the empty set to be a face of  $\Sigma(W, S)$  of dimension  $-1$ ), and  $d(C_0, F)$  is the distance between the fundamental chamber  $C_0$  of  $\Sigma(W, S)$  and the face  $F$ .

The notions of Coxeter complex, chamber, face and fundamental chamber will be recalled in Section 2.

**MAIN THEOREM.** *Let  $(W, S)$  be a Coxeter system. Then*

$$(1.1) \quad W_S(t_1, t_2) = \sum_{X \subseteq S} t_2^{|X|} \frac{W_S(t_1)}{W_X(t_1)}.$$

---

<sup>1)</sup> Supported by the FNSRS (Swiss national Foundation).

Solomon proved in [4] that, if  $(W, S)$  is a finite Coxeter system (i.e.  $W$  is finite), then

$$(1.2) \quad \sum_{X \subseteq S} (-1)^{|X|} \frac{W_S(t)}{W_X(t)} = t^m,$$

where

$$m = \max_{w \in W} l(w).$$

Later, in [1, §4.1, exercise 26], Bourbaki proved a similar formula for an infinite system  $(W, S)$ ; in that case,

$$(1.3) \quad \sum_{X \subseteq S} (-1)^{|X|} \frac{W_S(t)}{W_X(t)} = 0.$$

Several results on growth series of Coxeter groups are obtained by induction on  $|S|$  using (1.2) and (1.3). We refer to [3] for an exposition on those two equalities and their applications.

An immediate corollary of (1.1), (1.2) and (1.3) is: if  $(W, S)$  is a finite Coxeter system, then

$$(1.4) \quad W_S(t_1, -1) = t_1^m,$$

$m$  being the maximal length in  $W$ ; and if  $(W, S)$  is an infinite Coxeter system, then

$$(1.5) \quad W_S(t_1, -1) = 0.$$

In fact, these two equalities (1.4) and (1.5) can and will be proved independently of the formulas (1.1), (1.2) and (1.3) (Proposition 2).

As an illustration of the Main Theorem, let us give two explicit examples.

1) Assume

$$W = \langle s_1, s_2 \mid s_1^2 = s_2^2 = (s_1 s_2)^3 = 1 \rangle$$

to be a Coxeter group of type  $A_2$ . The geometric realisation of  $\Sigma(W, S)$  is an hexagon. We have

$$W_S(t_1, t_2) = (1 + t_1)(1 + t_1 + t_1^2) + 2(1 + t_1 + t_1^2)t_2 + t_2^2,$$

and

$$W_{\{s_1, s_2\}}(t) = (1 + t)(1 + t + t^2),$$

$$W_{\{s_1\}}(t) = W_{\{s_2\}}(t) = 1 + t,$$

$$W_{\emptyset}(t) = 1.$$

Thus the equality (1.1) holds in that case.

2) Assume

$$W = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^3 = (s_1 s_3)^3 = (s_2 s_3)^3 = 1 \rangle$$

to be a Coxeter group of type  $\tilde{A}_2$ . The geometric realisation of  $\Sigma(W, S)$  is a plane. We have

$$\begin{aligned} W_S(t_1, t_2) &= \frac{(1+t_1)(1+t_1+t_1^2)}{(1-t_1)(1-t_1^2)} + 3 \frac{1+t_1+t_1^2}{(1-t_1)(1-t_1^2)} t_2 \\ &\quad + 3 \frac{1}{(1-t_1)(1-t_1^2)} t_2^2 + t_2^3, \end{aligned}$$

and

$$\begin{aligned} W_S(t) &= \frac{(1+t)(1+t+t^2)}{(1-t)(1-t^2)}, \\ W_{\{s_1, s_2\}}(t) &= W_{\{s_1, s_3\}}(t) = W_{\{s_2, s_3\}}(t) = (1+t)(1+t+t^2), \\ W_{\{s_1\}}(t) &= W_{\{s_2\}}(t) = W_{\{s_3\}}(t) = 1+t, \\ W_\emptyset(t) &= 1. \end{aligned}$$

Thus the equality (1.1) holds in that case.

In Section 2 we will recall some definitions in the theory of Coxeter complexes, we will define the complex growth series of a Coxeter system  $(W, S)$ , we will prove that  $W_S(t_1, 0) = W_S(t_1)$  and  $W_S(0, t_2) = (1+t_2)^{|S|}$  (Proposition 1), we will prove the equalities (1.4) and (1.5) (Proposition 2), and we will prove the Main Theorem.

## 2. COMPLEX GROWTH SERIES

We assume the reader to be familiar with the notions of simplicial complex, chamber complex, adjacency between two chambers, gallery and labelling. We refer to [2, Chap. I, Appendix] for a good exposition of these notions.

Let  $(W, S)$  be a Coxeter system. A *special coset* of  $(W, S)$  is a coset  $wW_X$ , with  $w \in W$  and  $X \subseteq S$ . We denote by  $\Sigma = \Sigma(W, S)$  the poset of all special cosets, ordered by the reverse inclusion;  $B \leq A$  in  $\Sigma$  if  $B \supseteq A$  in  $W$ . The poset  $\Sigma$  is a labelled chamber simplicial complex (see [2, Chap. III, §1]).

A *chamber* of  $\Sigma$  is a singleton  $\{w\}$  with  $w \in W$ . A *vertex* of  $\Sigma$  is a special coset  $wW_{S-\{s\}}$  with  $w \in W$  and  $s \in S$ . The face of  $\Sigma$  of dimension  $-1$  is the