

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 38 (1992)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** COMPLEX GROWTH SERIES OF COXETER SYSTEMS  
**Autor:** Paris, Luis  
**Kapitel:** 2. Complex growth series  
**DOI:** <https://doi.org/10.5169/seals-59485>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Thus the equality (1.1) holds in that case.

2) Assume

$$W = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^3 = (s_1 s_3)^3 = (s_2 s_3)^3 = 1 \rangle$$

to be a Coxeter group of type  $\tilde{A}_2$ . The geometric realisation of  $\Sigma(W, S)$  is a plane. We have

$$\begin{aligned} W_S(t_1, t_2) &= \frac{(1+t_1)(1+t_1+t_1^2)}{(1-t_1)(1-t_1^2)} + 3 \frac{1+t_1+t_1^2}{(1-t_1)(1-t_1^2)} t_2 \\ &\quad + 3 \frac{1}{(1-t_1)(1-t_1^2)} t_2^2 + t_2^3, \end{aligned}$$

and

$$\begin{aligned} W_S(t) &= \frac{(1+t)(1+t+t^2)}{(1-t)(1-t^2)}, \\ W_{\{s_1, s_2\}}(t) &= W_{\{s_1, s_3\}}(t) = W_{\{s_2, s_3\}}(t) = (1+t)(1+t+t^2), \\ W_{\{s_1\}}(t) &= W_{\{s_2\}}(t) = W_{\{s_3\}}(t) = 1+t, \\ W_\emptyset(t) &= 1. \end{aligned}$$

Thus the equality (1.1) holds in that case.

In Section 2 we will recall some definitions in the theory of Coxeter complexes, we will define the complex growth series of a Coxeter system  $(W, S)$ , we will prove that  $W_S(t_1, 0) = W_S(t_1)$  and  $W_S(0, t_2) = (1+t_2)^{|S|}$  (Proposition 1), we will prove the equalities (1.4) and (1.5) (Proposition 2), and we will prove the Main Theorem.

## 2. COMPLEX GROWTH SERIES

We assume the reader to be familiar with the notions of simplicial complex, chamber complex, adjacency between two chambers, gallery and labelling. We refer to [2, Chap. I, Appendix] for a good exposition of these notions.

Let  $(W, S)$  be a Coxeter system. A *special coset* of  $(W, S)$  is a coset  $wW_X$ , with  $w \in W$  and  $X \subseteq S$ . We denote by  $\Sigma = \Sigma(W, S)$  the poset of all special cosets, ordered by the reverse inclusion;  $B \leq A$  in  $\Sigma$  if  $B \supseteq A$  in  $W$ . The poset  $\Sigma$  is a labelled chamber simplicial complex (see [2, Chap. III, § 1]).

A *chamber* of  $\Sigma$  is a singleton  $\{w\}$  with  $w \in W$ . A *vertex* of  $\Sigma$  is a special coset  $wW_{S-\{s\}}$  with  $w \in W$  and  $s \in S$ . The face of  $\Sigma$  of dimension  $-1$  is the

coset  $1 \cdot W = W$  (this face has 0 vertices). The *fundamental chamber* of  $\Sigma$  is  $\{1\}$ .

The Coxeter group  $W$  naturally acts on  $\Sigma$  by

$$(2.1) \quad w(vW_X) = (wv)W_X,$$

where  $w \in W$ , and  $vW_X$  is a face of  $\Sigma$  (i.e. a special coset).

The map which associates to a face  $F = wW_X$  the subset  $\lambda(F) = S - X$  of  $S$  determines a labelling on  $\Sigma$ , called the *canonical labeling* of  $\Sigma$ , where  $\lambda(F)$  is the *type* of a face  $F$ .

Two chambers  $\{w\} \neq \{w'\}$  are *adjacent* if they have a common codimension 1 face, namely, if there exists an  $s \in S$  such that  $w' = ws$ . A *gallery of length  $d$*  is a sequence  $\{C_i\}_{i=0}^d$  of  $d + 1$  chambers such that  $C_i$  and  $C_{i+1}$  are adjacent for  $i = 0, 1, \dots, d - 1$ . In fact, to give a gallery  $\{C_i\}_{i=0}^d$  is equivalent to give a source chamber  $C_0$  and a sequence  $s_1, \dots, s_d$  of elements of  $S$ ; the equivalence is given by  $C_i = s_i \dots s_2 s_1(C_0)$ . A gallery  $\{C_i\}_{i=1}^d$  joining two chambers  $C_0$  and  $C_d$  is called *minimal* if there is no gallery joining  $C_0$  and  $C_d$  with a smaller length.

The *distance*  $d(C, D)$  between two chambers  $C$  and  $D$  is the length of a minimal gallery joining  $C$  and  $D$ . We can easily see that, if  $C = \{w\}$  and  $D = \{v\}$ , then

$$(2.2) \quad d(C, D) = l(w^{-1}v).$$

The *distance*  $d(C, F)$  between a chamber  $C$  and a face  $F$  of  $\Sigma$  is

$$(2.3) \quad d(C, F) = \min \{d(C, D) \mid D \text{ a chamber having } F \text{ as face}\}.$$

As in (2.2), if  $C = \{w\}$  and  $F = vW_X$ , then

$$(2.4) \quad d(C, F) = \min \{l(u) \mid u \in w^{-1}vW_X\}.$$

The *complex growth series* of a Coxeter system  $(W, S)$  is the formal series in two variables

$$(2.5) \quad W_S(t_1, t_2) = \sum_F t_1^{d(C_0, F)} t_2^{\text{codim}(F)},$$

where the sum is over all the faces  $F$  of  $\Sigma$ , and where  $C_0 = \{1\}$  is the fundamental chamber.

Before stating and proving Propositions 1 and 2 and the Main Theorem, we are going to state two known results (Lemmas 1 and 2). A proof of Lemma 1 can be found either in [1, §4.1, exercise 3] or in [3, Lemma 1]. A proof of Lemma 2 can be found in [2, Chap. IV, §6].

Let  $X \subseteq S$  be a subset and let  $v \in W$ . The element  $v$  is called *X-minimal* if  $v$  is of minimal length among the elements of  $vW_X$ .

LEMMA 1. *Let  $X \subseteq S$  be a subset and let  $v \in W$  be an X-minimal element of  $W$ . Then*

- i)  $v$  is the unique *X-minimal* element of  $vW_X$ ,
- ii) for every  $w = vu \in vW_X$ , with  $u = v^{-1}w \in W_X$ , one has  $l(w) = l(v) + l(u)$ .

For an integer  $d \geq 0$ , we denote by  $\Sigma_d$  the subcomplex of  $\Sigma = \Sigma(W, S)$  generated by the chambers  $C$  of  $\Sigma$  at distance  $\leq d$  of  $C_0 = \{1\}$ .

$$\Sigma_d = \bigcup_F F,$$

where the union is over all the faces  $F$  of  $\Sigma$  such that  $d(C_0, F) \leq d$ . We denote by  $|\Sigma_d|$  the geometric realization of  $\Sigma_d$ .

LEMMA 2. i) Let  $(W, S)$  be a finite Coxeter system. Set  $m = \max_{w \in W} l(w)$ . Then  $|\Sigma_d|$  is contractible if  $d < m$ , and  $|\Sigma_d|$  is homotopic to the sphere  $S^{|S|-1}$  of dimension  $|S| - 1$  if  $d \geq m$ .

ii) Let  $(W, S)$  be an infinite Coxeter system. Then  $|\Sigma_d|$  is contractible.

PROPOSITION 1. *Let  $(W, S)$  be a Coxeter system. Then*

$$(2.6) \quad W_S(t_1, 0) = W_S(t_1) \quad \text{and}$$

$$(2.7) \quad W_S(0, t_2) = (1 + t_2)^{|S|}.$$

*Proof.*

$$W_S(t_1, 0) = \sum_F t_1^{d(C_0, F)},$$

where the sum is over all the faces of  $\Sigma$  of codimension 0, i.e. over all the chambers of  $\Sigma$ . Furthermore, if  $F = C = \{w\}$ , then, by (2.2),  $d(C_0, F) = l(w)$ . It follows that

$$W_S(t_1, 0) = \sum_{w \in W} t_1^{l(w)} = W_S(t_1).$$

Now,

$$W_S(0, t_2) = \sum_F t_2^{\text{codim}(F)},$$

where the sum is over all the faces  $F$  of  $\Sigma$  at distance 0 of  $C_0$ , i.e. over all the faces of  $C_0$ . Since  $C_0$  is an  $|S| - 1$  dimensional simplex, it has  $\binom{|S|}{i}$  faces of dimension  $i$  (where  $i = 0, 1, \dots, |S|$ ). It follows that

$$W_S(0, t_2) = \sum_{i=0}^{|S|} \binom{|S|}{i} t_2^i = (1 + t_2)^{|S|}. \quad \square$$

**PROPOSITION 2.** i) Let  $(W, S)$  be a finite Coxeter system. Then

$$(2.8) \quad W_S(t_1, -1) = t_1^m,$$

where  $m$  is the maximal length in  $W$ .

ii) Let  $(W, S)$  be an infinite Coxeter system. Then

$$(2.9) \quad W_S(t_1, -1) = 0.$$

*Proof.* Recall that  $\Sigma_d$  is the subcomplex of  $\Sigma$  generated by the chambers of  $\Sigma$  at distance  $\leq d$  of  $C_0$ , and that  $|\Sigma_d|$  is the geometric realization of  $\Sigma_d$ . We denote by  $E(|\Sigma_d|)$  the Euler characteristic of  $|\Sigma_d|$ . It is well known that  $E(|\Sigma_d|)$  can be computed as follows:

$$\begin{aligned} (-1)^{|S|-1} E(|\Sigma_d|) &= (-1)^{|S|-1} \sum_{\substack{d(C_0, F) \leq d \\ F \neq W}} (-1)^{\dim(F)} \\ &= \sum_{\substack{d(C_0, F) \leq d \\ F \neq W}} (-1)^{\text{codim}(F)}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} W_S(t_1, -1) &= \sum_F t_1^{d(C_0, F)} (-1)^{\text{codim}(F)} \\ &= \sum_{d=0}^{\infty} \left( \sum_{d(C_0, F) = d} (-1)^{\text{codim}(F)} \right) t_1^d. \end{aligned}$$

Thus

$$(2.10) \quad (-1)^{|S|-1} (E(|\Sigma_d|) - E(|\Sigma_{d-1}|))$$

is the coefficient of  $t_1^d$  in  $W_S(t_1, -1)$  for  $d \geq 1$ , and

$$(2.11) \quad (-1)^{|S|-1} E(|\Sigma_0|) + (-1)^{|S|}$$

is the coefficient of  $t_1^0$  in  $W_S(t_1, -1)$ . Lemma 2 implies that, if  $(W, S)$  is a finite Coxeter system, then

$$E(|\Sigma_d|) = \begin{cases} 1 & \text{if } d < m, \\ 1 + (-1)^{|S|-1} & \text{if } d \geq m, \end{cases}$$

where  $m$  is the maximal length in  $W$ ; and if  $(W, S)$  is an infinite Coxeter system, then

$$E(|\Sigma_d|) = 1,$$

for all  $d \geq 0$ . Replacing  $E(|\Sigma_d|)$  by its value in (2.10) and (2.11), we obtain the equalities (2.8) and (2.9).  $\square$

**MAIN THEOREM.** *Let  $(W, S)$  be a Coxeter system. Then*

$$(2.12) \quad W_S(t_1, t_2) = \sum_{X \subseteq S} t_2^{|X|} \frac{W_S(t_1)}{W_X(t_1)}.$$

*Proof.* Recall that the map which associates to a face  $F = wW_X$  the subset  $\lambda(F) = S - X$  of  $S$  determines a labelling on  $\Sigma$ , where  $\lambda(F)$  is the type of the face  $F$ . Clearly, if  $\lambda(F) = Y$ , then  $\dim(F) = |Y| - 1$  and  $\text{codim}(F) = |S| - |Y| = |S - Y|$ . Therefore

$$(2.13) \quad W_S(t_1, t_2) = \sum_{Y \subseteq S} t_2^{|S - Y|} \left( \sum_{F \in \mathcal{F}_Y} t_1^{d(C_0, F)} \right),$$

where  $\mathcal{F}_Y$  is the set of faces of  $\Sigma$  of type  $Y$ . Let us prove

$$(2.14) \quad \sum_{F \in \mathcal{F}_Y} t_1^{d(C_0, F)} = \frac{W_S(t_1)}{W_{S-Y}(t_1)},$$

for every  $Y \subseteq S$ . The equalities (2.13) and (2.14) clearly imply (2.12).

Let  $X = S - Y$ . Recall that an element  $v \in W$  is  $X$ -minimal if it is of minimal length in  $vW_X$ . Every face  $F \in \mathcal{F}_Y$  can be written  $F = vW_X$  with  $v$   $X$ -minimal (take any element of minimal length in  $F$ ). By (2.4), we have

$$d(C_0, F) = l(v).$$

Lemma 1 shows that, for every  $F \in \mathcal{F}_Y$ , there is an unique  $X$ -minimal element  $v$  in  $F$ . Therefore

$$(2.15) \quad \sum_{F \in \mathcal{F}_Y} t_1^{d(C_0, F)} = \sum_{v \in A_X} t_1^{l(v)},$$

where  $A_X$  is the set of all the  $X$ -minimal elements of  $W$ . Finally, Lemma 1 shows

$$\begin{aligned}
 W_S(t_1) &= \sum_{w \in W} t_1^{l(w)} \\
 &= \sum_{v \in A_X} \sum_{w \in vW_X} t_1^{l(w)} \quad (\text{Lemma 1.i}) \\
 &= \sum_{v \in A_X} \sum_{u \in W_X} t_1^{l(v) + l(u)} \quad (\text{Lemma 1.ii}) \\
 &= \left( \sum_{v \in A_X} t_1^{l(v)} \right) W_X(t_1).
 \end{aligned}$$

This and (2.15) imply (2.14)  $\square$

#### REFERENCES

- [1] BOURBAKI, N. *Groupes et algèbres de Lie*, Chap. 4, 5 et 6. Hermann, Paris, 1968.
- [2] BROWN, K. S. *Buildings*. Springer-Verlag, New York, 1989.
- [3] PARIS, L. Growth series of Coxeter groups. In: *Group Theory from a Geometrical Viewpoint*, (Trieste, 1990), pp. 302-310, World Scientific Publishing, Singapore, 1991.
- [4] SOLOMON, L. The order of the finite Chevalley groups. *J. of Algebra* 3 (1966), 376-393.
- [5] TITS, J. Groupes et géométries de Coxeter. 1961 (unpublished manuscript).
- [6] —— On buildings and their applications. In: *Proceedings of the International Congress of Mathematicians*, Vancouver, 1974, Vol. 1, Canad. Math. Congress, Montreal, 1975, pp. 209-220.

(Reçu le 1<sup>er</sup> août 1991)

Luis Paris

Department of Mathematics  
University of Wisconsin  
480 Lincoln dr.  
Madison, WI 53706, USA

Section de Mathématiques  
Université de Genève  
CP 240  
CH-1211 Genève 24, Switzerland