

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 38 (1992)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ANALOGUE OF HUBER'S FORMULA FOR RIEMANN'S ZETA FUNCTION
Autor: Williams, Floyd L.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-59488>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

AN ANALOGUE OF HUBER'S FORMULA
FOR RIEMANN'S ZETA FUNCTION

by Floyd L. WILLIAMS ¹⁾

To the memory of Michio Kuga

1. INTRODUCTION

A remarkable formula of H. Huber [11] relates the class 1 spectrum of a compact Riemann surface X and the spectral zeros of Selberg's zeta function ζ_X of X . More generally, if X is a space form (not necessarily compact) of a rank 1 symmetric space one can still assign to X a Selberg zeta function ζ_X and formulate a generalized version of Huber's formula [6], [18]. Here a decisive role is played by the Selberg trace formula.

On the other hand Weil's explicit sum formula for Riemann's zeta function ζ [16], [17] bears some striking similarity in appearance to the trace formula. It is now known, as a matter of fact, by a recent work of D. Goldfeld [7] that there exists a kernel function on a suitable space such that the conjugacy class sum in Selberg's trace formula is precisely the sum over the primes of Weil's formula; i.e. Weil's formula indeed can be interpreted as a trace formula.

Motivated by a certain "radial" function which occurs in semisimple Lie theory we consider a certain test function which we plug into Weil's formula, and we derive thereby a formula of Huber type for ζ . The formula, see Theorem 7.1, involves a sum over the "spectrum" of ζ — i.e. over its non-trivial zeros. We derive a second interesting formula in Theorem 7.10 by specializing the parameter s in Theorem 7.1.

Although Lie theory and spectral theory serve as a context and motivation, the lecture which is largely self-contained requires no familiarity with these subjects. We assume only a knowledge of basic real and complex analysis.

¹⁾ This is an expanded version of an invited Mathematical Association of America address delivered at the winter San Francisco meeting on January 16, 1991.

The author is thankful for and honored by the invitation extended by the Program Committee of the Mathematical Association of America to deliver this lecture. We dedicate the lecture to the memory of a very great mathematician — a kind and humble man — a friend — Professor Michio Kuga.

2. OUTLINE OF THE LECTURE

- I. Huber's formula (as a context)
- II. Riemann's zeta function — basic facts and the Riemann hypothesis (= RH)
- III. Test functions
- IV. Weil's explicit formula
- V. The Schwartz space and the RH
- VI. The main test function
- VII. An analogue of Huber's formula

I. HUBER'S FORMULA (as a context)

Since Huber's formula provides the motivation for this lecture we shall state (for the record) this remarkable result. Neither the result nor any understanding of it is required for later purposes.

Let G denote the group $SL(2, \mathbf{R})$ of real 2×2 matrices with determinant equal to 1:

$$(1.1) \quad G = SL(2, \mathbf{R}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid \begin{array}{l} a, b, c, d, \in \mathbf{R} \\ ad - cb = 1 \end{array} \right\}$$

where \mathbf{R} denotes the field of real numbers. Let $\Gamma \subset G$ be a discrete torsionfree¹⁾ subgroup such that the quotient $\Gamma \backslash G$ is compact. Euler's classical gamma function will also be denoted by Γ :

$$(1.2) \quad \Gamma(s) = \int_0^{\infty} e^{-t} t^{s-1} dt, \quad \text{Res} > 0.$$

The function $N: \Gamma - \{1\} \rightarrow \mathbf{R}$ defined by

$$(1.3) \quad N(P) = \max |c|^2, \quad c = \text{an eigenvalue of } P$$

¹⁾ That is, if $\gamma^n = 1$ for $\gamma \in \Gamma$, $n > 0$ an integer, then $\gamma = 1$.