Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A

SURVEY

Autor: Shallit, Jeffrey

Kapitel: 2. Numbers of Constant Type

DOI: https://doi.org/10.5169/seals-59489

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 10.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

If x is rational, then its continued fraction expansion terminates, and we can write $x = [a_0, a_1, ..., a_n]$. If we agree that $a_n = 1$ and $n \ge 1$, then this expansion is unique and we define

$$K(x) = \max_{1 \leq k \leq n} a_k ,$$

the largest partial quotient in the continued fraction for x.

If x is irrational, then its continued fraction expansion does not terminate. This expansion is unique. We write $x = [a_0, a_1, a_2, ...]$ and define

$$K(x) = \sup_{k \geqslant 1} a_k.$$

If $K(x) < \infty$, then we say that x has bounded partial quotients.

We define $\mathcal{B}_k = \{x \in \mathbf{R} \mid K(x) \leq k\}$, and $\mathcal{B} = \{x \in \mathbf{R} \mid K(x) < \infty\}$. Furthermore, let $\mathcal{E}_k = \mathcal{B}_k \cap (0, 1)$ and $\mathcal{E} = \mathcal{B} \cap (0, 1)$.

Real numbers with bounded partial quotients appear in many fields of mathematics and computer science: Diophantine approximation, fractal geometry, transcendental number theory, ergodic theory, numerical analysis, pseudo-random number generation, dynamical systems, and formal language theory. In this paper we survey some of these applications. Because of limited space, we cannot include a discussion of every result in detail. However, we have tried to include as complete a list of references as possible for those topics directly related to the main subject. Readers who know of other references are urged to contact the author (and provide a copy of the relevant paper, if possible). It is hoped that the list of references may contain some surprises even for experts in the field.

The author's interest in the subject arose from the material in Section 9. Because of this, the viewpoint presented in this article may be somewhat idiosyncratic.

2. Numbers of Constant Type

Let θ be an irrational number, and let $\|\theta\|$ denote the distance between θ and the closest integer.

Let $r \ge 1$ be a real number. We say that θ is of type < r if

$$q \parallel q\theta \parallel \geqslant \frac{1}{r}$$

for all integers $q \ge 0$. Then we have the following

THEOREM 1. If θ is of type < r, then $K(\theta) < r$. If $K(\theta) = r$, then θ is of type < r + 2.

For a proof, see Baker [20, p. 47] or Schmidt [272, p. 22].

If there exists an $r < \infty$ such that θ is of type < r, then θ is said to be of constant type. By the theorem, numbers of constant type and numbers with bounded partial quotients coincide, and we will use these terms interchangeably in what follows.

A classical theorem of Lagrange states that the continued fraction for x is ultimately periodic if and only if x is a real quadratic irrational, and so all real quadratic irrationals are of constant type; see, for example, Lagrange [178] or Hardy and Wright [135, Chapter 10]. We will not explicitly discuss quadratic irrationals further in this paper.

Since

$$e = [2, 1, 2, 1, 1, 4, 1, 1, 6, 1, 1, 8, 1, 1, 10, ...]$$

(see Cotes [58] and Euler [102]), we see that e is not of constant type. It is also known that the numbers $e^{2/n}$ and $\tan 1/n$ (n an integer ≥ 1) are not of constant type. The status of π and γ (Euler's constant) is presently unknown. In section 9 we will see some explicit examples of transcendental numbers of constant type.

One way to interpret Theorem 1 is to say that numbers with bounded partial quotients are *badly approximable* by rationals; this term is also used frequently in the literature.

Note that if θ is of constant type, and r is a rational number, then $r\theta$ is also of constant type [54]. In fact, it is not hard to prove the following: let r = a/b be a rational number, and suppose $K(\theta) = n$. Then $K(r\theta) \le |ab|(n+2)$, and $K(\theta+r) \le b^2(n+2)$; see Cusick and Mendès France [69].

From this, it follows that if a, b, c, d are integers with $ad - bc \neq 0$, then

$$\frac{a\theta + b}{c\theta + d}$$

has bounded partial quotients iff θ does. (See Shallit [278]. I would like to thank J. C. Lagarias for bringing this to my attention.) One can also deduce this result directly from the continued fraction, using results of Raney [256]. For another view of Raney's results, see van der Poorten [246].

Another related concept is the *Lagrange-Markoff* constant, denoted by $\mu(\theta)$. It is defined as follows:

$$\mu(\theta)^{-1} = \lim_{q \to \infty} \inf q \| q\theta \|.$$

Hurwitz [150] showed, among other things, that $\mu(\theta) \geqslant \sqrt{5}$; furthermore, $\mu\left(\frac{1+\sqrt{5}}{2}\right) = \sqrt{5}$. Perron [234] showed that if

$$\theta = [a_0, a_1, a_2, \ldots] ,$$

then

$$\mu(\theta) = \limsup_{i \to \infty} ([a_{i+1}, a_{i+2}, a_{i+3}, ...] + [0, a_i, a_{i-1}, ..., a_1]).$$

From this it follows that $\mu(\theta) < \infty$ if and only if θ is of constant type. The range of $\mu(\theta)$, as θ takes on all irrational values, is known as the Lagrange spectrum and has been extensively studied. We direct the reader to the work of Lagrange [178, pp. 26-27]; Markoff [203, 204]; Heawood [138]; Perron [235]; Vinogradov, Delone, and Fuks [295]; Freiman [111]; Kinney and Pitcher [166]; Berštein [29]; Davis and Kinney [78]; Cusick [59, 62]; Flahive [117]; Cusick and Mendès France [69]; Wilson [301]; Dietz [89]; Pavone [232]; Prasad [249]; and especially the books of Koksma [172] and Cusick and Flahive [67].

For more on approximation by rational numbers, see Cassels [52], Schmidt [272], Kraaikamp and Liardet [313], Larcher [312].

3. The Metric Theory of Continued Fractions

Recall that \mathscr{E} denotes the set of real numbers in (0, 1) with bounded partial quotients.

While it is easy to see \mathscr{E} has uncountably many elements, nevertheless "most" numbers do *not* have bounded partial quotients. More precisely, we have the following

THEOREM 2 (Borel-Bernstein). & is a set of measure 0.

The theorem is due to Borel [38]. The original proof was not complete, as discussed in Bernstein [27]; further details were provided in a later paper of Borel [39]. For other proofs, see Hardy and Wright [135, Thm. 196] or Khintchine [160]. Also see Dyson [96].

Here is a sketch of a more general theorem: first, let us equate probability with Lebesgue measure, and assume x is a real number in (0, 1). Then, expanding x as a continued fraction, we have

$$x = [0, a_1, a_2, a_3, ...],$$