Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A SURVEY

Autor: Shallit, Jeffrey

Kapitel: 5. Certain Sums in Diophantine Approximation

DOI: https://doi.org/10.5169/seals-59489

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

5. Certain Sums in Diophantine Approximation

Let us agree to write $\{\theta\}$ for the fractional part of θ, namely, $\theta - [\theta]$.

One of the earliest appearances of real numbers with bounded partial quotients is in the theory of Diophantine approximation.

For example, consider the sum

$$s_n'(\theta) = \sum_{1 \leqslant k \leqslant n} \left(\{k\theta\} - \frac{1}{2} \right) .$$

Clearly $s_n'(\theta) = O(n)$; but Lerch proved in 1904 that if θ has bounded partial quotients, we have $s_n'(\theta) = O(\log n)$. See [184]. (This result was also announced by Hardy and Littlewood in 1912; see [128].)

At the International Congress of Mathematicians in 1912, Hardy and Littlewood [128] announced several theorems on Diophantine approximation, some of which relate to the subject at hand. For example, they defined

$$s_n(\theta) = \sum_{1 \leqslant k \leqslant n} e^{\left(k - \frac{1}{2}\right)^2 \pi i \theta} ,$$

and stated that if θ has bounded partial quotients, then $s_n(\theta) = O(\sqrt{n})$. The proof appeared later; see [130].

At the same Congress, Hardy and Littlewood announced that

$$\sum_{1 \leqslant k \leqslant n} \left(\{k\theta\} - \frac{1}{2} \right)^2 = \frac{n}{12} + O(1) ,$$

for all irrational θ. This is incorrect, and the correct formulation was stated in a 1922 paper: the result holds for many, but not all irrationals, and in particular it holds for θ with bounded partial quotients. See [132] for the statement and [133] for a proof.

Hardy and Littlewood also examined other series of interest. They defined:

$$U_n(\theta) = \sum_{1 \leqslant k \leqslant n} \frac{(-1)^k}{k \sin k\pi\theta} ,$$

$$V_n(\theta) = \sum_{1 \leqslant k \leqslant n} \frac{(-1)^k}{\sin k\pi\theta} ,$$

and

$$W_n(\theta) = \sum_{1 \leqslant k \leqslant n} \frac{1}{(\sin k\pi\theta)^2} .$$

They showed that if θ has bounded partial quotients, then $U_n(\theta) = O(\log n)$, $V_n(\theta) = O(n)$, and $W_n(\theta) = O(n^2)$. See [134].

(Warning to the reader: in their papers, Hardy and Littlewood used the notation $\{x\}$ to mean $x - [x] - \frac{1}{2}$, not $x - [x]$, as is more standard today.)

For other related papers, see Hardy and Littlewood [129, 131]; the collected works of Hardy [127]; Ostrowski [230]; Khintchine [161]; Oppenheim [228]; Chowla [53, 54]; Walfisz [298, 299, 300], and Schoissengeier [314].

Others researchers have examined similar sums in connection with numbers with bounded partial quotients. See the papers of Faĭziev [104] Ivanov [151], and Schoissengeier [274].

6. Fractal Geometry

Numbers with bounded partial quotients provided an early example of a set with non-integral Hausdorff dimension.

Let $\dim S$ denote the Hausdorff dimension of the set S (for a definition, see, e.g. Falconer [103]). We use the definitions of $\mathscr{E}$ and $\mathscr{E}_k$ from section 1.

In 1928, Jarník [152] proved that $\dim \mathscr{E} = 1$,

$$\frac{1}{4} < \dim \mathscr{E}_2 < 1 ,$$

and

$$1 - \frac{4}{k \log 2} < \dim \mathscr{E}_k < 1 - \frac{1}{8k \log k},$$

for $k > 8$. An exposition of Jarník's work can be found in Rogers [263].

In 1941, Good proved the following result [118]:

$$\dim \mathscr{E}_k = \lim_{n \to \infty} \sigma_{k,n} ,$$

where $\sigma = \sigma_{k,n}$ is the real root of the equation

$$\sum_{1 \leqslant a_1, a_2, \ldots, a_n \leqslant k} Q(a_1, a_2, \ldots, a_n)^{-2\sigma} = 1$$

and $Q(\,)$ denotes Euler's continuant polynomial. (These are multivariate polynomials, defined by $Q(\,) = 1$, $Q(a_1) = a_1$, and

$$Q(a_1, a_2, \ldots a_n) = a_n Q(a_1, a_2, \ldots, a_{n-1}) + Q(a_1, a_2, \ldots, a_{n-2})$$

for $n \geqslant 2$.)