

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 38 (1992)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A SURVEY
Autor: Shallit, Jeffrey
Kapitel: 7. Schmidt's Game
DOI: <https://doi.org/10.5169/seals-59489>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 18.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Good also obtained the estimate $.5306 < \dim \mathcal{E}_2 < .5320$. This was improved by Bumby [48] in 1985 to $.5312 \leq \dim \mathcal{E}_2 \leq .5314$. More recently, Hensley [140] showed that $.53128049 < \dim \mathcal{E}_2 < .53128051$. For other results on the Hausdorff dimension of \mathcal{E}_k and related sets, see Jarník [153]; Besicovitch [30]; Rogers [262]; Baker and Schmidt [21]; Hirst [147, 148]; Billingsley and Henningsen [32]; Cusick [63, 64, 65]; Pollington [245]; Kaufman [158]; Marion [202]; Gardner and Mauldin [115]; Ramharter [253, 254]; and Hensley [139, 141, 308, 309].

7. SCHMIDT'S GAME

W. M. Schmidt [270] introduced the following two-player game, called an (α, β) game: let α, β be real numbers with $0 < \alpha, \beta < 1$. First Bob chooses a closed interval on the real line, called B_1 . Then Alice chooses a closed interval $A_1 \subset B_1$, such that the length of A_1 is α times the length of B_1 . Then Bob chooses a closed interval $B_2 \subset A_1$, such that the length of B_2 is β times the length of A_1 , and so on. If the intersection of all the intervals A_i is a number with bounded partial quotients, then Alice is declared the winner; otherwise Bob is declared the winner.

Schmidt showed that if $0 < \alpha < 1/2$, then Alice always has a winning strategy for this game. This is somewhat surprising, since as we have seen above, the set \mathcal{E} of numbers with bounded partial quotients has Lebesgue measure 0.

Using the theory of (α, β) games, Schmidt also reproved the result of Jarník that \mathcal{E} has Hausdorff dimension 1.

Several papers have proved other results on (α, β) games: see Schmidt [271]; Freiling [109, 110]; and Dani [70, 71, 72]. Also see Schmidt [272, Chapter 3].

8. HALL'S THEOREM

If S and T are sets, then by $S + T$ we mean the set

$$\{s + t \mid s \in S, t \in T\}.$$

Similarly, by $S \cdot T$ we mean the set

$$\{st \mid s \in S, t \in T\}.$$

If S is a set of Lebesgue measure zero, then it is quite possible for $S + S$ to have positive measure. For example, if C denotes the Cantor set (numbers