Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

**Band:** 38 (1992)

**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A

**SURVEY** 

Autor: Shallit, Jeffrey

**Kapitel:** 8. Hall's theorem

**DOI:** https://doi.org/10.5169/seals-59489

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Good also obtained the estimate .5306 < dim  $\mathscr{E}_2 <$  .5320. This was improved by Bumby [48] in 1985 to .5312  $\le$  dim  $\mathscr{E}_2 \le$  .5314. More recently, Hensley [140] showed that .53128049 < dim  $\mathscr{E}_2 <$  .53128051. For other results on the Hausdorff dimension of  $\mathscr{E}_k$  and related sets, see Jarník [153]; Besicovitch [30]; Rogers [262]; Baker and Schmidt [21]; Hirst [147, 148]; Billingsley and Henningsen [32]; Cusick [63, 64, 65]; Pollington [245]; Kaufman [158]; Marion [202]; Gardner and Mauldin [115]; Ramharter [253, 254]; and Hensley [139, 141, 308, 309].

# 7. SCHMIDT'S GAME

W. M. Schmidt [270] introduced the following two-player game, called an  $(\alpha, \beta)$  game: let  $\alpha, \beta$  be real numbers with  $0 < \alpha, \beta < 1$ . First Bob chooses a closed interval on the real line, called  $B_1$ . Then Alice chooses a closed interval  $A_1 \subset B_1$ , such that the length of  $A_1$  is  $\alpha$  times the length of  $B_1$ . Then Bob chooses a closed interval  $B_2 \subset A_1$ , such that the length of  $B_2$  is  $\beta$  times the length of  $A_1$ , and so on. If the intersection of all the intervals  $A_i$  is a number with bounded partial quotients, then Alice is declared the winner; otherwise Bob is declared the winner.

Schmidt showed that if  $0 < \alpha < 1/2$ , then Alice always has a winning strategy for this game. This is somewhat surprising, since as we have seen above, the set  $\mathscr{E}$  of numbers with bounded partial quotients has Lebesgue measure 0.

Using the theory of  $(\alpha, \beta)$  games, Schmidt also reproved the result of Jarník that  $\mathscr{E}$  has Hausdorff dimension 1.

Several papers have proved other results on  $(\alpha, \beta)$  games: see Schmidt [271]; Freiling [109, 110]; and Dani [70, 71, 72]. Also see Schmidt [272, Chapter 3].

## 8. Hall's theorem

If S and T are sets, then by S + T we mean the set

$$\{s+t \mid s \in S, t \in T\}$$
.

Similarly, by  $S \cdot T$  we mean the set

$$\{st \mid s \in S, t \in T\}$$
.

If S is a set of Lebesgue measure zero, then it is quite possible for S + S to have positive measure. For example, if C denotes the Cantor set (numbers

in [0, 1] containing only 0's and 2's in their ternary expansion), then C has measure 0, and it is not hard to show that C + C = [0, 2]; see Borel [40] or Pavone [233]. The result is due to Steinhaus [310]; I am most grateful to G. Myerson for bringing this to my attention.

As we have seen above, the set  $\mathcal{B}$ , and hence each  $\mathcal{B}_k$ , also has Lebesgue measure zero. In 1947 Hall proved the following theorem [126]:

THEOREM 3. Every real number x can be written as x = y + z, where  $y, z \in \mathcal{B}_4$ . Every real number  $x \ge 1$  can be written as x = yz, where  $y, z \in \mathcal{B}_4$ .

An exposition of Hall's result can be found in Cusick and Flahive [67]. Using the notation of the first paragraph of this section, we could rephrase the statement of Hall's theorem as follows:  $\mathcal{B}_4 + \mathcal{B}_4 = \mathbf{R}$ , and  $[1, \infty) \subseteq \mathcal{B}_4 \cdot \mathcal{B}_4$ .

In 1973, Cusick [61] proved that  $\mathcal{B}_3 + \mathcal{B}_3 + \mathcal{B}_3 = \mathbf{R}$ , and  $\mathcal{B}_2 + \mathcal{B}_2 + \mathcal{B}_2 + \mathcal{B}_2 = \mathbf{R}$ . He also observed that  $\mathcal{B}_3 + \mathcal{B}_3 \neq \mathbf{R}$ , and  $\mathcal{B}_2 + \mathcal{B}_2 + \mathcal{B}_2 \neq \mathbf{R}$ . These results were independently discovered by Diviš [90] and J. Hlavka 1) [149]. Hlavka also showed that  $\mathcal{B}_3 + \mathcal{B}_4 = \mathbf{R}$ , and similar results. Apparently the status of  $\mathcal{B}_2 + \mathcal{B}_5$  and  $\mathcal{B}_2 + \mathcal{B}_6$  is still open.

For results of a similar character, see Cusick [60]; Cusick and Lee [68]; and Bumby [47].

# 9. EXPLICIT EXAMPLES OF TRANSCENDENTAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS

In Lang [179] we find the following statement:

No simple example of [irrational] numbers of constant type, other than the one given above [real quadratic irrationals], is known. The best guess is that there are no other "natural" examples.

(Also see Lang [180].)

However, in 1979 Kmošek [167] and Shallit [275] independently discovered the following "natural" example of numbers of constant type.

THEOREM 4. Let  $n \ge 2$  be an integer and define

(1) 
$$f(n) = \sum_{i \geq 0} n^{-2i}.$$

<sup>1)</sup> Note this is not same person as E. Hlawka!