Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 38 (1992)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A SURVEY

Autor: Shallit, Jeffrey

Kapitel: 11. Properties of the sequence n (mod 1)

DOI: https://doi.org/10.5169/seals-59489

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 17.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

number. It follows from the results of Kmošek and Shallit cited above that $Z(2^{2^k-1}) \leqslant 2$ for all $k \geqslant 0$.

Borosh and Niederreiter [42] showed that $Z(2^k) \leqslant 3$ for $6 \leqslant k \leqslant 35$.

More recently, Niederreiter [223] proved that Zaremba's conjecture holds for all powers of 2; in fact, we have $Z(2^k) \leqslant 3$ for all $k \geqslant 0$.

Larcher [182, Corollary 2] proved the existence of a constant c, such that for every $n \geqslant 1$ there exists a positive integer $j \leqslant n$, relatively prime to n, such that if

$$j/n = [0, a_1, a_2, \ldots, a_m] ,$$

then

$$\sum_{1 \leqslant i \leqslant m} a_i < c(\log n)(\log \log n)^2 .$$

This is close to the best possible bound $O(\log n)$, which was reportedly conjectured by L. Moser (although I do not know a reference); the bound would be a consequence of Zaremba's conjecture.

For other results connected with Zaremba's conjecture, see the papers of Cusick [63, 66]; Niederreiter [224]; Sander [268]; and Hensley [315].

11. Properties of the sequence $n\theta$ (mod 1)

If θ is a real number, by θ (mod 1) we mean $\{\theta\} = \theta - [\theta]$, the fractional part of θ.

It has been known at least since Bernoulli [26] that properties of the sequence $\theta, 2\theta, 3\theta, \ldots$ are intimately connected with the continued fraction expansion for θ. The distribution of $n\theta$ (mod 1) is a vast subject, and we restrict ourselves to mentioning several results connected with numbers of constant type.

Let θ be an irrational number, and let

$$0 = a_0 < a_1 < a_2 < \cdots < a_n < a_{n+1} = 1$$

be the sequence of points $\{k\theta\}$, $1 \leqslant k \leqslant n$, arranged in ascending order. Define

$$\delta_\theta(n) = \max_{1 \leqslant i \leqslant n+1} a_i - a_{i-1} .$$

Then Graham and van Lint [119] proved the following theorem:

$$\limsup_{n\to\infty} n\delta_\theta(n) < \infty$$

if and only if θ is a number of constant type.

Boyd and Steele [43] introduced the function $l_n^+(\theta)$, the length of the longest increasing subsequence of $\{\theta\}, \{2\theta\}, \ldots, \{n\theta\}$. They proved that

$$\liminf_{n\to\infty} \frac{l_n^+(\theta)}{\sqrt{n}} > 0$$

and

$$\limsup_{n\to\infty} \frac{l_n^+(\theta)}{\sqrt{n}} < \infty$$

if and only if the partial quotients of θ are bounded.

For some other results on $\{n\theta\}$ connected with bounded partial quotients, see Ennola [100, 101]; Lesca [185]; Drobot [92]; and Strauch [288].

12. Discrepancy and Dispersion

Let $\omega = (x_1, x_2, x_3, \ldots)$ be a sequence of real numbers. Let $I \subseteq [0, 1)$ be an interval and let $|I|$ denote its length. Define the counting function $S_n(I) = S_n(I, \omega)$ as the number of terms x_k, $1 \leqslant k \leqslant n$, for which $\{x_k\} \in I$.

The *discrepancy* $D_n(x_1, x_2, \ldots, x_n)$ is a measure of how much the sequence $x_1, x_2, \ldots, x_n$ deviates from a uniform distribution. It is defined as follows:

$$D_n(\omega) = D_n(x_1, x_2, \ldots, x_n) = \sup_{I \subseteq [0,1)} \left| \frac{S_n(I, \omega)}{n} - |I| \right|.$$

Now consider the discrepancy of the sequence $\omega = (\theta, 2\theta, 3\theta, \ldots)$. If θ has bounded partial quotients, then the discrepancy of ω is small. In particular, we have the following estimate: If $K(\theta) \leqslant k$, then

$$nD_n(\omega) \leqslant 3 + \left(\frac{1}{\log \alpha} + \frac{k}{\log(k+1)} \right) \log n$$

for $\alpha = \frac{1}{2}(1 + \sqrt{5})$. See, for example, Kuipers and Niederreiter [173].