

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	38 (1992)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A SURVEY
Autor:	Shallit, Jeffrey
Kapitel:	12. Discrepancy and Dispersion
DOI:	https://doi.org/10.5169/seals-59489

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Then Graham and van Lint [119] proved the following theorem:

$$\limsup_{n \rightarrow \infty} n\delta_\theta(n) < \infty$$

if and only if θ is a number of constant type.

Boyd and Steele [43] introduced the function $l_n^+(\theta)$, the length of the longest increasing subsequence of $\{\theta\}, \{2\theta\}, \dots, \{n\theta\}$. They proved that

$$\liminf_{n \rightarrow \infty} \frac{l_n^+(\theta)}{\sqrt{n}} > 0$$

and

$$\limsup_{n \rightarrow \infty} \frac{l_n^+(\theta)}{\sqrt{n}} < \infty$$

if and only if the partial quotients of θ are bounded.

For some other results on $\{n\theta\}$ connected with bounded partial quotients, see Ennola [100, 101]; Lesca [185]; Drobot [92]; and Strauch [288].

12. DISCREPANCY AND DISPERSION

Let $\omega = (x_1, x_2, x_3, \dots)$ be a sequence of real numbers. Let $I \subseteq [0, 1]$ be an interval and let $|I|$ denote its length. Define the counting function $S_n(I) = S_n(I, \omega)$ as the number of terms x_k , $1 \leq k \leq n$, for which $\{x_k\} \in I$.

The *discrepancy* $D_n(x_1, x_2, \dots, x_n)$ is a measure of how much the sequence x_1, x_2, \dots, x_n deviates from a uniform distribution. It is defined as follows:

$$D_n(\omega) = D_n(x_1, x_2, \dots, x_n) = \sup_{I \subseteq [0, 1]} \left| \frac{S_n(I, \omega)}{n} - |I| \right|.$$

Now consider the discrepancy of the sequence $\omega = (\theta, 2\theta, 3\theta, \dots)$. If θ has bounded partial quotients, then the discrepancy of ω is small. In particular, we have the following estimate: If $K(\theta) \leq k$, then

$$nD_n(\omega) \leq 3 + \left(\frac{1}{\log \alpha} + \frac{k}{\log(k+1)} \right) \log n$$

for $\alpha = \frac{1}{2}(1 + \sqrt{5})$. See, for example, Kuipers and Niederreiter [173].

For other results connecting discrepancy and the boundedness of the partial quotients, see the papers of Niederreiter [218] and Dupain and Sós [94, 95]. Also see Beck and Chen [25] and Richert [258].

We can also consider the so-called L^2 discrepancy, T_n , defined as follows: let

$$R_n(t) = \frac{S_n([0, t), \omega)}{n} - t$$

and put

$$T_n(\omega) = \left(\int_0^1 R_n^2(t) dt \right)^{1/2}.$$

It is possible to generalize the definitions of D_n and T_n to the multi-dimensional case, though we omit the details. By appealing to numbers with bounded partial quotients, Davenport [73] constructed sequences in two dimensions with low L^2 discrepancy. Also see Proinov [250, 251, 252].

Another measure connected with sequences is called *dispersion*. Let $\omega = (x_1, x_2, \dots)$ and define the dispersion

$$d_n(\omega) = \sup_{x \in [0, 1]} \min_{1 \leq k \leq n} |x - x_k|,$$

essentially half the distance between the most widely separated points of the sequence x_1, x_2, \dots, x_n . (Compare with the function $\delta_\theta(n)$ in Section 11.)

Niederreiter [221] considered the dispersion of the sequence $\{n\theta\}$. He showed that if θ has bounded partial quotients, then $d_n(\omega) = O(1/n)$. He also gave a more detailed estimate, showing that $d_n(\omega)$ is approximately $K(\theta)/4n$. Also see Drobot [93] and Larcher [311].

13. CONNECTIONS WITH ERGODIC THEORY

Let θ be irrational, $\omega = (\theta, 2\theta, \dots)$ and $S_n(I, \omega)$ be defined as in the previous section. Veech [293] developed connections between S_n and ergodic theory. We mention one result that is number-theoretic in nature. Let $x_n = S_n(I, \omega) \bmod 2$, and define

$$\mu_\theta(I) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{1 \leq k \leq n} x_k,$$