

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 38 (1992)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REAL NUMBERS WITH BOUNDED PARTIAL QUOTIENTS: A SURVEY
Autor: Shallit, Jeffrey
Kapitel: 13. Connections with Ergodic Theory
DOI: <https://doi.org/10.5169/seals-59489>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

For other results connecting discrepancy and the boundedness of the partial quotients, see the papers of Niederreiter [218] and Dupain and Sós [94, 95]. Also see Beck and Chen [25] and Richert [258].

We can also consider the so-called L^2 discrepancy, T_n , defined as follows: let

$$R_n(t) = \frac{S_n([0, t), \omega)}{n} - t$$

and put

$$T_n(\omega) = \left(\int_0^1 R_n^2(t) dt \right)^{1/2}.$$

It is possible to generalize the definitions of D_n and T_n to the multi-dimensional case, though we omit the details. By appealing to numbers with bounded partial quotients, Davenport [73] constructed sequences in two dimensions with low L^2 discrepancy. Also see Proinov [250, 251, 252].

Another measure connected with sequences is called *dispersion*. Let $\omega = (x_1, x_2, \dots)$ and define the dispersion

$$d_n(\omega) = \sup_{x \in [0, 1]} \min_{1 \leq k \leq n} |x - x_k|,$$

essentially half the distance between the most widely separated points of the sequence x_1, x_2, \dots, x_n . (Compare with the function $\delta_\theta(n)$ in Section 11.)

Niederreiter [221] considered the dispersion of the sequence $\{n\theta\}$. He showed that if θ has bounded partial quotients, then $d_n(\omega) = O(1/n)$. He also gave a more detailed estimate, showing that $d_n(\omega)$ is approximately $K(\theta)/4n$. Also see Drobot [93] and Larcher [311].

13. CONNECTIONS WITH ERGODIC THEORY

Let θ be irrational, $\omega = (\theta, 2\theta, \dots)$ and $S_n(I, \omega)$ be defined as in the previous section. Veech [293] developed connections between S_n and ergodic theory. We mention one result that is number-theoretic in nature. Let $x_n = S_n(I, \omega) \bmod 2$, and define

$$\mu_\theta(I) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{1 \leq k \leq n} x_k,$$

if the limit exists. Then Veech showed that $\mu_\theta(I)$ exists for all $I \subseteq [0, 1)$ if and only if the partial quotients of θ are bounded.

For other connections with ergodic theory, see the papers of Stewart [286]; del Junco [154]; Dani [70, 72]; and Baggett and Merrill [14, 15].

14. PSEUDO-RANDOM NUMBER GENERATION

Lehmer [183] introduced the *linear congruential method* for pseudo-random number generation. Let X_0, m, a, c be given, and define

$$X_{k+1} = aX_k + c \pmod{m},$$

for $k \geq 0$. For this to be a good source of “random” numbers, we want the sequence X_k to be uniformly distributed, as well as the sequence of pairs (X_k, X_{k+1}) , triples, etc.

A test for randomness called the *serial test* on pairs (X_k, X_{k+1}) amounts to the two-dimensional version of the discrepancy mentioned above in Section 12. This turns out to be essentially the function $\rho(\mathbf{g}, m)$ defined in Section 10. Thus linear congruential generators that pass the pairwise serial test arise from rationals a/m having small partial quotients in their continued fraction expansion. See the papers of Dieter [87, 88]; Niederreiter [219, 220, 222]; Knuth [170, Section 3.3.3]; and Borosh and Niederreiter [42].

15. FORMAL LANGUAGE THEORY

Let $w = w_0w_1w_2 \cdots$ be an infinite word over a finite alphabet. We say that the finite word $x = x_0x_1 \cdots x_n$ is a *subword* of w if there exists $m \geq 0$ such that $w_{m+i} = x_i$, for $0 \leq i \leq n$. We say that w is *k-th power free* if x^k is never a subword of w , for all nonempty words x . Here is a classical example: let $s(n)$ denote the number of 1's in the binary expansion of n . Then the infinite word of Thue-Morse

$$t = t_0t_1t_2 \cdots = 0110100110010110 \cdots,$$

defined by $t_n = s(n) \pmod{2}$, is cube-free.

Another way to define infinite words is as the fixed point of a homomorphism on a finite alphabet. For example, the Thue-Morse word t is a fixed point of φ , where $\varphi(0) = 01$ and $\varphi(1) = 10$.