Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 39 (1993)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PONCELET'S THEOREM AND DUAL BILLIARDS

Autor: Tabachnikov, Serge

DOI: https://doi.org/10.5169/seals-60421

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025


ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

PONCELET'S THEOREM AND DUAL BILLIARDS

by Serge TABACHNIKOV

While in captivity as a prisoner of war in the Russian city of Saratov, from winter 1812-13 till June 1814, Jean-Victor Poncelet discovered his celebrated closure theorem. Its statement is as follows. Given two nested ellipses Γ_0 , Γ_1 in the plane, one plays the game illustrated in figure 1: choose a point x on Γ_1 , draw a tangent line to Γ_0 from x, find its intersection y with Γ_1 and iterate, taking y as a new starting point. The claim is that, if x returns back to the initial position after a finite number of iterations, then every point of Γ_1 will return back after the same number of iterations.

The prehistory of Poncelet's theorem is related to works of many mathematicians (Euler and Steiner among them). The original proof, given by Poncelet, was synthetic and rather complicated. Jacobi soon recognized a relation between Poncelet's theorem and the theory of elliptic functions, and published his analytic proof in 1828. In 1853 Cayley gave explicit conditions for Poncelet polygons to be closed. A modern conceptual proof of the theorem and its generalization to 3-space were found by Griffith and Harris some 15 years ago. We refer to [B.-K.-O.-R.] and [B], chap. 16.6 for the history of Poncelet's theorem and its classical proofs; and to [G.-H. 1, 2] for the modern one.






Fig.1

Fig.2

Here we give still another proof of Poncelet's theorem. Given two nested ellipses Γ_0 , Γ_∞ , we define an area form in the annulus A between them (so that the total area of A is infinite), and an area-preserving map T of A into itself. This map is integrable in the sense that A is foliated by T-invariant curves Γ_λ . A standard argument shows then that each Γ_λ carries an affine structure and T, restricted to Γ_λ , is a shift $t \to t + c$. It follows that, if T has an N-periodic orbit on Γ_λ , then all its orbits are N-periodic.

To define the area form and the map T identify the interior of Γ_{∞} with the hyperbolic plane (the Klein-Beltrami model). The distance between points is given by $\operatorname{dist}(x, y) = |\log[x, y, b, a]|$, where [] denotes the cross-ratio — see figure 2. The area form is the hyperbolic area, and the map T is the dual billiard transformation with respect to Γ_0 , defined as follows (see [M], [T1,2] for more information on this map). Given a point x outside of Γ_0 , draw a tangent line to Γ_0 through x (say, the right one from the view-point of x) and reflect x in the point of tangency W — see figure 3. Thus, $\operatorname{dist}(x, W) = \operatorname{dist}(W, Tx)$, where the distances are those in the hyperbolic metric.


We need two properties of the dual billiard map.

PROPOSITION 1. For any convex smooth curve Γ_0 (not necessarily an ellipse) the map T preserves hyperbolic area.

Two conics Γ_0 and Γ_∞ determine a pencil of conics Γ_λ , passing through the four intersection points $\Gamma_0 \cap \Gamma_\infty$ (imaginary in our case). The curves Γ_λ foliate the annulus A.

PROPOSITION 2. The map T preserves each ellipse of the pencil in A.

We will prove the propositions a little later; now we deduce Poncelet's theorem from them.








Fig.4

Recall that an affine structure on a curve is an atlas on it such that changes of charts are local affine transformations of the real line. Recall also that given a plane domain with an area form ω , a smooth function f in the domain has the symplectic gradient v associated to it (it is also called the Hamiltonian vector field with the Hamiltonian function f). In local coordinates (p, q) such


that
$$\omega = dp \wedge dq$$
, one has $v = -\frac{\partial f}{\partial q} \frac{\partial}{\partial p} + \frac{\partial f}{\partial p} \frac{\partial}{\partial q}$.

Given two initial ellipses Γ_0 , Γ_1 , include them into a pencil and take Γ_∞ to be an ellipse from this pencil outside of Γ_1 . Consider the dual billiard map in the annulus A between Γ_0 and Γ_∞ . Then A is foliated by invariant curves, and Γ_1 is one of them. Define an affine structure on Γ_1 . Let f be a smooth function in a neighbourhood of Γ_1 , which is constant on each curve Γ_λ . Its symplectic gradient v is tangent to Γ_1 . If one replaces f by another function $g = \phi(f)$, then the corresponding vector field on Γ_1 becomes $u = \phi'(f(\Gamma_1))v$. Hence a vector field is defined on Γ_1 up to a multiplication by a constant. Fix v requiring that its time-one map is the identity. Let t mod 1 be the coordinate on Γ_1 , such that $v = \partial/\partial t$. The parameter t is defined up to a shift $t \to t + a$ and it determines an affine structure on Γ_1 . Since T preserves the area form and the foliation Γ_λ leaf-wise, it is an affine transformation on Γ_1 . Its degree is one, hence it is a shift $t \to t + c$. Poncelet's theorem follows.

The above argument is a particular case of a more general well-known consideration. Given a Lagrangian foliation \mathcal{F} on a symplectic manifold (M, ω) , its leaves carry a canonical affine structure. This structure is defined on a leaf F by a locally free action of the additive group of the cotangent space $T_F^*(M/\mathcal{F})$, where M/\mathcal{F} is the (locally defined) space of leaves of the foliation. Namely, functions on M/\mathcal{F} , considered as Hamiltonians on M, define there commuting vector fields which are tangent to the leaves of \mathcal{F} . A function with zero differential at $F \in M/\mathcal{F}$ defines the zero field on F. If in addition a symplectic map is given, preserving \mathcal{F} leaf-wise, then its restriction to each leaf is an affine transformation therein. If, moreover, the leaves are tori, then the restriction of the map to each leaf is a shift (Arnold-Liouville theorem; see, e.g. [A.-G.]).

Now we prove the two propositions.

LEMMA 3. Given a smooth convex curve Γ in the hyperbolic plane, fix a number c > 0 and consider the family of straight lines which cut off segments of areas c from Γ . Then the envelope of this family bisects each segment of the family at the point of tangency (see figure 4).



Proof (see [F.-T.] for the Euclidean case). Let AB be a line from the family and O its tangency point to the envelope. Assume that AO > BO — see figure 5. Let A_1B_1 be a sufficiently close line from the family and $O' = AB \cap A_1B_1$. We have: Area $(AO'A_1) = \text{Area}(BO'B_1)$. Also AO' > BO' and $A_1O' > B_1O'$. Then the central symmetry in O' sends the "triangle" $BO'B_1$ inside $AO'A_1$; a contradiction.

Proposition 1 follows: the shaded areas in figure 6 are equal.

Consider now a pencil of conics Γ_{λ} and let l be a line in the plane. Intersections of l with the conics Γ_{λ} define an involution on l — see figure 7.

LEMMA 4 (Desargues' Theorem. [B], 14.2.8.3). This involution is a projective transformation of the line l.

Proof. Applying a projective transformation of the plane, we make the

conics Γ_{λ} concentric. Take the center as the origin of the plane V. Then the pencil consists of conics.

$$\Gamma_{\lambda} = \{ \langle A_{\lambda}(x), x \rangle = 1 \}$$

where $A_{\lambda} = A + \lambda E$ and $A, E: V \to V^*$ are selfadjoint operators. Let l be tangent to Γ_0 at x and u be the tangent vector there. Parametrize l by a real t such that points of l are x + tu.

The intersection $l \cap \Gamma_{\lambda}$ is given by

$$\langle (A + \lambda E) (x + tu), x + tu \rangle = 1$$
.

Since $\langle Ax, x \rangle = 1$ and $\langle Ax, u \rangle = 0$, we simplify to

$$<(A + \lambda E)u, u > t^2 + 2\lambda t < Ex, u > + \lambda < Ex, x > = 0.$$

It follows that

$$\frac{1}{t_1} + \frac{1}{t_2} = -2 \frac{\langle Ex, u \rangle}{\langle Ex, x \rangle}$$

independently of λ ; here t_1 and t_2 are the roots. Hence the correspondence $t_1 \leftrightarrow t_2$ is fractional-linear, that is, projective.

Note, that the two fixed points of the involution are the tangency points of *l* to conics of the pencil (one of them is an ellipse, another- a hyperbola). There are exactly two such tangency points by the Jacobi theorem.

To deduce proposition 2, consider figure 8. The involution sends x to y and U to V, preserving W. Since it preserves the cross-ratio, we have: dist(x, W) = dist(W, y).

One can deduce a little more from these proofs. Consider three nested ellipses from a pencil: Γ'' , Γ' and Γ , and identify the interior of the outer one with the hyperbolic plane. Let T' and T'' be the dual billiard transformations in the hyperbolic plane, associated to Γ' and Γ'' .




Fig.8

PROPOSITION 5. T'T'' = T''T' (see figure 9).

Proof. For any point of the annulus between Γ' and Γ there is an ellipse from the pencil through it. Both maps preserve this ellipse and both are shifts in the affine parameter on it. Since shifts commute, the proposition follows.

We refer to [T3] for a partial converse statement to the proposition.

One can also slightly generalize Poncelet's theorem (this generalization was known to Poncelet too). Consider a number of ellipses from a pencil: Γ , Γ' , Γ'' etc, and let Γ contain all the others. Choose a point $x \in \Gamma$, draw a tangent line to Γ' , find its intersection with Γ , draw a tangent line to Γ'' etc. Then, if x returns back after a number of iterations, any initial point on Γ does. It follows again from the fact that T', T'' etc. are shifts in the affine parameter on Γ .

We conclude with a conjecture. Let a smooth strictly convex curve Γ_0 be given in the plane. Assume that its outer neighbourhood is foliated by convex curves Γ_{λ} , $\lambda \in [0, \epsilon[$. Assume also that, for any line tangent to Γ_0 , the (local) involution, defined on it by its intersections with the curves Γ_{λ} , is a projective transformation. Then the curves Γ_{λ} belong to a pencil of ellipses.

Acknowledgements. I am grateful to M. Berger, E. Ghys, P. de la Harpe and J. Moser for stimulating discussions and to ENS de Lyon for its hospitality.

REFERENCES

- [A.-G.] ARNOLD, V. and A. GIVENTAL. Symplectic Geometry. Dynamical Syst. v.4. Springer (1990).
- [B] BERGER, M. Geometry. Springer (1987).
- [B.-K.-O.-R.] Bos, H., C. Kers, F. Oort and D. Raven. Poncelet's Closure Theorem. Expos. Math. 5 (1987), 289-364.
- [F.-T.] FUCHS, D. and S. TABACHNIKOV. Segments of Equal Areas. Quantum, N2 (1992), translated from Kvant (in Russian).
- [G.-H. 1] Griffith, P. and J. Harris. A Poncelet Theorem in Space. Comm. Math. Helv. 52 (1977), 145-160.
- [G.-H. 2] GRIFFITH, P. and J. HARRIS. On Cayley's Explicit Solution of Poncelet's Porism. L'Enseign. Math. 24 (1978), 31-40.
- [M] Moser, J. Is the Solar System Stable? Math. Intell. 1, N2 (1978).
- [T 1] TABACHNIKOV, S. On the Dual Billiard Problem. Adv. Math., to appear.
- [T 2] Dual Billiards. IHES preprint (1992).
- [T 3] Commuting Dual Billiard Maps. ETH preprint (1993).

Serge Tabachnikov

(Reçu le 5 octobre 1992)

Department of Mathematical Sciences University of Arkansas, 301 SCEN Fayetteville, AR 72701, USA