Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 39 (1993)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ELLIPTIC SPACES II
Autor: Felix, Yves / Halperin, Stephen / Thomas, Jean-Claude
Kapitel: 3. Elliptic spaces
DOI: https://doi.org/10.5169/seals-60412

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

It remains to deduce the inequality (2) from (2.3). If the inequality (2) holds for some power series $h(z)$ it will also hold for $h(z^k)$, at the cost of replacing K by $K^{\frac{1}{2k}}$. By (2.3) we are thus reduced to showing that the power series

$$\sum_{i=0}^{\infty} q_i z^i = \prod_{i=0}^{\infty} (1 + z^i)$$

satisfies (2). But this is an immediate consequence of a theorem of Hardy and Ramanujan [10]. □

COROLLARY OF PROOF. *If G satisfies the hypotheses of Theorem 2.1 (2) then for some $k \in \mathbf{N}$,*

$$G(z) \underset{c}{\gg} \prod_{i=1}^{\infty} [1 + (z^k)^i] . \quad \square$$

3. ELLIPTIC SPACES

In this section we establish the ellipticity of the spaces listed in the introduction.

3.1. *Finite simply connected H-spaces, X.*

Because X is an H-space, $H_*(\Omega X; \mathbf{F}_p)$ is commutative, all p. Since it has finite depth [3; Theorem A] it is elliptic [7; Prop. 3.2]. Hence X is elliptic.

3.2. *Simply connected homogeneous spaces, $G /\!/ H$.*

We may suppose that G is simply connected, and hence elliptic by § 3. The fibration $G \to G/H \to BH$ loops to the fibration $\Omega G \to \Omega(G/H) \to H$ in which $\pi_1(H)$ acts trivially in $H_*(\Omega G; \mathbf{F}_p)$ [1; Lemma 5.1]. Thus we can use the Serre spectral sequence to deduce polynomial growth for $H_*(\Omega(G/H); \mathbf{F}_p)$ from the same property for $H_*(\Omega G; \mathbf{F}_p)$.

3.3. *Fibrations $F \to X \to B$ with F, B elliptic.*

Here all spaces are simply connected and we can apply the Serre spectral sequence to deduce that $H_*(X; \mathbf{Z})$ is concentrated in finitely many degrees, and finitely generated in each. Hence X has the weak homotopy type of a finite CW complex. Loop the fibration $F \to X \to B$ and use the fact that $H_*(\Omega F; \mathbf{F}_p)$ and $H_*(\Omega B; \mathbf{F}_p)$ grow polynomially to deduce the same property for $H_*(\Omega X; \mathbf{F}_p)$.

3.4. Simply connected Poincaré complexes X with $H^*(X;\mathbf{F}_p)$ at most doubly generated.

Suppose $p \neq 2$ and $H = H^*(X;\mathbf{F}_p)$ contains an element of odd degree. Then it has an odd generator α. Using Poincaré duality it is easy to see that there are only three possibilities for the algebra H:

$$H = \Lambda\alpha \quad \text{or} \quad \Lambda\alpha \otimes \Lambda\beta \quad \text{or} \quad \Lambda\alpha \otimes \mathbf{F}_p[\beta]/\beta^k .$$

In each case a simple, classical computation [11] produces $\mathrm{Tor}^H(\mathbf{F}_p, \mathbf{F}_p)$ and shows that it grows polynomially. Since the Eilenberg-Moore spectral sequence converges from $\mathrm{Tor}^H(\mathbf{F}_p, \mathbf{F}_p)$ to $H^*(\Omega X;\mathbf{F}_p)$, $H^*(\Omega X;\mathbf{F}_p)$ also has this property.

In all other cases ($p = 2$ or H concentrated in even degrees) H is a commutative local ring in the classic sense. Because H satisfies Poincaré duality it is a Gorenstein ring. Now a theorem of Wiebe [12; Korollar p. 268] asserts (because H has at most two generators) that H is a polynomial algebra divided by a regular sequence. It is thus easy (and classical [11]) to compute $\mathrm{Tor}^H(\mathbf{F}_p, \mathbf{F}_p)$, and deduce that it grows polynomially. Hence so does $H_*(\Omega X;\mathbf{F}_p)$.

3.5. Simply connected Dupin hypersurfaces E in S^{n+1}.

In [9; Table 2.1] are listed the possibilities for $H_*(E;\mathbf{Z})$. We divide these into three cases, using the notation of [9].

Case (a): E has the same integral homology as S^k or as $S^k \times S^l$.

In this case Poincaré duality shows that E has the same integral cohomology ring as S^k or as $S^k \times S^l$, and we can apply 3.4.

Case (b): E has the rational homotopy type of $A_3(2)$, $A_3(4)$, $A_3(8)$, $A_4(2)$ or $A_6(2)$.

In these cases the calculations of [9; §6] show explicitly that the ring $H^*(E;\mathbf{Z})$ is torsion free and generated by two elements. Thus each $H^*(E;\mathbf{F}_p)$ is doubly generated, and we can apply Wiebe's result as in 3.4.

Case (c): E has the integral homology of $S^k \times S^l \times S^{k+l}$, with $k < l$.

We need, in this case, to recall from [9; §2] that there are linear sphere bundles

$$S^k \to E \xrightarrow{\pi_0} B \quad \text{and} \quad S^l \to E \xrightarrow{\pi_1} B_1$$

with B_0, B_1 simply connected focal submanifolds of S^{n+1}. Moreover if D_0, D_1 denote the corresponding disk bundles with boundary E then $S^{n+1} = D_0 \underset{E}{\cup} D_1$.

Fix $p \geqslant 0$ and consider the Serre spectral sequence for the fibration $S^k \to E \to B_0$ with coefficients in $\mathbf{F}_p$. If this fails to collapse then $H^k(\pi_0): H^k(B_0; \mathbf{F}_p) \to H^k(E; \mathbf{F}_p)$ is surjective. Since $l > k$ it is always true that $H^k(\pi_1)$ is surjective. Choose classes $\alpha \in H^k(B_0; \mathbf{F}_p)$, $\beta \in H^k(B_1; \mathbf{F}_p)$ mapping to the same non-zero class in $H^k(E; \mathbf{F}_p)$. The Mayer-Vietoris sequence for the decomposition $S^{n+1} = D_0 \underset{E}{\cup} D_1$ then gives a class $\gamma \in H^k(S^{n+1}; \mathbf{F}_p)$ restricting to α and β, which is absurd.

Thus the spectral sequence for $S^k \to E \to B_0$ collapses and so $H_*(B_0; \mathbf{F}_p) \cong H_*(S^l \times S^{l+k}; \mathbf{F}_p)$. Using Poincaré duality for B_0 we see that $H^*(B_0; \mathbf{F}_p)$ and $H^*(S^l \times S^{l+k}; \mathbf{F}_p)$ are isomorphic as graded algebras. Thus B_0 is elliptic by 3.4 and E is elliptic by 3.3.

3.6. Simply connected closed manifolds M with a smooth action by a compact Lie group G, having a simply connected codimension one orbit.

Here we may assume G is connected. Let the orbit be G/K, and convert the inclusion of G/K into a fibration $F \to G/K \to M$. From [9; Table 1.5] we see that for any p, $\dim H_i(F; \mathbf{F}_p) \leqslant 2$, all i. Thus applying the Serre spectral sequence to the fibration $\Omega(G/K) \to \Omega M \to F$ and using 3.1 for G/K we see that $H_*(\Omega M; \mathbf{F}_p)$ grows polynomially.

3.7. Simply connected manifolds $M \# N$ with each of the rings $H^*(M; \mathbf{Z})$, $H^*(N; \mathbf{Z})$ generated by a single class.

By Van Kampen's theorem both M and N are simply connected, and so their fundamental cohomology classes are not torsion. Since each ring is monogenic, $H^*(M; \mathbf{Z})$ and $H^*(N; \mathbf{Z})$ are torsion free. Thus $H^*(M; \mathbf{F}_p)$ and $H^*(N; \mathbf{F}_p)$ are also monogenic, and so $H^*(M \# N; \mathbf{F}_p)$ is doubly generated. Now apply 3.4.

REFERENCES

[1] BROWDER, W. On differential Hopf algebras. *Trans. Amer. Math. Soc. 107* (1963), 153-176.

[2] FELIX, Y., S. HALPERIN and J.-C. THOMAS. The homotopy Lie algebra for finite complexes. *Publ. de l'Institut des Hautes Etudes Scientifiques 56* (1983), 387-410.

[3] FELIX, Y., S. HALPERIN, J.-M. LEMAIRE and J.-C. THOMAS. Mod p loop space homology. *Invent. Math. 95* (1989), 247-262.

[4] FELIX, Y., S. HALPERIN and J.-C. THOMAS. Hopf algebras of polynomial growth. *J. of Algebra 125* (1989), 408-417.

[5] FELIX, Y., S. HALPERIN and J.-C. THOMAS. Loop space homology of spaces of *LS* category one and two. *Math. Ann. 287* (1990), 377-387.