

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	39 (1993)
<b>Heft:</b>	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	SMOOTHLY EMBEDDED 2-SPHERES AND EXOTIC 4-MANIFOLDS
<b>Autor:</b>	Lisca, Paolo
<b>Kapitel:</b>	2. The manifolds $M(k,l, m)$
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-60424">https://doi.org/10.5169/seals-60424</a>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 14.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

2. THE MANIFOLDS  $M(k, l, m)$ 

Gompf studied in [G] smooth simply connected minimal elliptic surfaces without multiple fibers. These are classified up to diffeomorphism by a positive integer. For each  $n > 0$  there is such a manifold  $V_n \rightarrow \mathbf{CP}^1$  with no multiple fibers,  $6n$  cusp fibers and a section with self-intersection  $-n$  which hits each fiber transversally once. Gompf defines  $N_n$ , the “nucleus” of  $V_n$ , to be a regular neighborhood of a cusp fiber together with the section. Performing a “differentiable logarithmic transform” of multiplicity  $p \geq 0$  inside  $N_n$  gives an elliptic surface  $V_n(p)$  with a multiple fiber, and corresponding nucleus  $N_n(p)$ . In [G] it is proved that  $V_n(0)$  decomposes as a connected sum of  $\pm \mathbf{CP}^2$ , and how this implies that for any fixed odd  $n \geq 3$   $N_n(0)$  and  $N_n$  are homeomorphic but non-diffeomorphic. The manifolds  $N_n(0)$  and  $N_n$  have the following handlebody descriptions:




FIGURE 3

As pointed out in [G], these examples can be generalized. In fact, let  $k, l, m$  be positive integers, and  $M(k, l, m)$  the manifold described in the introduction. The intersection form of  $M(k, l, m)$  is  $\begin{pmatrix} 0 & 1 \\ 1 & -k \end{pmatrix}$ , so  $\partial M(k, l, m)$  is a homology sphere. Moreover  $\partial M(k, l, m)$  is independent of the order of  $k, l, m$  (see below). Hence by [F] and the classification of the symmetric unimodular forms, if either  $s > 0$  or  $s = 0$  and  $k \equiv l \equiv m \pmod{2}$ , the homeomorphism type of  $M(k, l, m) \#^s \overline{\mathbf{CP}}^2$  is independent of the order of  $k, l, m$ . Analogously, it is possible to permute any pair of integers with the same parity without altering the homeomorphism type of  $M(k, l, m)$ . So, for instance,  $M(1, n, 1)$  and  $M(n, 1, 1)$  are homeomorphic for  $n$  odd, and since clearly  $N_n(0) = M(1, n, 1)$  and  $N_n = M(n, 1, 1)$ , they are not diffeomorphic for  $n \geq 3$ .

Let us prove that  $\partial M(k, l, m)$  is independent of the order of  $k, l, m$ . As observed in the introduction,  $M(k, l, m) = M(k, m, l)$  so it is enough to see that  $\partial M(k, l, m) = \partial M(m, l, k)$ . In fact,  $\partial M(k, l, m)$  is given by the first diagram of figure 4 after cancelling the 1-2 pairs, and similarly  $\partial M(m, l, k)$  is given by the second one. Moreover, there is an obvious isotopy between the two diagrams.




FIGURE 4

We introduce now an auxiliary result, needed for the proof of theorem 1.1. If  $q: \Lambda \rightarrow \mathbf{Z}$  is a quadratic form on the lattice  $\Lambda$ , and if  $\alpha \in \Lambda$ ,  $q(\alpha) = -1$ , then

$$R_\alpha(x) = x + 2(x \cdot \alpha)\alpha$$

is an integral isometry of  $\Lambda$ . Similarly, if  $q(\alpha) = -2$ , let

$$R_\alpha(x) = x + (x \cdot \alpha)\alpha.$$

Given an oriented 4-manifold  $M$ , let  $q_M: H^2(M) \rightarrow \mathbf{Z}$  its intersection form. Then there is the following elementary fact:

**PROPOSITION 2.1** (2.4, chapter III in [FM1]). *Let  $M$  be an oriented 4-manifold and  $S^2 \subseteq M$  be an embedded sphere with  $\alpha \in H^2(M; \mathbf{Z})$  the cohomology class dual to  $S^2$ . If  $q_M(\alpha) = -1$  or  $-2$ , there is an orientation-preserving diffeomorphism  $\varphi$  of  $M$  such that  $\varphi^* = R_\alpha$ .*

We shall use the notations  $\hat{N}_k = N_k \#^{l+m-2} \overline{\mathbf{CP}}^2$ ,  $\bar{N}_k = \hat{N}_k \#^s \overline{\mathbf{CP}}^2$ ,  $\hat{V}_k = V_k \#^{l+m-2} \overline{\mathbf{CP}}^2$ ,  $\bar{V}_k = \hat{V}_k \#^s \overline{\mathbf{CP}}^2$ , and  $\bar{M}(k, l, m) = M(k, l, m) \#^s \overline{\mathbf{CP}}^2$ .

**LEMMA 2.2.** *For all  $k, l, m > 0$  there is a smooth embedding  $i: M(k, l, m) \hookrightarrow \hat{N}_k$ , such that  $i_* H_2(M(k, l, m)) \subseteq H_2(N_k) \subseteq H_2(\hat{N}_k)$ .*

*Proof.* Blow-up  $l+m-2-1$ 's around the 0-framed knot in a link picture for  $N_k$  to get a framed link containing figure 1 plus algebraically unlinked — 1-framed unknots. QED

*Remark 2.3.* Since  $N_k \subset V_k$ , by the lemma

$$M(k, l, m) \hookrightarrow \hat{N}_k \subset \hat{V}_k, \quad \text{and} \quad \bar{M}(k, l, m) \hookrightarrow \bar{V}_k.$$

It is easy to see from the proof of the lemma that the image of  $H_2(M(k, l, m))$  under the above embeddings is orthogonal to the Poincaré duals of the exceptional classes, and it contains the class  $[f]$  of a smooth fiber of the elliptic fibration on  $\hat{V}_k$ . In fact  $[f]$  comes from the class given by the 2-handle attached to the 0-framed component in the link picture (see [G]).

**THEOREM 2.4.** *Let  $l, m$  be positive integers with  $l > 2$ , and  $s$  a non-negative integer. If either (i)  $s \neq 0$  or (ii)  $s = 0$  and  $l \equiv 0 \pmod{2}$ , the manifolds  $\bar{M}(2, l, m)$  and  $\bar{M}(l, 2, m)$  are homeomorphic but their interiors are not diffeomorphic.*

*Proof.* Suppose we are in case (i). Then, as pointed out above,  $\bar{M}(2, l, m)$  and  $\bar{M}(l, 2, m)$  are homeomorphic for any  $l$ .  $\bar{M}(2, l, m)$  contains an embedded sphere  $S^2$  of square  $-2$ : take a slicing disk for the  $-2$ -framed unknot in the link picture for  $M(2, l, m)$  union the core of the corresponding 2-handle. By contradiction, suppose  $\psi: \text{int}(\bar{M}(2, l, m)) \cong \text{int}(\bar{M}(l, 2, m))$  is a diffeomorphism. By remark 2.3  $\bar{M}(l, 2, m) \subset \bar{V}_l$ , so  $\psi(S^2) \subset \bar{V}_l$  is an embedded sphere. If  $\psi$  were orientation-reversing  $\psi(S^2)$  would have positive self-intersection. But for  $l \geq 2$   $\bar{V}_l$  does not contain such embedded spheres [FM2], hence  $\psi$  has to be orientation-preserving. So  $\psi(S^2)$  has self-intersection  $-2$ . Let  $\alpha \in H^2(\bar{V}_l)$  be the Poincaré dual of the homology class carried by  $\psi(S^2)$ . By proposition 2.1 there is an orientation-preserving self-diffeomorphism  $\varphi$  of  $\bar{V}_l$  such that  $\varphi^*(\xi) = \xi + (\xi, \alpha)\alpha$  for any  $\xi \in H^2(\bar{V}_l)$ . This implies that  $\alpha$  is orthogonal to the exceptional classes of  $\bar{V}_l$ , because by [FM2] if  $e$  is an exceptional class then  $\varphi^*(e) = e$ . Therefore  $[\psi(S^2)] \in H_2(M(l, 2, m)) \subset H_2(\hat{V}_l)$ . Observe that  $M(l, 2, m)$  has intersection form  $\begin{pmatrix} 0 & 1 \\ 1 & -l \end{pmatrix}$ , where the class of square zero may be taken to be the class of a smooth fiber  $f$ , hence  $[\psi(S^2)] \cdot [f] \neq 0$ . Hence  $\alpha \cdot k_{\bar{V}_l} \neq 0$ , where  $k_{\bar{V}_l}$  is the canonical class. Again by [FM2] any orientation-preserving self-diffeomorphism of  $\bar{V}_l$  preserves the canonical class  $k_{V_l}$  of  $V_l$  up to sign and the exceptional classes up to permutation and signs. But since  $\alpha$  is not a multiple of  $k_{V_l}$ ,  $\varphi$  cannot have this property. This gives a contradiction to the existence of  $\psi$  and proves the theorem in case (i). To prove the statement in case (ii) it is enough to observe that since, by what we have proved

already, the interiors of  $M(2, l, m) \# \overline{\mathbf{CP}}^2$  and  $M(l, 2, m) \# \overline{\mathbf{CP}}^2$  are not diffeomorphic, the interiors of  $M(2, l, m)$  and  $M(l, 2, m)$  cannot be diffeomorphic as well. QED

**THEOREM 2.5.** *Let  $l, m$  be positive integers with  $l > 1$ , and  $s$  a non-negative integer. If either (i)  $s \neq 0$  or (ii)  $s = 0$  and  $l \equiv 1 \pmod{2}$ , then the manifolds  $\bar{M}(1, l, m)$  and  $\bar{M}(l, 1, m)$  are homeomorphic but their interiors are not diffeomorphic.*

*Proof.* Use a sphere of square  $-1$  inside  $\bar{M}(1, l, m)$  to get a contradiction to the existence of a diffeomorphism exactly as in the proof of theorem 2.4. QED

*Proof of theorem 1.1.* Put theorems 2.4 and 2.5 together. QED

Now we give our proof of theorem 1.2.

*Proof of theorem 1.2.* In [A] it is proved that  $-W_1$  and  $-W_2$  are homeomorphic. To prove that they are not diffeomorphic, notice first that  $W_1$  can be embedded inside  $\hat{V}_k = V_k \# 2\overline{\mathbf{CP}}^2$  for any  $k \geq 1$ : blow-up two  $-1$ 's around the 0-framed knot in a link picture for  $N_k$ . This embedding sends a generator of  $H_2(W_1)$  to  $f + e \in H_2(\hat{V}_k)$ , where  $f$  is the class of a smooth fiber, and  $e$  is the Poincaré dual of an exceptional class. Next observe that a generator of  $H_2(W_2)$  is representable by a sphere  $S^2$  smoothly embedded in  $\text{int}(W_2)$  with self-intersection  $-1$ : the knot  $K_2$  is ribbon, hence slice, so take as  $S^2$  a slicing disk union the core of the corresponding 2-handle. Finally suppose, arguing by contradiction, that  $\psi: \text{int}(W_2) \cong \text{int}(W_1)$  is a diffeomorphism. The class  $f + e$  is therefore representable by the smoothly embedded sphere  $\psi(S^2)$  inside  $\hat{V}_k$ . Since  $\hat{V}_k$  does not contain embedded spheres with positive self-intersection [FM2], we may assume that  $\psi$  is orientation preserving and  $\psi(S^2)$  has self-intersection  $-2$ . Again by [FM2],  $V_k$  has big diffeomorphism group with respect to the canonical class, and it has some non-zero generalized Donaldson invariant. Moreover  $b_2^+(\hat{V}_k) \geq 3$  for  $k \geq 2$ . So when  $k \geq 2$  we may apply theorem 3.5 from [FM2], which implies that any class of square  $-1$  in  $H_2(\hat{V}_k)$  representable by a smoothly embedded 2-sphere is equal, up to sign, to the Poincaré dual of one of the exceptional classes. This is clearly not the case for  $f + e$ , so we get a contradiction to the existence of  $\psi$ . QED