

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 39 (1993)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REMARKS AND PROBLEMS ON FINITE AND PERIODIC CONTINUED FRACTIONS
Autor: Mendès France, Michel
Kapitel: §4. Related problems
DOI: <https://doi.org/10.5169/seals-60426>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

since it is b , the smallest of the two integers a and b that seems to control the behavior of $(a/b)^n$. It is also to be noticed that in Dixon's inequality, the $\ln a$ that appears on the left hand side could well be replaced by $\ln b$ since for "almost all" couples a, b , $\ln a \approx \ln b$.

Numerical evidence supports equality (1). Based on computer computation, Chr. Batut and M. Olivier showed that

$$\left| \frac{1}{n} \delta \left(\left(\frac{a}{b} \right)^n \right) - \frac{12}{\pi^2} \ln 2 \ln b \right|$$

is less than .02 for n in the range (4000, 5000) and for $a = 3, b = 2$ on the one hand and $a = 5, b = 2$ on the other hand.

§4. RELATED PROBLEMS

My initial (unsuccessful) attempts to prove Theorem 1 were based on the comparison of $\delta(ax/b)$ to $\delta(x)$. I was hoping that a relationship between both depths would give by induction some results on $\delta(xa^n/b^n)$. This turned out nonconclusive, yet I did obtain some results which I believe are interesting in themselves [10].

Let a, b, c, d be coprime integers and let $\Delta = |ad - bc|$. Consider the Möbius map

$$x \mapsto Tx = \frac{ax + b}{cx + d}.$$

THEOREM 2.

$$\limsup_{\delta(x) \rightarrow \infty} \frac{\delta(Tx)}{\delta(x)} = \Theta(\Delta)$$

where Θ takes odd integral values. As n increases to infinity $\Theta(n)$ behaves like $\ln n$. More precisely let

$$\alpha = \left(2 \ln \frac{1 + \sqrt{5}}{2} \right)^{-1}.$$

Then for all integer $n \geq 1$

$$1 + \alpha \ln n \leq \Theta(n) \leq 2(1 + \alpha \ln n).$$

Θ is linked to the depth by the formula

$$\Theta(n) = \max_{1 \leq b \leq n} \delta \left(\frac{b}{n} \right) + 1.$$

Finally

$$\liminf_{\delta(x) \rightarrow \infty} \frac{\delta(Tx)}{\delta(x)} = \frac{1}{\Theta(\Delta)}.$$

The following table gives the first values of Θ

n	1	2	3	4	5	6	7	8	9	10	...
$\Theta(n)$	1	3	3	3	5	3	5	5	5	5	...

Actually Theorem 1 can be improved. There exist two constants $C_1 = C_1(T)$ and $C_2 = C_2(T)$ such that for all rational x

$$\frac{1}{\Theta(\Delta)} \delta(x) - C_1 \leq \delta(Tx) \leq \Theta(\Delta) \delta(x) + C_2.$$

Both inequalities are sharp apart from the exact values of C_1 and C_2 .

5. MORE QUESTIONS

To every Möbius map T we associate the interval $I(T) = [\Theta^{-1}(\Delta), \Theta(\Delta)]$.

PROBLEM 3. *Is it true that for all $\zeta \in I(T)$ there exists a sequence of rational numbers x_n such that $\lim_{n \rightarrow \infty} \delta(x_n) = \infty$ and*

$$\lim_{n \rightarrow \infty} \frac{\delta(Tx_n)}{\delta(x_n)} = \zeta?$$

PROBLEM 4. *Let T_1, T_2, \dots, T_k be Möbius maps with pairwise coprime determinants $\Delta_1, \Delta_2, \dots, \Delta_k$.*

Is it true that for all

$$(\zeta_1, \zeta_2, \dots, \zeta_k) \in \prod_{i=1}^k I(T_i)$$

there exists a sequence of rational x_n with strictly increasing depths such that for all $i = 1, 2, \dots, k$

$$\lim_{n \rightarrow \infty} \frac{\delta(T_i x_n)}{\delta(x_n)} = \zeta_i?$$

Can k be infinite?

The following result should be mentioned at this point.