Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 39 (1993)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TREPREAU'S EXAMPLE, A PEDESTRIAN APPROACH
Autor: Rosay, Jean-Pierre
Bibliographie
DOI: https://doi.org/10.5169/seals-60427

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 06.03.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

VI. TREPREAU DOES MORE

1) Trepreau not only deals with wedge decomposability (which is related to wave front sets in acute cones), he shows in his example that the wave front set of a *CR* function at 0 is either empty or the whole conormal.

2) Our proof may not adapt to a slightly perturbed situation. But this is precisely the point! Let us compare with the theory of elliptic points for surfaces with isolated complex tangencies. The model case $(z_2 = |z_1|^2 + \alpha \operatorname{Re} z_1^2, 0 \leqslant \alpha < 1)$ is totally trivial to explore. Only the perturbed case needs Bishop's disks. We hope that the reader is convinced that the same is true here.

REFERENCES

[1] BAOUENDI, M.S. and L.P. ROTHSCHILD. Normal forms of generic manifolds and holomorphic extension of *CR* functions. *J. Diff. Geom. 25* (1987), 431-467.

[2] BAOUENDI, M.S. and L.P. ROTHSCHILD. Cauchy-Riemann functions on manifolds of higher codimension in complex space. *Inv. Math. 101* (1990), 45-56.

[3] BAOUENDI, M.S. and F. TREVES. A property of the functions and distributions annihilated by a locally integrable system of complex vector fields. *Ann. of Math. 113* (1981), 387-421.

[4] BOGGESS, A. *CR manifolds and the tangential Cauchy-Riemann Complex.* Studies in Adv. Math., CRC Press 1991.

[5] HÖRMANDER, L. *The Analysis of linear Partial Differential Operators I.* Grund. Math. Wis. 256, Springer Verlag (1983).

[6] POLETSKY, E. Holomorphic Currents. *Ind. Math. J. 42* (1993), 85-144.

[7] TREPREAU, J.M. Sur la propagation des singularités dans les variétés *CR*. *Bull. SMF 118* (1990), 403-450.

[8] TREVES, F. *Hypo-Analytic Structures: Local Theory.* Princeton Math. Series 40 (1992).

[9] TUMANOV, A. Extending *CR* functions on a manifold of finite type over a wedge. *Math. Sb. 136* (1988), 129-140.

[10] —— Connections and propagation of analyticity for *CR* functions. Preprint.

[11] WEINSTOCK, B.M. On the polynomial convexity of the union of two maximal totally real subspaces of $\mathbf{C}^n$. *Math. Ann. 282* (1988), 131-138.

(Reçu le 13 mai 1993)

Jean-Pierre Rosay

Department of Mathematics
University of Wisconsin
Madison, WI 53706
(USA)