

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 39 (1993)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MAXIMALLY COMPLETE FIELDS
Autor: Poonen, Bjorn
Kapitel: 3. Mal'cev-Neumann rings
DOI: <https://doi.org/10.5169/seals-60414>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

be embedded in a divisible one, namely its injective hull. Since an ordered group G is necessarily torsion-free, its injective hull \tilde{G} can be identified with the set of quotients g/m with $g \in G$, m a positive integer, modulo the equivalence relation $g/m \sim h/n$ iff $ng = mh$ in G . We make \tilde{G} an ordered group by setting $g/m \geq h/n$ iff $ng \geq mh$ in G . (One can check that this is the unique extension to \tilde{G} of the ordered group structure on G .)

If G is an ordered group, let $G_\infty = G \cup \{\infty\}$ be the ordered monoid containing G in which $g + \infty = \infty + g = \infty$ for all $g \in G_\infty$ and $g < \infty$ for all $g \in G$. As usual, a valuation v on a field F is a function from F to G_∞ satisfying for all $x, y \in F$

- (1) $v(x) = \infty$ iff $x = 0$
- (2) $v(xy) = v(x) + v(y)$
- (3) $v(x + y) \geq \min \{v(x), v(y)\}$.

The *value group* is G . The *valuation ring* A is $\{x \in F \mid v(x) \geq 0\}$. This is a local ring with maximal ideal $\mathcal{M} = \{x \in F \mid v(x) > 0\}$. The *residue field* is A/\mathcal{M} . We refer to the pair (F, v) (or sometimes simply F) as a *valued field*.

3. MAL'CEV-NEUMANN RINGS

This section serves not only as review, but also as preparation for the construction of the next section. Mal'cev-Neumann rings are generalizations of Laurent series rings. For any ring R (all our rings are commutative with 1), and any ordered group G , the Mal'cev-Neumann ring $R((G))$ is defined as the set of formal sums $\alpha = \sum_{g \in G} \alpha_g t^g$ in an indeterminate t with $\alpha_g \in R$ such that the set $\text{Supp } \alpha = \{g \in G \mid \alpha_g \neq 0\}$ is a well-ordered subset of G (under the given order of G). (Often authors suppress the indeterminate and write the sums in the form $\sum \alpha_g g$, as in a group ring. We use the indeterminate in order to make clear the analogy with the fields of the next section.) If $\alpha = \sum_{g \in G} \alpha_g t^g$ and $\beta = \sum_{g \in G} \beta_g t^g$ are elements of $R((G))$, then $\alpha + \beta$ is defined as $\sum_{g \in G} (\alpha_g + \beta_g) t^g$, and $\alpha\beta$ is defined by a "distributive law" as $\sum_{j \in G} \gamma_j t^j$ where $\gamma_j = \sum_{g+h=j} \alpha_g \beta_h$.

LEMMA 1. *Let A, B be well-ordered subsets of an ordered group G . Then*

- (1) *If $x \in G$, then $A \cap (-B + x)$ is finite.*
(We define $-B + x = \{-b + x \mid b \in B\}$.)

- (2) The set $A + B = \{a + b \mid a \in A, b \in B\}$ is well-ordered.
 (3) The set $A \cup B$ is well-ordered.

Proof. See [13]. \square

The lemma above easily implies that the sum defining γ_j is always finite, and that $\text{Supp}(\alpha + \beta)$ and $\text{Supp}(\alpha\beta)$ are well-ordered. Once one knows that the operations are defined, it's clear that they make $R((G))$ a ring.

Define $\nu: R((G)) \rightarrow G_\infty$ by $\nu(0) = \infty$ and $\nu(\alpha) = \min \text{Supp } \alpha$ for $\alpha \neq 0$. (This makes sense since $\text{Supp } \alpha$ is well-ordered.) If $\alpha \in R((G))$ is nonzero and $\nu(\alpha) = g$, we call $\alpha_g t^g$ the *leading term* of α and α_g the *leading coefficient*. If R is a field, then ν is a valuation on $R((G))$, since the leading term of a product is the product of the leading terms.

LEMMA 2. *If $\alpha \in R((G))$ satisfies $\nu(\alpha) > 0$, then $1 - \alpha$ is a unit in $R((G))$.*

Proof. One way of proving this is to show that for each $g \in G$, the coefficients of t^g in $1, \alpha, \alpha^2, \dots$ are eventually zero, so $1 + \alpha + \alpha^2 + \dots$ can be defined termwise. Then one needs to check that its support is well-ordered, and that it's an inverse for $1 - \alpha$. See [13] for this. An easier way [15] is to obtain an inverse of $1 - \alpha$ by successive approximation. \square

COROLLARY 1. *If the leading coefficient of $\alpha \in R((G))$ is a unit of R , then α is a unit of $R((G))$.*

Proof. Let rt^g be the leading term of α . Then α is the product of rt^g , which is a unit in $R((G))$ with inverse $r^{-1}t^{-g}$, and $(rt^g)^{-1}\alpha$, which is a unit by the preceding lemma. \square

COROLLARY 2. *If R is a field, then $R((G))$ is a field.*

So in this case, if we set $K = R((G))$, (K, ν) is a valued field. Clearly the value group is all of G , and the residue field is R . Note that $\text{char } K = \text{char } R$, since in fact, R can be identified with a subfield of K . (We will refer to these fields as being the "equal characteristic" case, in contrast with the p -adic fields of the next section in which the fields have characteristic different from that of their residue fields.) For example, if $G = \mathbf{Z}$, then $R((G))$ is the usual field of formal Laurent series.