

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 39 (1993)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** MAXIMALLY COMPLETE FIELDS  
**Autor:** Poonen, Bjorn  
**Kapitel:** 6. Applications  
**DOI:** <https://doi.org/10.5169/seals-60414>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

(1) Any equation of the form

$$x^{p^n} + a_1 x^{p^{n-1}} + \cdots + a_{n-1}x^p + a_n x + a_{n+1} = 0$$

with coefficients in  $R$  has a root in  $R$ .

(2) The value group  $G$  satisfies  $G = pG$ .

Also if  $G$  is discrete of arbitrary rank and  $\text{char } F = \text{char } R$ , then the extension is unique [6]. But Kaplansky gives examples where the extension is not unique. The exact conditions under which the extension is unique are not known.

## 6. APPLICATIONS

One application of Theorem 2 is to the problem of “glueing” two valued fields. (This result can also be proved directly without the use of Mal’cev-Neumann fields; it is equivalent to Exercise 2 for §2 in Chapter VI of [2]. Our method has the advantage of showing that the value group of the composite field can be contained in any divisible value group large enough to contain the value groups of the fields to be glued.)

**PROPOSITION 8.** *Suppose  $E, F, F'$  are valued fields and that we are given embeddings of valued fields  $\phi: E \rightarrow F, \phi': E \rightarrow F'$ . Then there exist a Mal’cev-Neumann field  $L$  (or  $K$ ) and embeddings of valued fields  $\Phi: F \rightarrow L, \Phi': F' \rightarrow L$  such that  $\Phi \circ \phi = \Phi' \circ \phi'$ .*

*Proof.* By the glueing theorem for ordered groups [14], we can assume the value groups of  $F$  and  $F'$  are contained in a single ordered group  $G$ . Also we can assume that their residue fields are contained in a field  $R$ . Moreover, we may assume  $G$  is divisible and  $R$  is algebraically closed. Then  $E$  can be embedded as a valued subfield of a power series field  $L$  (or  $K$ ) with value group  $G$  and residue field  $R$ , by Corollary 5. Finally, Theorem 2 gives us the desired embeddings  $\Phi, \Phi'$ .  $\square$

*Remark.* Transfinite induction can be used to prove the analogous result for glueing an arbitrary collection of valued fields.

Since a non-archimedean absolute value on a field can be interpreted as a valuation with value group contained in  $\mathbf{R}$ , we can specialize the results of Section 5 to get results about fields with non-archimedean absolute values. For example, Corollary 5 implies the following, which may be considered the non-archimedean analogue of Ostrowski’s theorem that any field with an archimedean absolute value can be embedded in  $\mathbf{C}$  with its usual absolute value (or one equivalent).

PROPOSITION 9. *Let  $(F, ||)$  be a field with a non-archimedean absolute value, and suppose the residue field is contained in the algebraically closed field  $R$ . Define  $K$  and  $L$  as the Mal'cev-Neumann fields with value group  $\mathbf{R}$  and residue field  $R$ . (Define the  $p$ -adic Mal'cev-Neumann field  $L$  only if  $\text{char } R > 0$ .) The valuations on  $K$  and  $L$  induce corresponding absolute values. Then there exists an absolute value-preserving embedding of fields  $\phi: F \rightarrow K$  or  $\phi: F \rightarrow L$ , depending on if the restriction of  $||$  to the minimal subfield of  $F$  is the trivial absolute value (on  $\mathbf{Q}$  or  $\mathbf{F}_p$ ) or the  $p$ -adic absolute value on  $\mathbf{Q}$ .*

Similarly, Proposition 8 above gives a glueing proposition for non-archimedean absolute values. In fact, this result holds for archimedean absolute values as well, in light of Ostrowski's theorem.

## 7. EXAMPLE: THE MAXIMALLY COMPLETE IMMEDIATE EXTENSION OF $\bar{\mathbf{Q}}_p$

For this section,  $(L, \nu)$  will denote the  $p$ -adic Mal'cev-Neumann field having value group  $\mathbf{Q}$  and residue field  $\bar{\mathbf{F}}_p$ . We have a natural embedding of  $\mathbf{Q}_p$  into  $L$ . By Corollary 4,  $L$  is algebraically closed, so this embedding extends to an embedding of  $\bar{\mathbf{Q}}_p$  into  $L$  (which is unique up to automorphisms of  $\bar{\mathbf{Q}}_p$  over  $\mathbf{Q}_p$ .) In fact this embedding is continuous, since there is a unique valuation on  $\bar{\mathbf{Q}}_p$  extending the  $p$ -adic valuation on  $\mathbf{Q}_p$ . Since  $\bar{\mathbf{Q}}_p$  has value group  $\mathbf{Q}$  and residue field  $\bar{\mathbf{F}}_p$ ,  $L$  is an immediate extension of  $\bar{\mathbf{Q}}_p$ . By Corollary 6,  $L$  is in fact the unique maximally complete immediate extension of  $\bar{\mathbf{Q}}_p$ . Also, any valued field  $(F, w)$  of characteristic 0 satisfying

- (1) The restriction of  $w$  to  $\mathbf{Q}$  is the  $p$ -adic valuation;
- (2) The value group is contained in  $\mathbf{Q}$ ;
- (3) The residue field is contained in  $\bar{\mathbf{F}}_p$ ;

can be embedded in  $L$ , by Corollary 5. For example, the completion  $\mathbf{C}_p$  of  $\bar{\mathbf{Q}}_p$  can be embedded in  $L$ . (This could also be proved by noting that  $L$  is complete by Corollary 4.)

We will always use as the set  $S$  of representatives for  $\bar{\mathbf{F}}_p$  the primitive  $k^{\text{th}}$  roots of 1, for all  $k$  not divisible by  $p$ , and 0. Then the elements of  $L$  have the form  $\sum_g \alpha_g p^g$  for some primitive  $k^{\text{th}}$  roots  $\alpha_g$  of 1, where the exponents form a well-ordered subset of  $\mathbf{Q}$ . In particular, the elements of  $\bar{\mathbf{Q}}_p$  can be expressed in this form. This was first discovered by Lampert [9].