Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 39 (1993)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HURWITZ QUATERNIONIC INTEGERS AND SEIFERT FORMS

Autor: Shastri, Parvati

Kapitel: §2. The root System \$D_4\$ and the Hurwitz quaternionic integers

DOI: https://doi.org/10.5169/seals-60415

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

1.3. PROPOSITION. A unimodular symmetric bilinear form S admits a decomposition S = B + B' with B unimodular if and only if S has a perfect isometry.

Thus, Question 1.1 reduces to the following.

1.4. QUESTION. Given a unimodular symmetric bilinear form S, does there exist a perfect isometry of S?

Note that if S is positive definite and even, then the rank of S is a multiple of S. M. Kervaire gives a complete answer to Question 1.4, for positive definite forms of rank less than or equal to 24. For forms of arbitrary rank, he proves the following partial result, using the theory of the associated root systems.

Let $R = \{x \in L \mid S(x, x) = 2\}$. Suppose that R is a root system in \mathbb{R}^n of rank n (= rank L). Then the irreducible components of R are of type A, D, or E; and we have:


- 1.5. THEOREM ([K], Cor. 3, Prop. 4).
- (a) If R has an irreducible component of type A_{2k-1} , E_7 or D_{k+4} , with $k \ge 1$, then there does not exist any perfect isometry of (L, S).
- (b) If $R = \coprod_{1 \le i \le p} A_{2k_i} \perp qE_6 \perp rE_8$, then there exists a perfect isometry
- of L, inducing a perfect isomorphism of the abelian group $\mathbb{Z}R^{\#}/\mathbb{Z}R$, which corresponds to multiplication by -1, where $\mathbb{Z}R^{\#}$ denotes the dual of the lattice $\mathbb{Z}R$.

Note that the case of R having an irreducible component of type D_4 is not covered by this theorem. In this paper we give an analogue of (b) for this case. In fact, we first consider the case in which R is of type nD_4 . In this case, we show (Th. 5.2) that (L, S) admits a perfect isometry if and only if the isometry class of (L, S) contains a symmetric bilinear space (L', S') of some hermitian space over the Hurwitz quaternionic integers. The analogue of Proposition 1.5 follows from this immediately (Theorem 5.3). In the final section we also give some examples.

$\S 2$. The root system D_4 and the Hurwitz quaternionic integers

The fact that the root lattice $\mathbf{Z}D_4$ can be identified with the lattice of Hurwitz quaternionic integers was long recognized: see for instance ([C-S]). However we give here a direct proof of this fact and recall some arithmetical facts about these quaternionic integers, which are needed in the sequel.

We first fix the following terminology. By a **Z**-lattice we mean a pair (L, b), where L is a finitely generated free **Z**-module and $b: L \times L \to \mathbf{Z}$ a positive definite, even, symmetric bilinear form. If the set $\{x \in L \mid b(x, x) = 2\}$ forms a root system of type nD_4 where the rank of L equals 4n, then we call it a **Z**-lattice of type nD_4 . If L is contained in \mathbf{R}^m and b is induced by the Euclidean inner product on \mathbf{R}^m , we call it a Euclidean **Z**-lattice. The symbol D_4 will always mean the root system in \mathbf{R}^4 with the Euclidean inner product, corresponding to the Dynkin diagram



Let $\mathscr{A} = \mathbf{Q} \oplus \mathbf{Q}i \oplus \mathbf{Q}j \oplus \mathbf{Q}k$ denote the quaternion division algebra over \mathbf{Q} , defined by

$$i^2 = j^2 = k^2 = -1$$
, $ij = -ji = k$.

Let $h: \mathcal{A}^n \times \mathcal{A}^n \to \mathcal{A}$ be the hermitian form defined by


$$h((x_1...x_n), (y_1, ..., y_n)) = \sum_{i=1}^{n} x_i \bar{y}_i,$$

where bar denotes the conjugation in \mathscr{A} . If $Tr: \mathscr{A} \to \mathbf{Q}$ denotes the trace map $Tr(x) = x + \bar{x}$, then $Tr \circ h$ is a positive definite symmetric bilinear form over \mathbf{Q} . Let \mathscr{H} denote the Hurwitz quaternionic integers in \mathscr{A} i.e. $\mathscr{H} = \{(a+bi+cj+dk)/2 \mid a,b,c,d \in \mathbf{Z}, \text{ with the same parity}\}$. Then, \mathscr{H} is a maximal order in \mathscr{A} and $(\mathscr{H}, Tr \circ h)$ is a \mathbf{Z} -lattice. It is trivial to verify that $\xi_1 = (1+i+j+k)/2$, $\xi_2 = (1+i+j-k)/2$, $\xi_3 = (1+i-j+k)/2$, and $\xi_4 = (1-i+j+k)/2$ form a \mathbf{Z} -basis of \mathscr{H} . Let \mathscr{H}^* denote the dual of \mathscr{H} in \mathscr{A} . Then we have

2.1. Proposition.

- (a) The **Z**-lattice $(\mathcal{H}, Tr \circ h)$ is isometric to the Euclidean lattice \mathbf{ZD}_4 .
- (b) The group of units of \mathcal{H} forms a root system isomorphic to D_4 .
- (c) Every **Z**-lattice of type nD_4 is isometric to a **Z**-lattice L such that $\mathcal{H}^n \subset L \subset \mathcal{H}^{*n}$, where the bilinear form on L is induced by $Tr \circ h$.

Proof. Let $\{\varepsilon_i\}$ denote the standard orthonormal basis in \mathbb{R}^4 , and let $\alpha_1 = \varepsilon_2 - \varepsilon_3$, $\alpha_2 = \varepsilon_1 - \varepsilon_2$, $\alpha_3 = \varepsilon_3 - \varepsilon_4$, $\alpha_4 = \varepsilon_3 + \varepsilon_4$. Then $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ is a basis for the root system D_4 . The associated Dynkin diagram is given by



If b denotes the Euclidean inner product on \mathbb{R}^4 , then, $\tau \colon \mathcal{H} \to \mathbf{Z}D_4$ defined by $\tau(\xi_1) = \alpha_1$, $\tau(\xi_i) = -\alpha_i$, $2 \le i \le 4$, is an isometry of $(\mathcal{H}, Tr \circ h)$ onto $(\mathbf{Z}D_4, b)$. This proves (a). We note that an element x in \mathcal{H} is a unit if and only if $Tr \circ h(x) = 2$. Hence (b) follows from the above isometry. Since $Tr \circ h$ is nondegenerate, the dual of \mathcal{H} in \mathcal{A} is the same as the dual of \mathcal{H} in $\mathcal{A} \otimes \mathbf{R} \simeq \mathbf{R}^4$. From (a) it follows that \mathcal{H}^* is isometric to $(\mathbf{Z}D_4)^\#$. Thus (c) follows from the fact that every \mathbf{Z} -lattice of type nD_4 is isometric to a Euclidean \mathbf{Z} -lattice L such that $(\mathbf{Z}D_4)^n \subset L \subset (\mathbf{Z}D_4^\#)^n$.

Let us now recall a few arithmetical facts about the Hurwitz quaternionic integers, details of which can be found in [R]. The dual \mathcal{H}^* is a two-sided full \mathcal{H} -module in \mathcal{A} i.e. an \mathcal{H} -submodule of \mathcal{A} which contains a **Q**-basis of \mathcal{A} . The set of all two-sided full \mathcal{H} -submodules of \mathcal{A} is a free abelian group with the set of all maximal ideals of \mathcal{H} as basis. Further the inverse of \mathcal{H}^* is a maximal ideal in \mathcal{H} . In fact, $(\mathcal{H}^*)^{-1} = \mathcal{P}$, $\mathcal{P} = (1+i)$, $\mathcal{P}^2 = (2)$, $\mathcal{P} = \bar{\mathcal{P}}$, and $\mathcal{H}/\mathcal{P} \simeq \mathbf{F}_4$. We have,

2.2. Proposition.

- (a) The quotient $\mathcal{H}^*/\mathcal{H}$ has the natural structure of a vector space of dimension one over \mathbf{F}_4 .
- (b) The hermitian form h induces a hermitian form $\eta(h)$ on $\mathcal{H}^*/\mathcal{H}$, with values in $\mathcal{H}^{*2}/\mathcal{H}^*$, which is isometric to the standard hermitian form on \mathbf{F}_4 .

Proof. (a) This follows from the fact that, \mathcal{H}^* is an \mathcal{H} -module of rank one and $\mathcal{PH}^* = \mathcal{H}^*\mathcal{P} = \mathcal{H}$.

(b) This follows from the commutativity of the diagram:

$$\mathcal{H}^*/\mathcal{H} \times \mathcal{H}^*/\mathcal{H} \to \mathcal{H}^{*2}/\mathcal{H}^*$$

$$\downarrow \qquad \qquad \downarrow$$

$$\mathcal{H}/\mathcal{P} \times \mathcal{H}/\mathcal{P} \to \mathcal{H}/\mathcal{P}$$

where the vertical arrows are the isomorphisms induced by multiplication by 1 + i and 2 respectively and the horizontal arrows are the respective hermitian forms.

From now on, we shall identify $\mathcal{H}^*/\mathcal{H}$ with \mathbf{F}_4 , as a one dimensional vector space for the choice of the basis 1/1 + i.

- 2.3. Proposition.
- (a) Let $\mathcal{H}^n \subseteq L \subseteq \mathcal{H}^{*n}$ be a **Z**-module. Then $(L, Tr \circ h)$ is integral if and only if $\eta(L)$ is a totally isotropic subspace of the symmetric bilinear space $(\mathbf{F}_4^n, Tr \circ \eta(h))$, where $\eta(h)$ is the standard hermitian form on \mathbf{F}_4^n .
- (b) The **Z**-lattice $(L, Tr \circ h)$ is unimodular if and only if $\eta(L)$ is a maximal totally isotropic subspace of $(\mathbf{F}_{4}^{n}, Tr \circ \eta(h))$.

Proof. (a) This follows easily from 2.2.

(b) This follows from (a), since L is unimodular if and only if L is maximal integral.

§3. Perfect isometries of \mathcal{H} -lattices

In this section we show that certain special class of **Z**-lattices admit perfect isometries. We begin with the following definition.

- 3.1. Definition. A **Z**-lattice (L, b) is called an \mathcal{H} -lattice if L is an \mathcal{H} -module and $b = Tr \circ h$ for some hermitian form h.
- 3.2. Proposition. Every \mathcal{H} -lattice has a perfect isometry.

Proof. Let $(L, Tr \circ h)$ be an \mathcal{H} -lattice. Let $\sigma: L \to L$ denote left (or right) multiplication by ξ where ξ is one of the units $(1 \pm i \pm j \pm k)/2$. Then,

$$Tr \circ h(\sigma(x), \sigma(y)) = Tr \circ h(\xi x, \xi y) = Tr(\xi h(x, y)\overline{\xi})$$

$$= \xi h(x, y)\overline{\xi} + \xi h(x, y)\overline{\xi} = \xi (h(x, y) + \overline{h(x, y)})\overline{\xi} = \xi \overline{\xi} (h(x, y) + \overline{h(x, y)})$$

$$= h(x, y) + \overline{h(x, y)} = Tr \circ h(x, y).$$

Therefore σ is an isometry. Since the minimal polynomial of σ is $x^2 - x + 1$, $det(1 - \sigma) = 1$ and hence σ is perfect.

As a special case of this we have:

- 3.3. COROLLARY. The \mathcal{H} -lattice $(\mathcal{H}, Tr \circ h)$ has a perfect isometry.
- 3.4. PROPOSITION. Every perfect isometry of $(\mathcal{H}, Tr \circ h)$ induces a perfect \mathbf{F}_2 -isomorphism of $\mathcal{H}^*/\mathcal{H} = \mathbf{F}_4$, which corresponds to multiplication by ω , where $\mathbf{F}_2(\omega) = \mathbf{F}_4$.