

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	39 (1993)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
<b>Artikel:</b>	QUICK LOWER BOUNDS FOR REGULATORS OF NUMBER FIELDS
<b>Autor:</b>	Skoruppa, Nils-Peter
<b>Rubrik</b>	
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-60417">https://doi.org/10.5169/seals-60417</a>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

(to compare this inequality to the inequality as originally stated by Zimmert one has to apply the gamma duplication formula). He chose  $s = 2$  to obtain

$$\frac{R}{w} \geq 0.02 \cdot \exp(0.46r_1 + 0.1r_2).$$

Zimmert deduced his regulator bounds by an ingenious, but quite involved, investigation of certain analytic properties of the partial Dedekind zeta function associated to the class of principal ideals of  $K$ .

In this note we show that it is possible to deduce the above theorem by a simple estimate from a certain, almost obvious, monotonicity property of Hecke's theta function associated to the maximal order of  $K$  (see below). Moreover, we indicate below how this method of proof can be refined to yield exactly Zimmert's bounds. The technique of estimating which we apply is a sort of simple variation of a method which is developed in [F-S] to obtain lower bounds for  $L^p$ -norms of a certain class of functions. It was found during a careful analysis of Zimmert's method and reflects, though it looks much easier, still very much the spirit of Zimmert's original proof.

2. PROOF. Let  $|\cdot|_j$  for  $1 \leq j \leq r := r_1 + r_2$  denote the archimedean absolute values of  $K$ , let  $G$  denote the  $r$ -fold direct product of the multiplicative group of the positive reals  $\mathbf{R}_+$ , and let  $V$  denote the image in  $G$  of the units of  $K$  under the map

$$\eta \mapsto (\dots, |\eta|_j^{n_j}, \dots),$$

where  $n_j$  equals 1 or 2 accordingly as  $|\cdot|_j$  is real or complex. Denote by  $\delta$  the group homomorphism

$$\delta: G \rightarrow \mathbf{R}_+, \quad \delta((\dots, x_j, \dots)) = x_1 \cdots x_r.$$

Its kernel contains  $V$ , and by Dirichlet's unit theorem  $\ker \delta/V$  is compact. We can thus fix a Haar measure  $\mu$  on  $G/V$  by requiring

$$\int_{G/V} g \circ \delta d\mu = \frac{R}{w} \int_0^\infty g(t) \frac{dt}{t}$$

for any integrable function on  $\mathbf{R}_+$ . Let

$$Z(s) := \gamma(s) \sum_{\alpha \in \mathfrak{R}} |\mathrm{N}_{K/\mathbf{Q}} \alpha|^{-s},$$

where  $\mathfrak{R}$  is a set of representatives for the non-zero elements of  $\mathfrak{O}$ , the ring

of integers in  $K$ , modulo units. According to Hecke [H] (and according to the choice of  $\mu$ ) one has, for  $\operatorname{Re}(s) > 1$ , the integral representation

$$Z(s) = \int_{G/V} \Theta \delta^s d\mu .$$

Here  $\Theta$  is a smooth, non-negative and  $V$ -invariant function on  $G$ , which is given by

$$\Theta(x) = \sum_{\substack{\alpha \in \mathfrak{D} \\ \alpha \neq 0}} \exp \left( - \sum_{j=1}^r |\alpha|_j^2 x_j^{2/n_j} \right) .$$

The main observation for the proof of the theorem is the

**LEMMA.** *The function  $(1 + \Theta)\delta$  is increasing in each argument.*

*Proof.* This follows from Hecke's theta formula [H, p. 165-166]

$$(1 + \Theta(x)) \delta(x) = \frac{\pi^{\frac{n}{2}} 2^{r_2}}{\sqrt{|d|}} \sum_{\alpha \in \mathfrak{D}^{-1}} \exp \left( - \pi^2 \sum_{j=1}^r n_j^2 |\alpha|_j^2 x_j^{-2/n_j} \right) ,$$

i.e. by applying Poisson summation to the series defining  $\Theta(x)$  (here  $\mathfrak{D}$  and  $d$  denote the different and discriminant of  $K$ ).  $\square$

We can now give the

*Proof of the theorem.* For  $a \in \mathbf{R}_+$  set

$$I(a) := \int_{G/V} (1 + \Theta(x)) \delta(x) w((a\delta(x))^{s-1}) d\mu(x) ,$$

where we use

$$w(t) = t \max(0, \log(1/t)) ,$$

and where  $s > 1$  as in the theorem.

For any  $\varepsilon > 0$ , one has  $w(t) = O(t^\varepsilon)$  and  $|w(t+h) - w(t)| / |h| \leq |w'(t)| = O(t^{-\varepsilon})$  as  $t \rightarrow 0$ . Thus, using the convergence of the integral representation of  $Z(s)$  for  $s > 1$ , we deduce that the integral defining  $I(a)$  is finite, and, on applying Lebesgue's theorem, that  $I(a)$  is differentiable and its derivative is obtained by differentiating under the integral sign. Here we agree to use  $w'(1)$  for the derivative on the right, i.e.  $w'(1) = 0$ .

On replacing  $x$  by  $x/a^{1/r}$  in the integral defining  $I(a)$  we deduce from the lemma that  $I(a)$  is decreasing. Hence  $I'(a) \leq 0$ , from which we obtain, writing  $\sigma = s - 1$ ,

$$\begin{aligned} & -\frac{d}{da} \int_{G/V} \delta w((a\delta)^\sigma) d\mu \geq \frac{d}{da} \int_{G/V} \Theta \delta w((a\delta)^\sigma) d\mu \\ &= \sigma a^{\sigma-1} \int_{G/V} \Theta \delta^s w'((a\delta)^\sigma) d\mu \geq -\sigma a^{\sigma-1} \int_{G/V} \Theta \delta^s (1 + \log(a\delta)^\sigma) d\mu \\ &= -\sigma^2 a^{\sigma-1} Z(s) \left( \frac{1}{\sigma} + \log a + \frac{Z'}{Z}(s) \right). \end{aligned}$$

By the choice of  $\mu$  the left-hand side equals  $R\sigma/(ws^2a^2)$ . Multiplying the above inequality by  $s^2a^2/\sigma$  and then maximizing the right hand side, i.e. choosing

$$\frac{1}{\sigma} + \log a + \frac{Z'}{Z}(s) = -\frac{1}{s},$$

we find

$$(1) \quad \frac{R}{w} \geq \frac{s(s-1)}{e} \exp \left( -\frac{s}{s-1} \right) Z(s) \exp \left( -s \frac{Z'}{Z}(s) \right).$$

Finally,  $Z(s) \geq \gamma(s)$ , since the Dirichlet series  $D(s)$  in the definition of  $Z(s)$  satisfies  $D(s) > 1$ , and  $\frac{Z'}{Z}(s) \leq \frac{\gamma'}{\gamma}(s)$ , since  $D'(s) < 0$ . Thus, (1) implies the claimed inequality.  $\square$

**3. CONCLUDING REMARKS.** To obtain a lower bound as sharp as Zimmert's one can proceed as above, but with a variant  $\Theta_1$  of the function  $\Theta$ . Namely, fix a real number  $s > 1$ , and define  $\Theta_1(x)$  by the same series as  $\Theta(x)$  but with the term

$$\exp \left( - \sum |\alpha|_j x_j^{2/n_j} \right)$$

replaced by

$$\prod_{j=1}^r f_j(|\alpha|_j^{n_j} x_j),$$