

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ERGODIC ADDING MACHINE ON THE CANTOR SET
Autor: Collas, Peter / Klein, David
DOI: <https://doi.org/10.5169/seals-61113>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

AN ERGODIC ADDING MACHINE ON THE CANTOR SET

by Peter COLLAS and David KLEIN

ABSTRACT. We calculate all ergodic measures for a specific function F on the unit interval. The supports of these measures consist of periodic orbits of period 2^n and the classical ternary Cantor set. On the Cantor set, F is topologically conjugate to an “adding machine” in base 2. We show that F is representative of the class of functions with zero topological entropy on the unit interval, already analyzed in the literature, and its behavior is therefore typical of that class.

I. INTRODUCTION

The dynamical behavior of the quadratic function $f_c(x) = x^2 - c$ has been extensively studied as the parameter c is varied. For example, $c_0 = 1.401155189\dots$ is the smallest value of c for which $f_c(x)$ has infinitely many distinct periodic orbits [1-3]. As c approaches this number through smaller values, the dynamical system, $x \rightarrow f_c(x)$, progresses through the famous period doubling route to chaos. When $c = c_0$, the dynamical behavior of $f(x) \equiv f_c(x)$ includes the following properties:

1. There is a Cantor set K which is an attractor and $f: K \rightarrow K$
2. All periodic points of f have period 2^n for some n .
3. There are periodic points which are arbitrarily close to K .
4. With the restriction of $f(x)$ to an appropriate interval I such that $f(I) \subset I$, there are just two possibilities for the orbit of a point $x_0 \in I$: either $f^k(x_0)$ is in a periodic orbit for some k , or $f^k(x_0)$ converges to K as k increases.
5. The restriction of f to K is topologically equivalent to a function on 2-adic integers which adds 1 to its argument (this “adding machine” will be described in detail below).

The Cantor set K is sometimes called a Feigenbaum attractor. When $\mu = 3.57\dots$, the well-known logistic function $g_\mu(x) = \mu x(1 - x)$ exhibits the same dynamical properties [4]. In fact, a large class of dynamical systems exhibiting the properties 1 through 5 has been studied and the ergodic properties analyzed [2, 3, 5].

A particularly simple example of a dynamical system on the interval $[0, 1]$ satisfying 1 through 5 was given and studied by Delahaye [6] and, in a slightly different form, its topological properties (including 1-5) were given in the statements of a series of exercises by Devaney [7]. The function may be described through the concept of the "double" of a function (cf. [7]) as follows: Let $f_0(x) \equiv \frac{1}{3}$ and define $f_n(x)$ recursively by

$$(1) \quad f_{n+1}(x) = \begin{cases} \frac{1}{3} f_n(3x) + \frac{2}{3} & \text{if } x \in [0, \frac{1}{3}] \\ -\frac{7}{3}x + \frac{14}{9} & \text{if } x \in [\frac{1}{3}, \frac{2}{3}] \\ x - \frac{2}{3} & \text{if } x \in [\frac{2}{3}, 1] \end{cases},$$

and

$$(2) \quad F(x) = \lim_{n \rightarrow \infty} f_n(x).$$

It follows that F is continuous on $[0, 1]$ and that it is its own double, i.e.,

$$(3) \quad F(x) = \begin{cases} \frac{1}{3} F(3x) + \frac{2}{3} & \text{if } x \in [0, \frac{1}{3}] \\ -\frac{7}{3}x + \frac{14}{9} & \text{if } x \in [\frac{1}{3}, \frac{2}{3}] \\ x - \frac{2}{3} & \text{if } x \in [\frac{2}{3}, 1] \end{cases}.$$

The function F is shown in Figure 1. The notion of the double of a function and its use in studying dynamical systems goes back to Sharkovskii [8]. A general definition of the double of a function, however, will not be needed here.

We will show in the sequel that the function F , like $x^2 - c_0$, is not chaotic. F closely models the behavior of the quadratic function $x^2 - c$ at the critical value c_0 (and many other functions at corresponding critical values of an associated parameter as well) beyond which chaos is present. In addition, the sequence f_n , in its approach to F , exhibits the classical period doubling bifurcations, characteristic of the onset of chaos [7]. The function F is a simple model for understanding the point of transition from nonchaotic behavior to chaotic behavior. In this note we summarize the topological




FIGURE 1
Equation (3), the adding machine

properties of the dynamical system $x \rightarrow F(x)$ in the form of Theorems 1.1 and 1.2 below and then show how ergodic theory may be used to further analyze the dynamical system. We then indicate how this system may be understood from a more general context developed by Misiurewicz [2, 3] involving topological entropy.

We refer to the following commonly used terms (cf. ref. 7). The point y_0 is a fixed point of F if $F(y_0) = y_0$. The point y is a periodic point of period n if $F^n(y) = y$. The least positive n for which $F^n(y) = y$ is called the *prime period* of y . Hereafter when we refer to a periodic point of period n it shall be understood that n is the prime period. The set of all iterates of a

periodic point is a periodic orbit. We shall denote the set of periodic points of period n by $\text{Per}_n(F)$. Finally a point x is *eventually periodic* of period n if x is not periodic but there exists a $j > 0$ such that $F^{n+i}(x) = F^i(x)$ for all $i \geq j$. In other words although x is not itself periodic, an iterate of x is.




FIGURE 2

Three stages on the way to the Cantor set

Theorem 1.1 below makes reference to the classical ternary Cantor set in $[0, 1]$. We will use the following labeling: The “middle third” intervals that are removed on the way to obtaining the Cantor set are labeled A_n^k , for

example, $A_0^1 \equiv A_0 = (\frac{1}{3}, \frac{2}{3})$, $A_2^4 = (\frac{25}{27}, \frac{26}{27})$, (Figure 2). Set $A_n = \bigcup_{k=1}^{2^n} A_n^k$.

Thus A_n consists of 2^n intervals which we number from left to right. We

let $(A_{n-1})^c \equiv I_n$, and $I_n = \bigcup_{k=1}^{2^n} I_n^k$, so that I_n also consists of 2^n intervals

which we again number from left to right. So, for example, $I_1^1 = [0, \frac{1}{3}]$, $I_3^8 = [\frac{26}{27}, 1]$, (Figure 2), but $I_0 = [0, 1]$. Denote the ternary Cantor set

by $I_\infty = \bigcap_{n=0}^{\infty} I_n$. It is well-known, and easily deduced, that a real number

in $[0, 1]$ is in the Cantor set I_∞ if and only if it has a ternary expansion (“base 3 decimal expansion”) of the form $0.\alpha_1\alpha_2\alpha_3\dots$, where $\alpha_k = 0$ or 2 for each k .

Delahaye’s results and Devaney’s exercises are slightly extended by Theorem 1.1 below.

THEOREM 1.1. *The function $F: [0, 1] \rightarrow [0, 1]$ given by (2) satisfies the following properties:*

- (a) *For each n , F is a cyclic permutation on the collection of intervals $\{I_n^k: k = 1, \dots, 2^n\}$, i.e., for given k , $F^{2^n}(I_n^k) = I_n^k$, and for any $p \neq k$, $p = 1, \dots, 2^n$, $F^j(I_n^k) = I_n^p$ for precisely one j between 1 and $2^n - 1$.*
- (b) *For each $n = 0, 1, 2, \dots$, F has exactly one periodic orbit with period 2^n and no other periodic orbits.*
- (c) *Every periodic orbit is repelling. $\text{Per}_{2^n}(F) \subset A_n$ and A_n^k contains exactly one point from $\text{Per}_{2^n}(F)$ for each $k = 1, \dots, 2^n$ and each nonnegative integer n .*
- (d) *Each point is eventually periodic or converges to I_∞ under repeated iterations of F .*

We briefly sketch part of the proof of Theorem 1.1. For $n \geq 2$, it can be shown, using induction on n , that

$$F(I_n^k) = I_n^{G(k)},$$

where

$$G(k) = \begin{cases} k - 2^{n-1} & \text{for } 2^{n-1} + 1 \leq k \leq 2^n \\ 2^n & \text{for } k = 1 \\ k + (2^{N+1} - 3)2^{n-N-1} & \text{for } 2 \leq k \leq 2^{n-1} \end{cases}$$

and where $N = \left[n - \frac{\log k}{\log 2} \right]$ (and $[]$ denotes "integer part"). Part (a) of Theorem 1.1 may now be deduced using this formula and (3).

To check part (b), observe that if $x \in \left[\frac{1}{3}, \frac{2}{3} \right]$, then iterates of x by F will eventually move out of $\left[\frac{1}{3}, \frac{2}{3} \right]$ and never return (see Figure 1). If $x \in \left[0, \frac{1}{3} \right]$, then $F(x) \in \left[\frac{2}{3}, 1 \right]$, and if $x \in \left[\frac{2}{3}, 1 \right]$, then $F(x) \in \left[0, \frac{1}{3} \right]$. Therefore F has no odd periods. An induction argument shows that if $x \in \left[0, \frac{1}{3} \right]$, then $F^{2^n}(x) = \frac{1}{3}F^n(3x)$. To show F does not have any even period orbits other than period 2^n orbits, suppose that there is a period 2^nk orbit, where $k > 1$ is an odd number and $n \geq 1$. If $x \in \left[0, \frac{1}{3} \right]$ and $F^{2^nk}(x) = x$, then $F^{2^{n-1}k}(3x) = 3x$ and $3x \in \text{Per}_{2^{n-1}k}(F)$. Therefore there is an $x \in \left[0, \frac{1}{3} \right]$ such that $F^{2^{n-1}k}(x) = x$. Continuing in this way we will reach a point such that $F^k(y) = y$, which is impossible since there are no odd period orbits. The existence of a unique orbit of period 2^n follows from (3) and induction on n .

The proofs of parts (c) and (d) use similar ideas and are outlined in the exercises in [7]. \square

We turn now to a description of the “adding machine” on the ternary Cantor set and its relationship to F .

A 2-adic integer is an infinite sequence $x = (x_0, x_1, x_2, \dots)$ where $x_i = 0$ or 1. The collection S of all 2-adic integers is a metric space with the metric $d(x, y) = 2^{-n}$ where $y = (y_0, y_1, y_2, \dots)$ and n is the smallest integer for which $x_n \neq y_n$. S is the completion of the nonnegative integers with this metric under the identification of the (base 2) integer

$$m = x_0 + x_1 2^1 + x_2 2^2 + \dots + x_n 2^n$$

with the sequence

$$(4) \quad (x_0, x_1, x_2, \dots, x_n, 0, 0, 0, \dots).$$

Define a base 2 addition on S by

$$x + y = z = (z_0, z_1, z_2, \dots),$$

where $z_0 = x_0 + y_0$ if $x_0 + y_0 \leq 1$, $z_0 = 0$ if $x_0 + y_0 = 2$ in which case 1 is added to $x_1 + y_1$, which otherwise follows the same rules. The numbers z_2, z_3, \dots are successively determined in the same manner. Thus, if $x = (x_0, x_1, x_2, \dots, x_n, 0, 0, 0, \dots)$ and $y = (y_0, y_1, y_2, \dots, y_k, 0, 0, 0, \dots)$, then $x + y$ corresponds to the usual base 2 arithmetic addition of integers under the identification (4). S is a commutative, compact topological group with this addition. Let us denote the element $(1, 0, 0, \dots)$ of S by $\mathbf{1}$. Define a map $h: I_\infty \rightarrow S$ as follows: If $x \in I_\infty$ has base 3 expansion $0.\alpha_0\alpha_1\alpha_2\dots$, where each $\alpha_i = 0$ or 2, then $h(x) = (x_0, x_1, x_2, \dots)$, where $x_i = 1 - \frac{\alpha_i}{2}$. For example,

$$h(0.02022\dots) = (1, 0, 1, 0, 0, \dots).$$

Theorem 1.2 below was also stated in the same set of exercises in Devaney [7]. We supply the proof for the convenience of the reader.

THEOREM 1.2.

- (a) *The function h is a homeomorphism from the ternary Cantor set I_∞ to the 2-adic integers S .*
- (b) *$F(I_\infty) = I_\infty$, and F restricted to I_∞ is topologically conjugate by h to the addition of $\mathbf{1}$ on 2-adic integers, i.e., $h(F(x)) = h(x) + \mathbf{1}$.*
- (c) *The F -orbit of each point in I_∞ is dense in I_∞ .*

Proof. (a) h is clearly one-to-one and onto. To see that h^{-1} is continuous, let $\varepsilon > 0$ be given and choose n so that $3^{-n-1} < \varepsilon$ and let $\delta = 2^{-n}$. If $x, y \in S$ and $d(x, y) \leq \delta$, then

$$(5) \quad |h^{-1}(x) - h^{-1}(y)| = |0.000\dots 0\alpha_n\alpha_{n+1}\dots|$$

where the first $n - 1$ digits on the right side of (5) are zeros and the number on the right is expressed in base 3 so that $\alpha_k = 0$ or 2. Consequently $|h^{-1}(x) - h^{-1}(y)| \leq 3^{-n-1} < \varepsilon$. Since h^{-1} is a continuous bijection and I_∞ is compact, it follows from a well-known theorem in topology that h is continuous and therefore h is a homeomorphism.

(b) Suppose $x \in I_\infty \cap [\frac{2}{3}, 1]$ and let the base 3 expansion of x be given by $x = 0.2\alpha_1\alpha_2\alpha_3\dots$, thus $h(x) = (0, x_1, x_2, \dots)$, where $x_i = 1 - \frac{\alpha_i}{2}$. Then $F(x) = x - \frac{2}{3}$ has base 3 expansion given by $F(x) = 0.0\alpha_1\alpha_2\alpha_3\dots$. Therefore $h(F(x)) = (1, x_1, x_2, \dots)$ which is the same as $h(x) + 1$. If $x = 0$, then $h(F(0)) = (0, 0, 0, \dots) = h(0) + 1$. If $x \in I_\infty \cap (0, \frac{1}{3}]$, then $x \in [\frac{2}{3^i}, \frac{1}{3^{i-1}})$ for some $i \geq 2$. A brief calculation shows that

$$(6) \quad F(x) = x + 0.\overbrace{2\dots 2}^{i-2}1\underbrace{10}_i$$

where the number on the right is expressed in base 3 and the second "1" occurs i places after the decimal point. It now follows from (6) and base 3 addition that $h(F(x)) = h(x) + 1$ and therefore $F(I_\infty) = I_\infty$.

(c) Since S is the completion of the nonnegative integers under the identification (4), the set

$$\{(x_0, x_1, x_2, \dots, x_n, 0, 0, 0, \dots) : n = 0, 1, 2, \dots, x_i = 0 \text{ or } 1\}$$

which equals $\{n\mathbf{1} : n = 0, 1, 2, \dots\}$ is dense in S . It is easy to establish that the map which takes x to $x + z$ is a homeomorphism from S to S for any fixed $z \in S$. Thus

$$\{n\mathbf{1} : n = 0, 1, 2, \dots\} + z$$

is dense in S for any z . But $h^{-1}(\{n\mathbf{1} : n = 0, 1, 2, \dots\} + z)$ is precisely the F -orbit of $h^{-1}(z)$. Therefore the F -orbit of any $y = h^{-1}(z) \in I_\infty$ is dense in I_∞ . \square

II. ERGODIC MEASURES FOR F

A measure μ on a set X is called a probability measure if $\mu(X) = 1$; the pair (X, μ) is then called a probability space. Given a measurable transformation $T: X \rightarrow X$ on a probability space (X, μ) , μ is T -invariant if $\mu = \mu \circ T^{-1}$, i.e., for any measurable set $B \subset X$, $\mu(B) = \mu(T^{-1}(B))$. The probability measure μ is ergodic if $T^{-1}(A) = A$ implies that $\mu(A)$ is 0 or 1.

One way to study the attracting, often fractal, sets of a dynamical system (for example the ternary Cantor set for F) is to study the invariant probability measures which are supported on the attractors. (The support of a measure on $[0, 1]$ is the intersection of all closed subsets of $[0, 1]$ whose complements have zero measure.) This is an especially fruitful approach when the attracting set lies in a high dimensional space so that a purely geometric description is unfeasible (cf. [9]). Particularly important, though generally difficult to find, are the invariant ergodic probability measures for the dynamical system. For Devaney's transformation $F: [0, 1] \rightarrow [0, 1]$, we will find all F -invariant ergodic probability measures.

To construct a probability measure on I_∞ , we borrow some ideas from the theory of iterated function systems as developed by Barnsley [10]. Let $P[0, 1]$ denote the set of all Borel probability measures on $[0, 1]$, i.e., probability measures on the σ -algebra generated by all open sets on $[0, 1]$. For any $\mu, \nu \in P[0, 1]$, define

$$d(\mu, \nu) \equiv \sup \left\{ \left| \int f d\mu - \int f d\nu \right| : f : [0, 1] \rightarrow [0, 1] \right. \\ \left. \text{and } |f(x) - f(y)| \leq |x - y|, \forall x, y \in [0, 1] \right\}.$$

Then d is a complete metric on $P[0, 1]$. Let $w_1, w_2 : [0, 1] \rightarrow [0, 1]$ by $w_1(x) = \frac{1}{3}x$ and $w_2(x) = \frac{1}{3}x + \frac{2}{3}$. Let $M: P[0, 1] \rightarrow P[0, 1]$ by $M(\mu) = \frac{1}{2}\mu \circ w_1^{-1} + \frac{1}{2}\mu \circ w_2^{-1}$. M is called the Markov operator for an iterated function system defined by w_1, w_2 . It follows as a special case of a more general theorem from ref. 10 that

$$(7) \quad d(M(\mu), M(\nu)) \leq \frac{1}{3} d(\mu, \nu),$$

so that by the well-known contraction mapping theorem, M has a unique fixed point ν_∞ , i.e., $M(\nu_\infty) = \nu_\infty$.

Notice that the intervals I_n^k defined above are in one-to-one correspondence with iterations of the form $w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n}([0, 1])$, where $i_k = 1$ or 2 . For example, $w_1 \circ w_2([0, 1]) = \left[\frac{2}{9}, \frac{1}{3}\right] \equiv I_2^2$.

LEMMA 2.1. For all $n \geq 0$ and $k \leq 2^n$, $\nu_\infty(I_n^k) = 2^{-n}$.

Proof. The proof follows by induction. For $n = 0$, $I_0^1 = [0, 1]$ and $\nu_\infty([0, 1]) = 1 = 2^0$ because ν_∞ is a probability measure. Assume $M(\nu_\infty) = \nu_\infty$ and the induction hypothesis, $\nu_\infty(I_n^k) = \nu_\infty(w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n}[0, 1]) = 2^{-n}$

for any sequence i_1, \dots, i_n and any $k \leq 2^n$. For any $j \leq 2^{n+1}$, there is a sequence i_1, \dots, i_{n+1} such that

$$\begin{aligned} v_\infty(I_{n+1}^j) &= v_\infty(w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n} \circ w_{i_{n+1}}[0, 1]) \\ &= \frac{1}{2} v_\infty \circ w_1^{-1}(w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n} \circ w_{i_{n+1}}[0, 1]) \\ (8) \quad &+ \frac{1}{2} v_\infty \circ w_2^{-1}(w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n} \circ w_{i_{n+1}}[0, 1]). \end{aligned}$$

Notice that $v_\infty \circ w_j^{-1}(w_{i_1} \circ w_{i_2} \circ \cdots \circ w_{i_n} \circ w_{i_{n+1}}[0, 1]) = 2^{-n}$ or 0 depending on whether or not $j = i_1$. Combining this observation with (8) shows $v_\infty(I_{n+1}^j) = \frac{1}{2} 2^{-n} = 2^{-n-1}$. \square

PROPOSITION 2.1. *The measure v_∞ is supported on the Cantor set I_∞ .*

Proof. By Lemma 2.1 $v_\infty(I_n) = \sum_{k=1}^{2^n} v_\infty(I_n^k) = 2^n 2^{-n} = 1$ for all n .

Since $I_0 \supset I_1 \supset I_2 \supset \cdots$ is a decreasing sequence, $v_\infty(I_\infty) = v_\infty\left(\bigcap_{n=1}^{\infty} I_n\right) = \lim_{n \rightarrow \infty} v_\infty(I_n) = 1$. If C is a proper closed subset of I_∞ , there is an open

interval U in the complement of C whose intersection with I_∞ is nonempty. Let $x \in U \cap I_\infty$. For some positive integers n and k , $x \in I_n^k \subset U$. Then since I_n^k is in the complement of C , $v_\infty(C) \leq 1 - 2^{-n} < 1$. \square

The next step is to show that v_∞ is an invariant measure for the dynamical system $F: [0, 1] \rightarrow [0, 1]$. This means that $v_\infty = v_\infty \circ F^{-1}$, i.e., for any Borel set $B \subset [0, 1]$, $v_\infty(B) = v_\infty(F^{-1}(B))$.

LEMMA 2.2. *If μ is any probability measure on $[0, 1]$ and $\mu(I_n^k) = 2^{-n}$ for all n and k , then $\mu = v_\infty$.*

Proof. The condition $\mu(I_n^k) = 2^{-n}$ implies that $\mu\{x\} = 0$ for every $x \in [0, 1]$. This is clearly true if $x \notin I_\infty$ since $\mu(I_\infty) = 1$, as follows from the proof of Proposition 2.1. If $x \in I_\infty$, then for every n there exists a k such that $x \in I_n^k$. Thus $\mu\{x\} \leq 2^{-n}$ for every n and hence $\mu\{x\} = 0$.

Any interval of the form $[a, b] \subset [0, 1]$ where a and b have finite ternary expansions (finite “decimal” expansions in base 3) is a disjoint union of sets of the following form:

- 1) I_n^k for some n and k
- 2) intervals in the complement of I_n for some n
- 3) $\{a\}, \{b\}$.

Since μ and ν_∞ agree on each of the sets in 1, 2, and 3, $\mu[a, b] = \nu_\infty[a, b]$. It is a standard result in measure theory that two probability measures which are equal on a collection of measurable sets, closed under finite intersections, are equal on the σ -algebra generated by that collection. In this case the σ -algebra generated by sets of the form $[a, b] \subset [0, 1]$ where a and b have finite ternary expansions is just the Borel σ -algebra on $[0, 1]$. \square

It is well-known [11] that the “adding machine” is uniquely ergodic. This fact may also be deduced from Proposition 2.2 below.

PROPOSITION 2.2. ν_∞ is invariant under F . If μ is a probability measure invariant under F and $\mu(I_\infty) = 1$, then $\mu = \nu_\infty$.

Proof. To show that ν_∞ is invariant under F , it suffices by Lemma 2.2 to show that

$$\nu_\infty(F^{-1}(I_n^k)) = 2^{-n}$$

for all k and n . By Theorem 1.1 (a), given k there exists a unique integer j such that

$$(9) \quad I_n^j \subset F^{-1}(I_n^k) \subset I_n^j \cup [0, 1] \setminus I_\infty$$

(because F is a permutation on the intervals in I_n). Thus,

$$2^{-n} = \nu_\infty(I_n^j) \leq \nu_\infty(F^{-1}(I_n^k)) \leq \nu_\infty(I_n^j \cup [0, 1] \setminus I_\infty) = 2^{-n}.$$

Hence ν_∞ is invariant under F . Suppose μ is a probability measure invariant under F and $\mu(I_\infty) = 1$. Then by (9) and F -invariance,

$$(10) \quad \mu(I_n^j) \leq \mu(F^{-1}(I_n^k)) = \mu(I_n^k) \leq \mu(I_n^j \cup [0, 1] \setminus I_\infty) = \mu(I_n^j).$$

Since F is a cyclic permutation on the intervals in I_n , for any n and any positive integers $j, k \leq 2^n$, $\mu(I_n^j) = \mu(I_n^k)$. As there are 2^n intervals of the form I_n^k in I_n , it follows that $\mu(I_n^k) = 2^{-n}$ for all n and k . Therefore, $\mu = \nu_\infty$, by Lemma 2.2. \square

Before investigating the ergodicity of ν_∞ , we introduce some other invariant measures for F . For a fixed $x \in [0, 1]$, let δ_x be the probability measure on $[0, 1]$ which assigns 1 to any Borel measurable set containing x and assigns 0 to all other measurable sets. The probability measure δ_x is sometimes called the “point mass at x ” or the “Dirac delta function at x .”

For each nonnegative integer n , let y_n be the smallest number in $[0, 1]$ which lies in the unique orbit with prime period 2^n for the function F and define

$$(11) \quad \nu_n = \frac{1}{2^n} \sum_{k=0}^{2^n-1} \delta_{F^k(y_n)}.$$

The measure ν_n assigns mass 2^{-n} to each point in the unique orbit with period 2^n of F , i.e., $\nu_n(B) = k2^{-n}$ if B contains exactly k points from $P_n \equiv \text{Per}_{2^n}(F)$. It is not difficult to check that for each $n = 0, 1, 2, \dots$, ν_n is invariant with respect to F , and ν_n is the only F -invariant probability measure on the set of points P_n .

Let $C[0, 1]$ be the Banach space consisting of all continuous functions with the maximum norm $\| \cdot \|$ given by $\|f\| \equiv \max\{|f(x)| : x \in [0, 1]\}$. Let $M_F[0, 1] \equiv \{\mu \in P[0, 1] : \mu = \mu \circ F^{-1}\}$ be the set of invariant probability measures on $[0, 1]$. $M_F[0, 1]$ may be identified in a natural way with a metrizable, compact, convex subset of the dual space of $C[0, 1]$. The compact, metrizable topology on $M_F[0, 1]$ is the weakest topology which makes the map $\mu \rightarrow \int f(x) \mu(dx)$ continuous for each $f \in C[0, 1]$; it is called the vague or weak-* topology. An extreme point of the convex set $M_F[0, 1]$ is a measure μ which is not a convex combination of any other two points in $M_F[0, 1]$, i.e., μ is extreme if whenever $\mu = \alpha\mu_1 + (1 - \alpha)\mu_2$, $0 < \alpha < 1$, and $\mu_1, \mu_2 \in M_F[0, 1]$, then $\mu = \mu_1 = \mu_2$. It is a consequence of the Krein-Milman Theorem that the set of extreme points of $M_F[0, 1]$ is non-empty. The following theorem is a specialization of a well-known result in ergodic theory (see for example [12]).

THEOREM 2.1. *The F -invariant measure μ is an extreme point of $M_F[0, 1]$ if and only if μ is ergodic with respect to F on $[0, 1]$.*

As a consequence of Theorem 2.1, we have the following proposition.

PROPOSITION 2.3. *For each $n = 0, 1, 2, \dots, \infty$, ν_n is an extreme point of $M_F[0, 1]$ and is therefore ergodic with respect to F .*

Proof. Consider the case $n = \infty$. Suppose there exist F -invariant probability measures μ_1 and μ_2 and $\alpha \in (0, 1)$ such that $\nu_\infty = \alpha\mu_1 + (1 - \alpha)\mu_2$. Then since $\nu_\infty(I_\infty) = 1$, $\mu_1(I_\infty) = \mu_2(I_\infty) = 1$. Then by Proposition 2.2, $\mu_1 = \mu_2 = \nu_\infty$. Thus ν_∞ is an extreme point of $M_F[0, 1]$ and is ergodic by Theorem 2.1. The cases, $n = 0, 1, 2, \dots$ are handled in the same manner. \square

PROPOSITION 2.4. *The measures $\nu_0, \nu_1, \nu_2, \dots, \nu_\infty$ are the only probability measures on $[0, 1]$ ergodic with respect to F .*

Proof. Let μ be an F -invariant probability measure with support A_μ . Let $\mathcal{P} \equiv \{x \in [0, 1] : F^k(x) \in P_n \text{ for some } k, n\}$ be the set of periodic or eventually periodic points. By Theorem 1.1 $A_\mu \setminus \mathcal{P} \subset \bigcup_{k=0}^{\infty} F^{-k}(I_n)$ for each n . Thus

$$\mu(A_\mu \setminus \mathcal{P}) \leq \mu\left(\bigcup_{k=0}^{\infty} F^{-k}(I_n)\right).$$

Since $I_n \subset F^{-1}(I_n) \subset F^{-2}(I_n) \subset \dots$ is an increasing sequence of sets,

$$\mu\left(\bigcup_{k=0}^{\infty} F^{-k}(I_n)\right) = \lim_{k \rightarrow \infty} \mu(F^{-k}(I_n)) = \mu(I_n) \geq \mu(A_\mu \setminus \mathcal{P})$$

for all n . Since $I_\infty = \bigcap_{n=1}^{\infty} I_n$, is a decreasing sequence of sets,

$$\mu(I_\infty) \geq \mu(A_\mu \setminus \mathcal{P}).$$

Similarly,

$$\mu\left(\bigcup_{k=0}^{\infty} F^{-k}(P_n)\right) = \lim_{k \rightarrow \infty} \mu(F^{-k}(P_n)) = \mu(P_n) \geq \mu\left(A_\mu \cap \bigcup_{k=0}^{\infty} F^{-k}(P_n)\right)$$

Thus,

$$\mu\left(\bigcup_{n=0}^{\infty} P_n\right) = \sum_{n=0}^{\infty} \mu(P_n) \geq \mu(A_\mu \cap \mathcal{P}).$$

Since $A_\mu = (A_\mu \setminus \mathcal{P}) \cup (A_\mu \cap \mathcal{P})$, it follows that

$$\begin{aligned} 1 &\geq \mu\left(I_\infty \cup \bigcup_{n=0}^{\infty} P_n\right) = \mu(I_\infty) + \mu\left(\bigcup_{n=0}^{\infty} P_n\right) \\ &\geq \mu(A_\mu \setminus \mathcal{P}) + \mu(A_\mu \cap \mathcal{P}) = \mu(A_\mu) = 1. \end{aligned}$$

Hence $\mu\left(I_\infty \cup \bigcup_{n=0}^{\infty} P_n\right) = 1$.

From Theorem 1.1, $\bigcup_{n=0}^{\infty} A_n \setminus \bigcup_{n=0}^{\infty} P_n$ is a union of open intervals. Therefore

$$\left(I_\infty \cup \left(\bigcup_{n=0}^{\infty} P_n\right)\right)^c = \bigcup_{n=0}^{\infty} A_n \setminus \bigcup_{n=0}^{\infty} P_n$$

is open. Hence $I_\infty \cup \left(\bigcup_{n=0}^{\infty} P_n \right)$ is a closed subset of $[0, 1]$. Since

$\mu \left(I_\infty \cup \bigcup_{n=0}^{\infty} P_n \right) = 1$ the definition of the support of a measure implies that

$$A_\mu \subset I_\infty \cup \bigcup_{n=0}^{\infty} P_n.$$

Since F is a continuous function and A_μ is closed, $F^{-1}(A_\mu)$ is also closed. By F -invariance, $\mu(F^{-1}(A_\mu)) = 1$ and therefore $A_\mu \subset F^{-1}(A_\mu)$. Hence $F(A_\mu) \subset A_\mu$. Suppose $A_\mu \cap I_\infty \neq \emptyset$ and let $x \in A_\mu \cap I_\infty$. Then $F^k(x) \in A_\mu \cap I_\infty$ for all k . Since by Theorem 1.2 the orbit of x is dense in I_∞ and $A_\mu \cap I_\infty$ is closed, it follows that $A_\mu \cap I_\infty = I_\infty$, i.e., $I_\infty \subset A_\mu$. A similar argument shows that if $A_\mu \cap P_n \neq \emptyset$, then $P_n \subset A_\mu$.

Thus A_μ is a union of one or more of $I_\infty, P_0, P_1, P_2, \dots$. Furthermore, if $\mu(I_\infty) = 1$ then by Proposition 2.2, $\mu = \nu_\infty$. Similarly if $\mu(P_n) = 1$ for some n , then $\mu = \nu_n$. If $0 < \mu(I_\infty) < 1$, then

$$F^{-1} \left(\bigcup_{k=0}^{\infty} F^{-k}(I_\infty) \right) = \bigcup_{k=0}^{\infty} F^{-k}(I_\infty)$$

is an invariant set for F and

$$\mu \left(\bigcup_{k=0}^{\infty} F^{-k}(I_\infty) \right) = \lim_{k \rightarrow \infty} \mu(F^{-k}(I_\infty)) = \mu(I_\infty).$$

Therefore μ is not ergodic. Similarly if $0 < \mu(P_n) < 1$, then μ is not ergodic. \square

Using the above results and some general theorems from ergodic theory, it is possible to give alternative descriptions of the measure ν_∞ . The following theorem may be found in [12].

THEOREM 2.2. *Let X be a compact metric space, $T: X \rightarrow X$ a continuous map, and assume ν is the unique probability measure on X which is invariant with respect to T . Then for any continuous real valued function f on X .*

$$(12) \quad \frac{1}{N} \sum_{k=0}^{N-1} f(T^k(x)) \rightarrow \int f d\nu$$

uniformly for all $x \in X$.

Note that by the Birkhoff Ergodic Theorem the convergence in (12) holds for any integrable function f pointwise for almost all $x \in X$. Before applying Theorem 2.2 to F , we cite the following special case of a theorem of Choquet [13].

THEOREM 2.3. *For any $\mu \in M_F[0, 1]$ there exists a Borel probability measure m_μ on the set of extreme points of $M_F[0, 1]$ such that $\mu = \int v m_\mu(dv)$.*

PROPOSITION 2.5. *For any continuous function f on $[0, 1]$*

(a) $\int f(x) v_\infty(dx) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=0}^{N-1} f(F^k(x_0))$ *for all x_0 not eventually periodic, and uniformly for all x_0 in I_∞ .*

(b) $\int f(x) v_\infty(dx) = \lim_{n \rightarrow \infty} \int f(x) v_n(dx) = \lim_{n \rightarrow \infty} \frac{1}{2^n} \sum_{k=0}^{2^n-1} f(F^k(y_n))$
where y_n is the smallest number in P_n .

Proof. Part (a) follows from Theorem 2.2 with $X = I_\infty$ and Theorem 1.1. To prove part (b) consider the sequence $\{v_n\}$ in $M_F[0, 1]$. Since $M_F[0, 1]$ is compact and metrizable in the vague topology any subsequence of $\{v_n\}$ has a convergent subsequence. Let $\{v_{n_k}\}$ be a subsequence of $\{v_n\}$ converging to $\mu \in M_F[0, 1]$. By Proposition 2.4 and Theorem 2.3,

$$\mu = \alpha_0 v_0 + \alpha_1 v_1 + \alpha_2 v_2 + \cdots + \alpha_\infty v_\infty$$

for some sequence of nonnegative real numbers $\{\alpha_n\}$ such that $\sum_{n=0}^{\infty} \alpha_n + \alpha_\infty = 1$. For ease of notation, denote by $v_n(f)$ the integral $\int f(x) v_n(dx)$. Then for any continuous function f on $[0, 1]$,

$$(13) \quad v_{n_k}(f) \rightarrow \alpha_0 v_0(f) + \alpha_1 v_1(f) + \alpha_2 v_2(f) + \cdots + \alpha_\infty v_\infty(f)$$

as $k \rightarrow \infty$. Let f be continuous on $[0, 1]$, $f \equiv 0$ outside of the open interval A_0 and $f(y_0) = 1$ (where as before $F(y_0) = y_0$ is the unique fixed point and smallest number in the period 1 orbit). By Theorem 1.1 $v_n(f) = 0$ when $n \geq 1$ and $v_0(f) = 1$. It follows that $\alpha_0 = 0$. Choosing a continuous function f such that $f(y_n) = 1$ and $f \equiv 0$ outside of the open set A_{n-1} and using a similar argument shows that $\alpha_n = 0$ for each integer n . Since μ is a probability measure, it follows that $\alpha_\infty = 1$. Thus $\mu = v_\infty$. Since every subsequence of $\{v_n\}$ has a subsequence converging to v_∞ , it follows that

$$(14) \quad v_n \rightarrow v_\infty$$

in the vague topology of $M_F[0, 1]$, which is equivalent to part (b) of Proposition 2.5. \square

Part (a) of Proposition 2.5 may be understood in an intuitive way. Consider a system with initial "state" $x_0 \in [0, 1]$, whose state at integer time n is given recursively by $x_n = F(x_{n-1})$. Let f be an observable, i.e., a continuous function from $[0, 1]$ to \mathbf{R} . Proposition 2.5 (a) then says that the time average $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=0}^{N-1} f(F^k(x_0))$ of f is equal to the "phase space average"

$\int f(x) \nu_\infty(dx)$ of f for any initial state x_0 which is not eventually periodic. The identification of time averages with phase space averages is the theoretical foundation of the statistical mechanical derivation of thermodynamics.

One way to measure chaos is to calculate Liapunov exponents. These exponents measure the rate of separation of nearby points under iterations of the map defining a dynamical system. When x and x_0 are close,

$$F^n(x) - F^n(x_0) \approx D_{x_0} F^n \cdot (x - x_0).$$

If we also require that $F^n(x) - F^n(x_0) \approx \exp(n\lambda(x_0))$ asymptotically as n increases (for x_0 "infinitesimally close" to x), then a natural definition for $\lambda(x_0)$ is

$$\begin{aligned} \lambda(x_0) &\equiv \lim_{n \rightarrow \infty} \frac{1}{n} \log \left| D_{x_0} F^n \cdot (x - x_0) \right| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \left| F'(F^{n-1}(x_0)) \cdot F'(F^{n-2}(x_0)) \cdots F'(x_0)(x - x_0) \right| \\ (15) \quad &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \log \left| F'(F^k(x_0)) \right| \end{aligned}$$

assuming, of course, that F is differentiable at $F^k(x_0)$ for all $k \geq 0$. In our case, F is differentiable at all but countably many points. For a point x in $[0, 1]$ where F fails to be differentiable, let us make the convention that $D_x F \equiv 1$, the smaller in magnitude of the one-sided derivatives. If x_0 is in I_∞ , (15) can be calculated directly or via Proposition 2.5 (a) with $f(x) \equiv \log |F'(x)|$ so that

$$(16) \quad \lambda(x_0) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \log \left| F'(F^k(x_0)) \right| = \int \log |F'(x)| \nu_\infty(dx).$$

In either case, $\lambda(x_0) = 0$ for all x_0 in I_∞ because $F'(x) \equiv 1$ on I_∞ . Similar

reasoning shows that if x is eventually in the period 2^n orbit of F , i.e., $x \in \bigcup_{k=0}^{\infty} F^{-k}(P_n)$ then

$$(17) \quad \lambda(x) = \frac{1}{2^n} \log \left(\frac{7}{3} \right).$$

There is no generally accepted mathematical definition of a chaotic map. One widely accepted definition [9] requires that the Liapunov exponent be positive. It follows that $F: I_{\infty} \rightarrow I_{\infty}$ is not chaotic in this sense. It is not difficult to verify that F is also not chaotic according to the definition given in Devaney [7]. The qualitative behavior that nearby points in I_{∞} do not separate with increasing iterations of F is also manifested by the fact, established in Theorem 1.1 (a), that F is a permutation on the intervals I_n^k for any fixed n .

A measurable partition of a probability space (X, ν) is a collection

$$\xi = \{B_1, B_2, \dots, B_n\}$$

of measurable subsets of X whose union is X and which are pairwise disjoint. The entropy $H(\xi)$ is given by

$$(18) \quad H(\xi) = - \sum_{i=1}^n \nu(B_i) \log [\nu(B_i)]$$

with the convention that $0 \log 0 = 0$. If $T: X \rightarrow X$ is a fixed measurable transformation, define ξ^m to be the measurable partition of X consisting of all sets of the form $B_{i_1} \cap T^{-1}(B_{i_2}) \cap \dots \cap T^{-m-1}(B_{i_m})$ where $B_{i_k} \in \xi$. For a T -invariant probability measure ν , the entropy $h(\nu, \xi)$ of ν relative to ξ is defined by

$$(19) \quad h(\nu, \xi) = \lim_{m \rightarrow \infty} \frac{1}{m} H(\xi^m),$$

where it can be shown that the limit in (19) exists. The entropy $h(\nu)$ is then defined by

$$(20) \quad h(\nu) = \sup h(\nu, \xi)$$

where the supremum is over all measurable partitions of X . The entropy $h(\nu)$ is sometimes called the Kolmogorov-Sinai invariant and it measures the asymptotic rate of creation of information by iterating T . It is invariant under

measure preserving isomorphisms. If X is a compact metric space (with the Borel σ -algebra) and T is continuous, then [14]

$$(21) \quad h(\nu) = \lim_{\text{diam } \xi \rightarrow 0} h(\nu, \xi),$$

where $\text{diam } \xi = \max_i \{\text{diameter of } B_i \in \xi\}$.

We apply (21) to $F: I_\infty \rightarrow I_\infty$ with the invariant probability measure ν_∞ . For any positive integer n , let $\xi(n) = \{I_n^1 \cap I_\infty, I_n^2 \cap I_\infty, \dots, I_n^{2^n} \cap I_\infty\}$. As $n \rightarrow \infty$, $\text{diam } \xi(n) \rightarrow 0$. By Theorem 1.1, $(\xi(n))^m = \xi(n)$ for every m and

$$\begin{aligned} H(\xi(n)) &= - \sum_{k=1}^{2^n} \nu_\infty(I_n^k \cap I_\infty) \log [\nu_\infty(I_n^k \cap I_\infty)] \\ &= - \sum_{k=1}^{2^n} 2^{-n} \log 2^{-n} \\ (22) \quad &= n \log 2. \end{aligned}$$

Therefore $h(\nu_\infty, \xi(n)) = \lim_{m \rightarrow \infty} \frac{1}{m} n \log 2 = 0$ for all n . Thus

$$h(\nu_\infty) = \lim_{n \rightarrow 0} h(\nu_\infty, \xi(n)) = 0.$$

Essentially the same argument shows that $h(\nu_\infty) = 0$ when we regard $F: [0, 1] \rightarrow [0, 1]$. The only modification needed is to add terms to the partition $\xi(n)$ which partition the complement of I_∞ in such a way that the diameters of these zero measure pieces decrease to zero as $n \rightarrow \infty$.

A similar calculation shows that $h(\nu_n) = 0$ for $F: [0, 1] \rightarrow [0, 1]$ for any nonnegative integer n . Because $[0, 1]$ is a compact metric space, the topological entropy $h_t(f)$ for our map $F: [0, 1] \rightarrow [0, 1]$ may be defined [4] as

$$h_t(F) = \sup \{h(\nu) : \nu \text{ is an ergodic } F\text{-invariant probability measure}\}.$$

From the above analysis, the topological entropy is clearly zero.

To what extent is the behavior of the function F generic? Let $T: [0, 1] \rightarrow [0, 1]$ be a continuous map with zero topological entropy, and let ν be an ergodic T -invariant probability measure on $[0, 1]$ which is not supported on any periodic orbit of T . Misiurewicz [2] pointed out that all periodic orbits of T have periods which are powers of 2. He proved that the dynamical system determined by T and ν on $[0, 1]$ is isomorphic to the adding machine on 2-adic integers explained in Theorem 1.2 together with the measure $\nu_\infty \circ h^{-1}$ (where h is the homeomorphism defined in Theorem 1.2). It follows that the dynamical system determined by T and ν on $[0, 1]$ is therefore isomorphic to the dynamical system determined by F and ν_∞ . An example of such a map with zero topological entropy is the function $x^2 - c_0$ discussed in the introduction.

ACKNOWLEDGMENTS

The authors would like to thank Professor Michal Misiurewicz for help in tracing the history of the “adding machine” and Professor David Ostroff for extensive discussions on the subject matter of this paper.

REFERENCES

- [1] MILNOR, J. On the concept of attractor. *Commun. Math. Phys.* 99 (1985), 177-195.
- [2] MISIUREWICZ, M. Structure of mappings of an interval with zero entropy. *Publ. Math. de l'Institut des Hautes Etudes Scientifiques* 53 (1981), 5-17.
- [3] ——— Invariant measures for continuous transformations of $[0, 1]$ with zero topological entropy. *Lecture notes in mathematics* No. 729, 144-152, edited by M. Denker and K. Jacobs, Springer-Verlag, 1979.
- [4] ECKMANN, J.-P. and D. RUELLE. Ergodic theory of chaos and strange attractors. *Rev. Mod. Phys.* 57 (1985), 617-656.
- [5] COLLET, P. and J.-P. ECKMANN. *Iterated maps on the interval as dynamical systems*, Birkhauser, 1980.
- [6] DELAHAYE, J.-P. Fonctions admettant des cycles d'ordre n importe quelle puissance de 2 et aucun autre cycle. *Compte Rendus de l'Académie des Sciences, Paris*, 291A (1980), 323-325.
- [7] DEVANEY, R.L. *An introduction to chaotic dynamical systems*. Addison-Wesley, 1989, 2nd ed.
- [8] SHAROVSKIĬ, A.N. Co-existence of the cycles of a continuous mapping of the line into itself. *Ukrain. Math. Zh.* 16 (1) (1964), 61-71 [in Russian].
- [9] RUELLE, D. *Chaotic evolution and strange attractors*. Cambridge U. Press, 1989.
- [10] BARNESLEY, M. *Fractals Everywhere*. Academic Press, 1988.
- [11] BROWN, J.R. *Ergodic theory and topological dynamics*. Academic Press, 1976.
- [12] FURSTENBERG, H. *Recurrence in ergodic theory and combinatorial number theory*. Princeton U. Press, 1981.
- [13] PHELPS, R.R. *Lectures on Choquet's theorem*. Van Nostrand, 1966.
- [14] BOWEN, R. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*. Lecture notes in mathematics No. 470. Springer-Verlag, 1975.

(Reçu le 15 février 1994)

Peter Collas

Department of Physics and Astronomy
California State University, Northridge
Northridge, CA. 91330-8268, U.S.A.

E-mail address: pcollas@huey.csun.edu

David Klein

Department of Mathematics
California State University, Northridge
Northridge, CA. 91330-8313, U.S.A.

E-mail address: dklein@huey.csun.edu