

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ERGODIC ADDING MACHINE ON THE CANTOR SET
Autor: Collas, Peter / Klein, David
Kurzfassung
DOI: <https://doi.org/10.5169/seals-61113>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

AN ERGODIC ADDING MACHINE ON THE CANTOR SET

by Peter COLLAS and David KLEIN

ABSTRACT. We calculate all ergodic measures for a specific function F on the unit interval. The supports of these measures consist of periodic orbits of period 2^n and the classical ternary Cantor set. On the Cantor set, F is topologically conjugate to an “adding machine” in base 2. We show that F is representative of the class of functions with zero topological entropy on the unit interval, already analyzed in the literature, and its behavior is therefore typical of that class.

I. INTRODUCTION

The dynamical behavior of the quadratic function $f_c(x) = x^2 - c$ has been extensively studied as the parameter c is varied. For example, $c_0 = 1.401155189\dots$ is the smallest value of c for which $f_c(x)$ has infinitely many distinct periodic orbits [1-3]. As c approaches this number through smaller values, the dynamical system, $x \rightarrow f_c(x)$, progresses through the famous period doubling route to chaos. When $c = c_0$, the dynamical behavior of $f(x) \equiv f_c(x)$ includes the following properties:

1. There is a Cantor set K which is an attractor and $f: K \rightarrow K$
2. All periodic points of f have period 2^n for some n .
3. There are periodic points which are arbitrarily close to K .
4. With the restriction of $f(x)$ to an appropriate interval I such that $f(I) \subset I$, there are just two possibilities for the orbit of a point $x_0 \in I$: either $f^k(x_0)$ is in a periodic orbit for some k , or $f^k(x_0)$ converges to K as k increases.
5. The restriction of f to K is topologically equivalent to a function on 2-adic integers which adds 1 to its argument (this “adding machine” will be described in detail below).