

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON HAUSDORFF-GROMOV CONVERGENCE AND A THEOREM OF PAULIN
Autor: Bridson, M.R. / Swarup, G. A.
Kapitel: Section 2: The Proof of Paulin's Theorem
DOI: <https://doi.org/10.5169/seals-61114>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 22.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Thus, as $i \rightarrow \infty$, we see that x'_i, y'_i, z'_i converge to the same point, say z' , on $[x, y]$. Thus $d(x_i, y'_i) + d(y_i, x'_i) - d(x_i, y_i)$ converges to zero. Since $d(x_i, z_i) + d(y_i, z_i) - d(x_i, y_i)$ also converges to zero, we have that $d(y'_i, z_i) + d(z_i, x'_i)$ converges to zero. Since $d(z_i, z'_i) \leq d(z'_i, y'_i) + d(y'_i, z_i) \leq 4\delta_i + d(y'_i, z_i)$ we see that the z_i converge to the point z' on our original geodesic segment $[x, y]$. Thus z , the midpoint of our arbitrary geodesic from x to y , coincides with the midpoint of our fixed geodesic. Repeating the argument we see that these geodesics must agree at a dense set of points, and hence everywhere. Since geodesic triangles in C_i are δ_i -slim, and geodesics in C all arise as limits of geodesics in C_i , we see that geodesic triangles in C must be 0-slim, and hence C is an **R**-tree. \square

Remark. If one has a sequence of δ_i -hyperbolic spaces C_i , with $C_i \rightarrow C$ and $\delta_i \rightarrow \delta > 0$, then one can extend the preceding argument to show that C is δ' -hyperbolic (with $\delta' = 19\delta$, for example).

SECTION 2: THE PROOF OF PAULIN'S THEOREM

In this section we shall prove the following theorem of F. Paulin [P4].

2.1 THEOREM (Paulin). *If Γ is a word hyperbolic group and $Out(\Gamma)$ is infinite, then Γ acts by isometries on an **R**-tree with virtually cyclic segment stabilizers and no global fixed points.*

In its outline, the proof given below is very similar to Paulin's original proof, except that we use Hausdorff-Gromov convergence instead of the equivariant Gromov convergence used by Paulin. In particular, this allows us to avoid the difficulties discussed in the next section.

Let S be a finite set of generators for Γ and let $X = X(\Gamma, S)$ denote the Cayley graph of Γ with respect to S , as defined in the introduction. Γ is the vertex set of X and receives the induced metric. The hypothesis that Γ is word hyperbolic means precisely that there exists $\delta > 0$ such that X is a δ -hyperbolic geodesic metric space. Note that with our definition of a Cayley graph, the endpoints of each edge are distinct, and there is at most one edge joining each pair of vertices; hence the action of Γ on itself by left multiplication can be extended linearly across edges in a unique way to give an isometric action of Γ on X .

The proof of Theorem 2.1 will be broken into a number of smaller results. We begin by noting that, because $Out(\Gamma)$ is infinite, we can choose a sequence

of automorphisms $\{\phi_i\}_{i \in \mathbb{N}}$ such that none of the ϕ_i is an inner automorphism and no two of the ϕ_i have the same image in $Out(\Gamma)$. For each $i \in \mathbb{N}$ we consider the function $f_i : X \rightarrow [0, \infty)$ defined by:

$$(2.2) \quad f_i(x) = \max_{s \in S} d(x, \phi_i(s)x) .$$

This function has been used by Bestvina in his study of degeneration of real hyperbolic structures [B], and our use of this function is similar to his. (A similar idea was used earlier in a different context by Thurston [T, Prop. 1.1].)

Note that f_i takes on integer values at vertices and midpoints of edges in X , and its restriction to half-edges is linear. It follows that f_i attains its infimum (which is an integer) at some point, $x_i \in X$ say. (In the case where Γ is not virtually cyclic one can also see this by showing that f_i is a proper map, i.e., a map with the property that the inverse image of a compact set is compact.)

Let

$$(2.3) \quad \begin{aligned} \lambda_i &= \max_{s \in S} d(x_i, \phi_i(s)x_i) \\ &= \inf_{x \in X} \max_{s \in S} d(x, \phi_i(s)x) . \end{aligned}$$

We fix a definite choice of points x_i with the above property.

For future reference, we note that by passing to a subsequence of the ϕ_i we may assume there is a single element $s_0 \in S$ such that $\lambda_i = d(x_i, \phi_i(s_0)x_i)$ for all $i \in \mathbb{N}$. We also note that with the above choice of x_i , the triangle inequality yields:

$$(2.4) \quad d(x_i, \phi_i(\gamma)x_i) \leq \lambda_i d(e, \gamma) .$$

Following Paulin, we next note that because $Out(\Gamma)$ is infinite, the sequence λ_i must be unbounded. For suppose that there were a uniform bound, ρ say, on the value of λ_i . Then for any vertex $y_i \in X$ closest to x_i , we would have $d(e, y_i^{-1}\phi_i(s)y_i) = d(y_i, \phi_i(s)y_i) \leq \rho + 2$ for all $s \in S, i \in \mathbb{N}$. But there are only finitely many vertices in the ball of radius $\rho + 2$ about e , so this bound would imply the existence of integers $n \neq m$ such that $y_n^{-1}\phi_n(s)y_n = y_m^{-1}\phi_m(s)y_m$ for all $s \in S$. Whence ϕ_n and ϕ_m would be equal in $Out(\Gamma)$, contrary to hypothesis. Thus we have shown that the sequence of numbers $\{\lambda_i\}_{i \in \mathbb{N}}$ is unbounded, so we may pass to a subsequence $\{\lambda_n\}_{n \in \mathbb{N}}$ which is *strictly increasing* and assume that $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$.

Consider the sequence of metric spaces $X_k = (X, d_k)$, where $d_k := d/\lambda_k$ is the original metric on X scaled down by λ_k . In what follows we shall intermittently use both the original metric d and the scaled metric d_k , specifying which on each occasion and, where appropriate, using the formal notation (Y, d) for a metric space which consists of the set Y together with a distance function d . But for the moment, the most important distinction between the X_k will be that we shall regard Γ as acting on X_k via ϕ_k , and think of our chosen point x_k , at which the minimax λ_k is attained, as a *basepoint* in X_k . More precisely, we consider the sequence of pointed Γ -spaces (X_k, x_k) , where the action of $\gamma \in \Gamma$ on X_k is $x \mapsto \phi_k(\gamma)x$.

We wish to use the hyperbolic nature of X_k to approximate it by a sequence of star-like compact subsets $X_k(P_i)$ centred at x_k . To this end, we fix a sequence of finite subsets $\{e\} = P_0 \subseteq P_1 \subseteq P_2 \cdots \subseteq P_i \subseteq \cdots$ which exhaust Γ . Let $n_i = |P_i|$ denote the cardinality of P_i . The desired subsets of X_k are defined inductively as follows: $X_k(P_0) = \{x_k\}$, and $X_k(P_i)$ is the union of $n_i - 1$ geodesic segments, those in $X_k(P_{i-1})$ together with a choice of geodesic segment from x_k to each element of $\{\phi_k(\gamma)x_k \mid \gamma \in P_i - P_{i-1}\}$.

We next ‘fatten-up’ each of the sets $X_k(P_i)$ by taking its closed δ -neighbourhood in the metric d . Henceforth we shall denote this neighbourhood V_k^i . Let $d_{i,k}$ be the induced *path metric* on V_k^i . As we discussed in Section 1, $(V_k^i, d_{i,k})$ is a geodesic metric space. It is also important to notice that the induced path metric which V_k^i receives from d_k is $d_{i,k}/\lambda_k$. The following lemma is suggested by an argument of B. Bowditch [Bo].

2.5 LEMMA. *With the above notation, for all $x, y \in V_k^i$ we have:*

$$d(x, y) \leq d_{i,k}(x, y) \leq d(x, y) + 4\delta.$$

Proof. The left-most inequality comes from the general fact that for any subspace of a geodesic metric space the induced metric is dominated by the induced path metric. In order to establish the other inequality, we first note that $X_k(P_i)$ is δ -convex in (X_k, d) , in the sense that if a geodesic segment in X_k joins a pair of points $x, y \in X_k(P_i)$, then this geodesic segment lies entirely within the closed δ -neighbourhood V_k^i of $X_k(P_i)$.

Given $x, y \in V_k^i$, we fix points $z, w \in X_k(P_i)$ closest to x and y respectively. (Such points are not unique in general.) Let $[x, z]$, $[z, w]$ and $[w, y]$ be choices of geodesic segments joining x to z , z to w and w to y , respectively. Each is contained in V_k^i , and hence so is the broken geodesic $[x, z, w, y]$ obtained by concatenating them. The length of this broken geodesic is at most $d(z, w) + 2\delta \leq d(x, y) + 4\delta$. Hence $d_{i,k}(x, y) \leq d(x, y) + 4\delta$. \square

The subspace V_k^i forms a good substitute for the notion of a convex hull for $\phi_k(P_i)x_i$ in X_k . According to the above lemma, geodesics in $(V_k^i, d_{i,k})$ are $(1, 4\delta)$ -quasigeodesics in (X_k, d) , and hence by [GH, p. 82] there exists a constant $\eta = \eta(\delta)$ (independent of k, i) such that geodesic triangles in $(V_k^i, d_{i,k})$ are η -slim. Thus we have proved the first part of:

2.6 LEMMA. *There exists a constant $\eta = \eta(\delta)$ such that, for all $k \in \mathbb{N}$, with respect to the path metric $d_{i,k}$ on V_k^i , geodesic triangles in V_k^i are η -slim. Moreover, for fixed i , with respect to the (scaled) path metrics $d_{i,k}/\lambda_k$, the metric spaces $\{V_k^i\}_{k \in \mathbb{N}}$ are uniformly compact.*

Proof. It remains to prove the assertion of the second sentence. We follow an argument of Bestvina [B]. Until further notice we work with the metric d . Let μ_i be the maximum of the integers $\{d(e, \gamma) \mid \gamma \in P_i\}$. Each of the geodesic segments used to define $X_k(P_i)$ has length at most $\mu_i \lambda_k$ (by (2.4)). Therefore, given $\varepsilon > 0$, we can cover $X_k(P_i)$ by $2n_i \mu_i / \varepsilon$ segments of length at most $\lambda_k \varepsilon / 2$. (Recall that $n_i = |P_i|$.) Hence, if $\lambda_k \varepsilon > 2\delta$, then in order to cover V_k^i we need at most $2n_i \mu_i / \varepsilon$ balls of radius $\lambda_k \varepsilon$. But we arranged that $\lambda_k \rightarrow \infty$ as $k \rightarrow \infty$, so this is true for large k .

Now we change viewpoints and work with the scaled metric d_k on X_k , and the induced path metric on V_k^i . In this setting, the preceding argument shows that for large k one needs only $2n_i \mu_i / \varepsilon$ balls of radius ε to cover V_k^i . Since the path metric on V_k^i and the restriction to V_k^i of d_k differ by at most an additive constant of $4\delta/\lambda_k$, we have thus established the existence of a uniform ε -count for the $\{V_k^i\}_{k \in \mathbb{N}}$ both when equipped with the restriction of the metrics d_k and when equipped with the induced path metrics. Because they are *path* metric spaces, a uniform ε -count also yields a bound on the diameter of the V_k^i . \square

Continuing with the proof of Paulin's theorem, we fix an integer j and suppose that we are given a positive constant ε . According to the preceding lemma, we can choose ε -nets $N_\varepsilon(k, j)$ for V_k^j on whose cardinalities there is a bound independent of k . We may also assume that the set $N_\varepsilon(k, j)$ includes $\phi_k(P_j)x_k$. Since, for fixed j , the $N_\varepsilon(k, j)$ are finite metric spaces of uniformly bounded cardinality and diameter, we can pass to a subsequence (using a diagonal type argument, as in Section 1) so as to assume that, for all $\gamma, \gamma' \in P_j$, the sequence of numbers $d_{j,k}(\phi_k(\gamma)x_k, \phi_k(\gamma')x_k)$ converges as $k \rightarrow \infty$. Passing to a further subsequence which is convergent in the Hausdorff-Gromov topology we obtain a limit metric space $L_{\varepsilon,j}$ (whose cardinality will be no greater than that of the $N_\varepsilon(k, j)$). As a basepoint in the

limit space we choose the limit of the sequence x_k , and we christen this point x_∞ . For each $\gamma \in P_j$, we denote the limit of the sequence $\phi_k(\gamma)x_k$ by γx_∞ .

We next take an $\varepsilon/2$ -net for V_k^j which is constructed so as to include the previously chosen ε -net. Passing to a subsequence if necessary, we obtain a finite limit metric space $L_{\varepsilon/2,j}$. We proceed in this manner, taking finer ε -nets, and at each stage including the previous (coarser) ones and extracting convergent subsequences to obtain finite limit metric spaces. The natural inclusions of each ε -net into its refinements gives a natural identification of points in the limit, so it is not too abusive a notation to write:

$$L_{\varepsilon,j} \subset L_{\varepsilon/2,j} \cdots \subset L_{\varepsilon/2^n,j} \subset \cdots$$

We define L_j to be the direct limit of this sequence, that is, $L_j = \bigcup \{L_{\varepsilon/2^n,j} \mid n \in \mathbf{N}\}$. We denote by \hat{L}_j the metric completion of L_j . Since the diameters of the V_k^j are uniformly bounded in the scaled metrics, we see that \hat{L}_j is a complete space of finite diameter, and hence is compact.

By choosing a diagonal type subsequence and renumbering, we obtain the following array of spaces with convergence in both the horizontal and vertical directions:

$$\begin{array}{ccccccc} N_\varepsilon(1,j) & \subseteq & N_{\varepsilon/2}(2,j) & \subseteq & \cdots & \subseteq & N_{\varepsilon/2^n}(1,j) & \subseteq & \cdots & \subseteq & V_1^j & \subseteq & X_1 \\ N_\varepsilon(2,j) & \subseteq & N_{\varepsilon/2}(2,j) & \subseteq & \cdots & \subseteq & N_{\varepsilon/2^n}(2,j) & \subseteq & \cdots & \subseteq & V_2^j & \subseteq & X_2 \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ N_\varepsilon(m,j) & \subseteq & N_{\varepsilon/2}(m,j) & \subseteq & \cdots & \subseteq & N_{\varepsilon/2^n}(m,j) & \subseteq & \cdots & \subseteq & V_m^j & \subseteq & X_m \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ L_{\varepsilon,j} & \subseteq & L_{\varepsilon/2,j} & \subseteq & \cdots & \subseteq & L_{\varepsilon/2^n,j} & \subseteq & \cdots & \subseteq & \hat{L}_j \end{array}$$

Our next goal is to show that as $k \rightarrow \infty$ the V_k^j actually converge to \hat{L}_j in the Hausdorff-Gromov topology. We have that $N_{\varepsilon/2^n}(m,j)$ is $\varepsilon/2^{n-1}$ close to V_m^j for all m . After passing to yet another diagonal type subsequence, we may assume that $N_{\varepsilon/2^n}(m,j)$ is $\varepsilon/2^{m-1}$ close to $L_{\varepsilon/2^n,j}$ for all $m \geq n$. Thus V_m^j and $L_{\varepsilon/2^n,j}$ are $\varepsilon/2^{n-2}$ close for $m \geq n$. On the other hand, $L_{\varepsilon/2^n,j}$ and $L_{\varepsilon/2^{n+1},j}$ are $\varepsilon/2^{n+1}$ close (since any choice of $\varepsilon/2^n$ and $\varepsilon/2^{n+1}$ nets of V_k^j are $\varepsilon/2^{n+1}$ close). Thus $L_{\varepsilon/2^n,j}$ is $\sum_{i \geq n} \varepsilon/2^i$ close to L_j and \hat{L}_j . Hence V_n^j and \hat{L}_j are $\varepsilon/2^{n-3}$ close, so V_n^j converges to \hat{L}_j , in the Hausdorff-Gromov topology, as $n \rightarrow \infty$.

Notice that, by (1.9) and (2.6), the spaces \hat{L}_j are \mathbf{R} -trees of finite diameter, because V_k^j is η/λ_k -hyperbolic and $\lambda_k \rightarrow \infty$. It is also useful to observe that \hat{L}_j is spanned by γx_∞ , with $\gamma \in P_j$. Furthermore, the $X_k(P_j)$ themselves converge to \hat{L}_j because $X_k(P_j)$ and V_k^j are $4\delta/\lambda_k$ -close and $\lambda_k \rightarrow \infty$. However, in what follows it is most convenient to still work with V_k^j rather than $X_k(P_j)$ when we need to take a choice of geodesic between two points of $X_k(P_j)$. Also, because the scaled path metric on V_k^j and the induced metric d_k/λ_k differ only by $4\delta/\lambda_k$, which tends to 0 as $k \rightarrow \infty$, henceforth it is not important to keep track of the difference between these two metrics.

By construction, all of our $\varepsilon/2^n$ -nets include the set $\{\phi(\gamma)x_k \mid \gamma \in P_j\}$ and each of the sequences $d_k(\phi(\gamma)x_k, \phi(\gamma')x_k)$ converges. Thus, if we denote by $x_\infty \in \hat{L}_j$ the 'limit' of the x_k , and by γx_∞ the limit of the $\phi(\gamma)x_k$, then we see that $d(\gamma x_\infty, \gamma' x_\infty)$ (distance in \hat{L}_j) is independent of j . Since the tree \hat{L}_j is the convex hull of the points γx_∞ , we can define an isometric embedding of \hat{L}_j into \hat{L}_{j+1} for all j and hence obtain an \mathbf{R} -tree by taking the direct limit of the resulting system of inclusions. We denote the direct limit metric space with basepoint (which as the limit of \mathbf{R} -trees is itself an \mathbf{R} -tree) by $(X_\infty; x_\infty)$. The final important observation to make is that Γ acts isometrically on X_∞ , because it acts isometrically on the subset $\{\gamma x_\infty\}_{\gamma \in \Gamma}$ (by left translation), and the convex hull of this subset is the whole of X_∞ .

Let us now examine the nature of the action of Γ on X_∞ . We claim that it has the following properties:

- (1) There is no point of X_∞ whose stabilizer is the whole of Γ .
- (2) The stabilizer of every non-trivial segment in X_∞ is virtually cyclic.

To see that (1) is true, let us see what would happen if it were to fail. Suppose that Γ were to stabilize a point $z_\infty \in X_\infty$. We fix a segment $z_\infty \in [\gamma x_\infty, \gamma' x_\infty] \subseteq \hat{L}_j$. Up to the taking of subsequences, we have that the closures in \hat{L}_j of the images of the geodesic segments $[\gamma x_k, \gamma' x_k] \subseteq V_k^j$ converge (in the Hausdorff metric) to $[\gamma x_\infty, \gamma' x_\infty]$, and we fix points $z_k \in [\gamma x_k, \gamma' x_k]$ which converge to z_∞ . We then choose j large enough to ensure that $S \subset P_j$ (recall that S is our fixed finite generating set for Γ), and l large enough to ensure that $P_j P_j \subset P_l$.

We have, for every $s \in S$, geodesics $[s\gamma x_k, s\gamma' x_k] := s \cdot [\gamma x_k, \gamma' x_k]$ in V_k^l , and (by definition of the action on X_∞) the closures of their images in $\hat{L}_l \subseteq X_\infty$ converge to $[s\gamma x_\infty, s\gamma' x_\infty]$. Moreover, $\{s z_k\}_{k \in \mathbf{N}}$ converges to $s \cdot z_\infty = z_\infty$, so for large k we have that $d_k(s \cdot z_k, z_k) < 1/4$ in the *scaled*

metric of X_k . Hence $d(s \cdot z_k, z_k) < \lambda_k/4$, for large k , in the original metric on X_k . But this contradicts the definition of λ_k .

Remark. The preceding argument actually shows that for every finite set $P \subseteq \Gamma$ which fixes z_∞ , given any $\varepsilon > 0$ one has that for k sufficiently large z_k and γz_k are ε -close, in the scaled metric d_k , for every $\gamma \in P$.

We next need to show that segment stabilizers are virtually cyclic. This seems to be the place where some sort of discreteness assumption on Γ is needed. In the classical real-hyperbolic case, Margulis' Lemma implies the result for discrete actions (see [B] and [P2]). Since we are using Cayley graphs and the group actions are (almost) free there is still some sort of discreteness and Paulin gives a delicate argument to show that segment stabilizers are virtually cyclic. The following algebraic lemma is taken from [P4]:

2.7 LEMMA. *Let G be a finitely generated group. If the set of commutators $\{aba^{-1}b^{-1} \mid a, b \in G\}$ is finite, then G is virtually abelian.*

Proof. The action of G on itself by conjugation determines a map $G \rightarrow \text{Aut}(\Gamma)$, whose image is $\text{Inn}(G)$ and whose kernel is the centre of G ; it suffices to prove that $\text{Inn}(G)$ is finite. If A is a finite generating set for G , then the action of $g \in G$ by conjugation is determined by its action on the elements $a \in A$. But $g^{-1}ag = (g^{-1}aga^{-1})a$, and by hypothesis there are only finitely many possibilities, M say, for the commutator $g^{-1}aga^{-1}$. Hence the cardinality of $\text{Inn}(G)$ is at most $M^{|A|}$. \square

We proceed with the proof of assertion (2) on segment stabilizers. We call a subgroup *large* if it contains a non-abelian free subgroup (for hyperbolic groups this is equivalent to not having a cyclic subgroup of finite index). Suppose that a large subgroup G of Γ stabilizes a non-trivial segment $e \subseteq X_\infty$. If e is finite, then a subgroup of index 2 in G fixes e pointwise. If e is infinite, a subgroup of index 2 in G acts as translations on a ray in e and thus a large subgroup of G , obtained by taking commutators, fixes a segment of positive length in e pointwise. Thus, in any case, if a large subgroup of Γ stabilizes a segment, then a (perhaps smaller) large subgroup of Γ fixes a segment e of positive length pointwise. Therefore, in order to complete the proof of Paulin's theorem, it suffices to show that if a subgroup of Γ fixes a segment of X_∞ pointwise, then that subgroup is virtually cyclic. Let D denote the length of such a segment which is fixed pointwise by the subgroup $G \subset \Gamma$, and let z and z' denote the endpoints of the segment.

We fix $\varepsilon > 0$ small (to be estimated later) compared to D , and k so large that if $z_k, z'_k \in X_k$ correspond to $z, z' \in X$ then $|d(z_k, z'_k) - D| < \varepsilon$. We fix

a geodesic segment $[z_k, z'_k]$ from z_k to z'_k in X_k . Given any finite subset $P \subseteq G$, we choose a finite subset $Q \subseteq G$ which contains all products of length ≤ 4 in s, t, s^{-1}, t^{-1} , as s and t vary over P . We choose k large enough so that z_k, z'_k are moved by less than ε by each $\gamma \in Q$ with respect to the scaled metric $d_k = d/\lambda_k$. If $D > 3\varepsilon + (24\delta/\lambda_k)$ then if we omit segments of d -length $\lambda_k\varepsilon + 12\delta$ from the ends of $[z_k, z'_k]$, the remaining sub-segment is non-empty; call this segment C_k . We assume that ε is small enough to satisfy the above inequality; we shall place further restrictions on ε later.

Now we use the original metric d on X_k . From the proof 'slim \Rightarrow thin' (see [Sho] p. 17), if $x \in C_k$ then γx is within 12δ of $[z_k, z'_k]$. We denote by γ_*x the projection of γx on $[z_k, z'_k]$. Of course, the 'projection' is not uniquely defined, but the preceding sentence is true no matter which closest point on $[z_k, z'_k]$ one chooses — we fix a definite choice for each $x \in C_k$, thus defining a map $\gamma_*: C_k \rightarrow [z_k, z'_k]$ for each γ . Next, we omit segments of length $5(\lambda_k\varepsilon + 12\delta)$ from the ends of $[z_k, z'_k]$ and denote the remaining long segment by $E_k \subseteq C_k$. The map $C_k \rightarrow [z_k, z'_k]$ just defined restricts to a map $E_k \rightarrow [z_k, z'_k]$; we continue to denote this map by γ_* . Notice that this map is a 24δ -isometry, that is to say, it distorts distances by at most an additive constant of 24δ ; in fact it is 24δ close to a translation of E_k along $[z_k, z'_k]$. (Here, and in what follows, the terminology η -close is used to describe functions f, g with the same domain such that $d(f(x), g(x)) < \eta$ for all points in their common domain.)

Note that on E_k the maps $s_*, s_*t_*, s_*t_*(s^{-1})_*, s_*t_*(s^{-1})_*(t^{-1})_*$ etc. are well-defined and uniformly close to translations. Choose $M = \text{Max}\{5(\lambda_k\varepsilon + 12\delta), 600\delta\}$. We will denote by e_k the segment obtained from $[z_k, z'_k]$ by omitting segments of length M from the ends. We have $e_k \subset E_k$. To make sure that $e_k \neq \emptyset$ we assume $D - \varepsilon > 5\varepsilon + (60\delta/\lambda_k)$, we also assume $D - \varepsilon > (600\delta/\lambda_k)$. Since $\lambda_k \rightarrow \infty$, we can choose large enough k and small enough ε so that the above conditions are satisfied.

We shall consider the restrictions $\gamma_*: e_k \rightarrow C_k$ to e_k of the maps γ_* defined above; we retain the notation γ_* for these restricted maps. Our goal is to obtain a bound (independent of $|Q|$) on the number commutators $tst^{-1}s^{-1}$ in Q by estimating how close the action of such a commutator on e_k is to the identity map. We first compare $t_*s_*(t^{-1})_*(s^{-1})_*$ to $tst^{-1}s^{-1}$. Observe that, since the maps s and s_* are 12δ close, ts and $t(s_*)$ are 12δ close (the left-action of Γ on X_k is by isometries in the metric d). Hence, $(ts)_*$ and t_*s_* are 36δ close. Comparing successively $tst^{-1}s^{-1}$, $(tst^{-1}s^{-1})_*$, $(ts)_*(t^{-1}s^{-1})_*$, $t_*s_*(t^{-1})_*(s^{-1})_*$ shows that $tst^{-1}s^{-1}$ and $t_*s_*(t^{-1})_*(s^{-1})_*$ are $(12 + 36 + 108)\delta$ close.

Next, we compare $t_*s_*(t^{-1})_*(s^{-1})_*$ to the identity map on e_k . Since $s_*(t^{-1})_*$ and $(t^{-1})_*s_*$ are 72δ close to the same translation, and translations commute, we have that $t_*s_*(t^{-1})_*(s^{-1})_*$ and $t_*(t^{-1})_*s_*(s^{-1})_*$ are $(144 + 24)\delta$ close. Moreover, $t_*(t^{-1})_*$ and $s_*(s^{-1})_*$ are 36δ close to the identity. Thus $t_*(t^{-1})_*s_*(s^{-1})_*$ is 108δ close to the identity. Hence $t_*s_*(t^{-1})_*(s^{-1})_*$ is 276δ close to the identity. Combining this with the estimate in the previous paragraph we have that the restriction of $(tst^{-1}s^{-1})$ to e_k is 532δ close to the identity on e_k . Therefore, a vertex close to the midpoint of e_k is moved by less than $532\delta + 2$ by $tst^{-1}s^{-1}$. Thus $tst^{-1}s^{-1}$ lies in the ball of radius $532\delta + 2$ about the identity in Γ , and we have the desired bound on the number of commutators in the arbitrary finite subset $P \subset G$.

Now Lemma 2.7 implies that G is virtually abelian. But every abelian subgroup of a hyperbolic group is virtually cyclic. Hence the segment stabilizers for the action of Γ on X_∞ must be virtually cyclic. This completes the proof of Paulin's theorem. \square

SECTION 3: CONVEX HULLS

A subset Σ of a geodesic metric space X is said to be *geodesically convex* if for all $p, q \in \Sigma$ every geodesic segment from p to q is completely contained in Σ . Given a bounded set $Y \subset X$, perhaps the most natural way to define its convex hull is as the intersection of all geodesically convex sets containing Y .

If X is simply connected and non-positively curved then round balls are geodesically convex and hence the convex hull of a bounded set is bounded. However, for more general geodesic metric spaces, even δ -hyperbolic spaces, it may happen that the convex hull of a finite set is the whole of the ambient space X . The following example illustrates how general this problem is.

3.1 PROPOSITION. *Given any finitely generated group Γ there exists a finite generating set S and a finite subset $Y \subset \Gamma$ such that the convex hull of Y in the Cayley graph $X(\Gamma, S)$ is the whole of $X(\Gamma, S)$.*

Proof. Let A be any finite generating set for Γ , and take S to be the set of those elements of Γ which are a distance 1 or 4 from the identity in the Cayley graph of Γ with respect to S . Let Y be the set of elements of Γ which are a distance at most 3 away from the identity in the Cayley graph associated to A .