

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PROOF OF MARGULIS' THEOREM ON VALUES OF QUADRATIC FORMS, INDEPENDENT OF THE AXIOM OF CHOICE
Autor: Dani, S. G.

Bibliographie
DOI: <https://doi.org/10.5169/seals-61104>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

in $C(X)$ this readily implies that $n_k^{-1} \sum_{i=0}^{n_k-1} f \circ \varphi^i(x_0)$ converges for all $f \in C(X)$; let c_f be the limit corresponding to f . Then it can be verified that $\Lambda: C(X) \rightarrow \mathbf{R}$ defined by $\Lambda(f) = c_f$, for all $f \in C(X)$, is a φ -invariant integral on $C(X)$. Also clearly Λ is not identically zero and therefore by our observations above, the support, say X' , is a nonempty closed φ -invariant subset of X and further $C(X')$ admits an integral with full support (namely X') which is invariant under the restriction of φ to X' . Replacing X as in the hypothesis by X' we may without loss of generality assume that $C(X)$ admits a φ -invariant integral whose support is X ; in the rest of the argument we let Λ be any such integral.

Now suppose that there do not exist any recurrent points for φ . Let $\rho(\cdot, \cdot)$ be the metric on X . Let θ be the function on X defined by $\theta(x) = \inf\{\rho(\varphi^i(x), x) \mid i = 1, 2, \dots\}$, for all $x \in X$. There being no recurrent points means that $\theta(x) > 0$ for all $x \in X$. For each natural number k let $E_k = \{x \in X \mid \theta(x) \geq 1/k\}$. Then each E_k is a closed subset of X and $X = \cup E_k$. Therefore by the Baire category theorem there exists a k such that E_k has an interior point in X . In particular, there exists an open ball, say A , of radius at most $1/3k$ contained in E_k . The definition of E_k and the condition on the radius of A then imply that the sets $\varphi^i(A)$, $i \in \mathbf{Z}$, are mutually disjoint. Now let $x \in A$ and let $f \in C(X)^+$ be such that $f(x) > 0$ and the support of f (the closure of the set $\{y \in X \mid f(y) > 0\}$) is contained in A . For each natural number n let $S_n(f) = \sum_{i=0}^{n-1} f \circ \varphi^i \in C(X)$. The disjointness of $\varphi^i(A)$, $i \in \mathbf{Z}$, implies that, for any n , $\|S_n(f)\| = \|f\|$. Also, by the φ -invariance of Λ we have $\Lambda(S_n(f)) = n\Lambda(f)$. Hence $\Lambda(f) = \Lambda(S_n(f))/n \leq \|S_n(f)\| \Lambda(1_X)/n = \|f\| \Lambda(1_X)/n$ for all n , where 1_X denotes the constant function with value 1. But this implies that $\Lambda(f) = 0$ contradicting the assumption that the support of Λ is the whole of X . This proves the proposition.

Acknowledgement: Thanks are due to J. Aaronson and M.G. Nadkarni for a discussion on recurrent points and to Gopal Prasad and Nimish Shah for useful comments on an earlier version.

REFERENCES

- [DM-1] DANI, S.G. and G.A. MARGULIS. Values of quadratic forms at primitive integral points. *Invent. Math.* 98 (1989), 405-424.
- [DM-2] DANI, S.G. and G.A. MARGULIS. Values of quadratic forms at integral points; an elementary approach. *L'Enseignement Math.* 36 (1990), 143-174.

- [DM-3] DANI, S.G. and G.A. MARGULIS. Limit distributions of orbits of unipotent flows and values of quadratic forms. *Advances in Soviet Math.* 16 (1993), 91-137.
- [DGS] DENKER, M., C. GRILLENBERGER and K. SIGMUND. *Ergodic Theory on Compact Spaces*. Springer-Verlag, 1976.
- [D] DIEUDONNÉ, J. *Foundations of Modern Analysis*. Academic Press, 1969.
- [H] HENLE, J.M. *An Outline of Set Theory*. Springer-Verlag, 1986.
- [M] MAÑÉ, R. *Ergodic Theory and Differentiable Dynamics*. Springer-Verlag, 1983.
- [M-1] MARGULIS, G.A. Formes quadratiques indéfinies et flots unipotents sur les espaces homogènes. *C.R. Acad. Sci. Paris, Série I*, 304 (1987), 249-253.
- [M-2] ——— Discrete subgroups and ergodic theory; in: *Number Theory, Trace Formulas and Discrete Subgroups*. Academic Press, 1989.
- [M-3] ——— Orbits of group actions and values of quadratic forms at integral points. *Israel J. Math.* 3 (1990), 127-150.
- [M-4] ——— Dynamical and ergodic properties of subgroup actions on homogeneous spaces with applications to number theory; in *Proceedings of the International Congress of Mathematicians (Kyoto 1990)*, The Mathematical Society of Japan, Tokyo; Springer-Verlag, 1991.
- [MZ] MONTGOMERY, D. and L. ZIPPIN. *Topological Transformations Groups*. Interscience Publishers, New York, 1955.
- [R] RATNER, M. Raghunathan's topological conjecture and distributions of unipotent flows, *Duke Math. J.* 63 (1991), 235-280.
- [S] SIKORAV, J.-C. Valeurs des formes quadratiques indéfinies irrationnelles; in *Séminaire de Théorie des Nombres, Paris 1987-88*, Progress in Math. 81, Birkhauser 1990.

(Reçu le 28 avril 1993)

S. G. Dani

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road
Bombay 400 005
India