

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNIMODULAR LATTICES WITH A COMPLETE ROOT SYSTEM
Autor: Kervaire, Michel
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-61105>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

UNIMODULAR LATTICES WITH A COMPLETE ROOT SYSTEM

by Michel KERVAIRE

1. INTRODUCTION

Let \mathbf{Q}^n be the n -dimensional euclidean space (over the field \mathbf{Q} of rational numbers) endowed with the standard scalar product

$$(x, y) = \sum_{i=1}^n x_i y_i ,$$

where $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$.

A lattice $L \subset \mathbf{Q}^n$ is a \mathbf{Z} -submodule of rank n of \mathbf{Q}^n , i. e.

$$L = \{ \sum_{i=1}^n a_i v_i : a_i \in \mathbf{Z} \} ,$$

where v_1, \dots, v_n is some basis of \mathbf{Q}^n . We are interested in *integral* lattices, i. e. lattices L satisfying $(x, y) \in \mathbf{Z}$ for all $x, y \in L$.

An integral lattice L is said to be *unimodular* if

$$\det(S) = \pm 1 ,$$

where S is the $n \times n$ matrix of scalar products

$$S = ((v_i, v_j)), \quad 1 \leq i, j \leq n ,$$

v_1, \dots, v_n being a \mathbf{Z} -basis of L . The number $\det(S)$ is called the *determinant* of L and is denoted $\det(L)$. It does not depend on the choice of the \mathbf{Z} -basis v_1, \dots, v_n of L .

If L is an integral lattice, the set

$$R = \{x \in L : (x, x) = 2\}$$

is a *root system*. (For the general notion of a root system see [B], p. 142.)

The author gratefully acknowledges partial support from the Fonds National Suisse de la Recherche Scientifique during the preparation of this paper. In particular the FNSRS provided the necessary computer.

The root system R will be said to be *complete* in L if the sublattice $N = \mathbf{Z}R$ of L generated by the roots R is a subgroup of finite index in L .

Our purpose is to study unimodular lattices with a complete root system.

It is well known that there are finitely many isomorphism classes of unimodular lattices $L \subset \mathbf{Q}^n$ for a given n . (See [MH], p. 18.)

The subcollection consisting of the lattices with a complete root system is particularly interesting, e.g. in view of the connection with the theory of error-correcting codes as we shall recall below.

We begin by setting up some necessary conditions that a root system must satisfy in order to be a complete root system in a unimodular lattice (Sections 3, 4 and 5).

We are particularly interested in even unimodular lattices, i. e. (x, x) is even for every $x \in L$. In this case, as is well known, the rank of L has to be divisible by 8. In dimensions 8, 16 and 24, where the classification of even unimodular lattices is available, it turns out that every such lattice has a complete root system, with the sole exception of the 24-dimensional Leech lattice. (History and relevant literature in e.g. [N], p. 142.)

In dimension 32, there are millions of even unimodular lattices. (See [Se], p. 95.) Among them as we shall see, only a small subcollection have a complete root system. In this paper, we endeavour to provide the complete list of such lattices.

There are 132 indecomposable even unimodular 32-dimensional lattices with a complete root system. In some cases several lattices happen to have the same root system. Thus, only a total number of 119 root systems correspond to these lattices. They are listed in Section 6.

The enumeration of the lattices and their root systems could only be completed using a computer, thanks to the generous help of Shalom Eliahou who patiently explained to me the use of mulisp programming language. Of course any mistake in the programs is my sole responsibility. It is a pleasure to express to him here my warmest gratitude.

I am also deeply indebted to Boris Venkov for very valuable discussions, in particular on the use of the notion of deficiency. (See Section 5.)

2. RELATIONSHIP WITH CODES

As is customary we shall use codes to describe lattices. We briefly recall how this can be done.

If $X \subset \mathbf{Q}^n$ is any finitely generated \mathbf{Z} -submodule of \mathbf{Q}^n , we set

$$X^\# = \{u \in \mathbf{Q}X : (u, x) \in \mathbf{Z} \text{ for all } x \in X\}.$$