Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNIMODULAR LATTICES WITH A COMPLETE ROOT SYSTEM
Autor: Kervaire, Michel
Kapitel: 5. The deficiency
DOI: https://doi.org/10.5169/seals-61105

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Thus,

$$f(u) = \prod_{i \in S} (1 + (q_i - 1)x_i) \cdot \prod_{i \in S'} (1 - x_i) ,$$

where $S \subset \{1, \ldots, s\}$ is the set of indices i for which $u_i = 0$, and $S' \subset \{1, \ldots, s\}$ the set of indices i for which $u_i \neq 0$.

Another way of writing $f(u)$ is

$$f(u) = \prod_{i=1}^{s} (1 - x_i)^{w(u_i)} \cdot (1 + (q_i - 1)x_i)^{1 - w(u_i)} .$$

Plugging this formula into $\sum_{u \in M} f(u)$, we get

$$\begin{aligned} \textstyle\sum_{u \in M} f(u) &= \textstyle\prod_{i=1}^{s} (1 + (q_i - 1)x_i) \cdot \sum_{u \in M} \prod_{i=1}^{s} \left(\frac{1 - x_i}{1 + (q_i - 1)x_i}\right)^{w(u_i)} \\ &= \textstyle\prod_{i=1}^{s} (1 + (q_i - 1)x_i) \cdot P_M \left(\frac{1 - x_1}{1 + (q_1 - 1)x_1}, \ldots, \frac{1 - x_s}{1 + (q_s - 1)x_s}\right) . \end{aligned}$$

Comparing the two expressions for $\sum_{u \in M} f(u)$, we get the theorem.

5. The deficiency

The main further necessary condition for a root system to be contained in an even unimodular lattice of the same rank is provided by the notion of deficiency (Defekt) introduced and studied in [KV].

If R is a root system of rank n, the *deficiency* of R, denoted $d(R)$, is the difference $n - m$, where m is the maximal cardinality of a set $\{a_1, \ldots, a_m\} \subset R$ of mutually orthogonal roots

$$(a_i, a_j) = 2\delta_{ij}, \qquad \text{for all } 1 \leqslant i, j \leqslant m .$$

We use this notion only if all roots in R have the same scalar square 2.

If $R = R_1 \boxplus R_2$, then $d(R) = d(R_1) + d(R_2)$. The values of the deficiency for the irreducible root systems are

$$d(\mathbf{A}_l) = \left[\tfrac{l}{2}\right] ,$$

$$d(\mathbf{D}_l) = \begin{cases} 0 & \text{for } l \text{ even}, \\ 1 & \text{for } l \text{ odd} , \end{cases}$$

$$d(\mathbf{E}_6) = 2, \; d(\mathbf{E}_7) = d(\mathbf{E}_8) = 0 .$$

By Satz 5 of [KV], if R is the (complete) root system of an even unimodular lattice of rank 32, then

$$d(R) = 0, 8, 12, 14, 15 \text{ or } 16 .$$

The proof consists in constructing from the given lattice a new lattice L, still of rank 32 and containing the orthogonal sum of $m = 32 - d(R)$ copies of $\mathbf{Z}$. Thus, $L = \mathbf{Z}^m \boxplus L_0$, where L_0 is again unimodular and of rank $d(R)$. (Hence, $\operatorname{rank}(L_0) \leqslant 16$.)

By Martin Kneser's classification of unimodular (positive definite) lattices of rank $\leqslant 16$, the rank of L_0, i.e. $d(R)$ can only take the above values. (See [Kn], Satz 1.)

In setting up the tables we conveniently use the deficiency to discriminate the various root systems R according to the value of $d(R)$.

6. The tables

We now proceed to list the *indecomposable* even unimodular lattices L of rank 32 with a complete root system R.

The presence in R of a factor of type $\mathbf{E}_8$ would produce a unimodular sublattice $\mathbf{ZE}_8 = L_0 \subset L$, and hence a decomposition $L = L_0 \boxplus L_1$ for some (even) unimodular L_1 of rank 24. Hence, we assume throughout that R has the form

$$R = \mathbf{A}_{l_1} \boxplus \ldots \boxplus \mathbf{A}_{l_r} \boxplus \mathbf{D}_{m_1} \boxplus \ldots \boxplus \mathbf{D}_{m_s} \boxplus \mathbf{mE}_6 \boxplus \mathbf{nE}_7 ,$$

with no factor of type $\mathbf{E}_8$.

Altogether there are $N = 88523$ such systems (of rank 32). The possible dimensions for $\mathbf{mE}_6 \boxplus \mathbf{nE}_7$ are

$$D = \{0, 6, 7, 12, 13, 14, 18, 19, 20, 21, 24, 25, 26, 27, 28, 30, 31, 32\}$$

and for $d \in D$, there is a unique pair (m, n) such that $d = 6m + 7n$. Hence

$$N = \textstyle\sum_{d \in D} \sum_{i=0}^{32-d} p(i) q(32 - d - i) ,$$

where $p(i)$ is the number of partitions of i and $q(j)$ is the number of partitions $(j_1, \ldots, j_t)$ of j with $4 \leqslant j_1 \leqslant \ldots \leqslant j_t$. (Of course, we use the convention $p(0) = q(0) = 1$.)

Among these, only 21209 have an acceptable deficiency, i.e. $d = 0, 8, 12, 14, 15$ or 16. They are distributed as follows: