Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ENUMERATIVE COMBINATORICS AND CODING THEORY
Autor: Eliahou, Shalom
Kapitel: 5. The number of Hadamard matrices of order n
DOI: https://doi.org/10.5169/seals-61109

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

and therefore

$$\alpha_\gamma = \frac{1}{2^{N-\alpha-\dim L}\,\alpha!} P_K^{(\alpha)}(-1) .$$

Multiplying both sides by $2^{n-\dim L}$, and plugging in equation (1), we obtain the claimed formula for $|f^{-1}(v)|$. □

COROLLARY 5. *Let* v_{min} *be the least value assumed by* f *on binary points. Then*

$$\frac{1}{2}(N + v_{min}) = \textit{the order of } -1 \textit{ as a root of } P_K(T) . \qquad \square$$

5. THE NUMBER OF HADAMARD MATRICES OF ORDER n

A *Hadamard matrix* is a square matrix H of order n with entries in $\{+1, -1\}$, satisfying the relation

$$H \cdot H^\top = nI_n .$$

($H^\top$ denotes the transpose of H, and I_n the identity matrix of order n.)

It is well known that the order of a Hadamard matrix can only be 1, 2 or a multiple of 4. Conversely, the existence of a Hadamard matrix of order n for every $n \equiv 0 \bmod 4$ is a longstanding conjecture, due to Jacques Hadamard [H]. The smallest open case currently occurs at $n = 428$. For a survey on Hadamard matrices, see [SY].

The theory exposed above yields a counting formula for Hadamard matrices of order n, in terms of the weight enumerator of a certain binary linear code of length $\binom{n}{2}^2$.

STEP 1. *Defining equations for Hadamard matrices.*

We represent binary matrices of order n as points $p = (p_{i,j}) \in \{1, -1\}^{n^2}$. Considering n^2 variables $\{x_{i,j}\}_{1 \leqslant i,j \leqslant n}$, let

$$g_{k,l} = \sum_{r=1}^{n} x_{k,r} x_{l,r} .$$

If $p = (p_{i,j})$ is a binary matrix, then $g_{k,l}(p)$ is the dot product of the k-th and l-th rows of p. Thus, a binary matrix p is Hadamard if and only if

$$g_{k,l}(p) = 0 \quad \text{for all } 1 \leqslant k < l \leqslant n .$$

STEP 2. *Reduction to a single equation.*

Let

$$g = \sum_{1 \leqslant k < l \leqslant n} g_{k,l}^2 .$$

By construction, we have the following properties:

(1) $g(p) \geqslant 0$ for every binary matrix p;

(2) $g(p) = 0$ if and only if p is Hadamard.

Developing the expression for g, we obtain:

$$\begin{aligned} g &= \textstyle\sum_{k<l} g_{k,l}^2 \\ &= \textstyle\sum_{k<l} (\sum_r x_{k,r} x_{l,r})^2 \\ &= \textstyle\sum_{k<l} (n + 2 \sum_{r<s} x_{k,r} x_{l,r} x_{k,s} x_{l,s}) \\ &= n \binom{n}{2} + 2f, \end{aligned}$$

where

$$f := \sum_{k<l} \sum_{r<s} x_{k,r} x_{l,r} x_{k,s} x_{l,s} .$$

(Of course, the above computation is performed modulo the relations $x_{i,j}^2 = 1$ for all i, j.)

The following properties of $f = \frac{1}{2}\left(g - n\binom{n}{2}\right)$ derive instantly from those of g:

(1) $f(p) \geqslant -\frac{1}{2} n \binom{n}{2}$ for every binary matrix p;

(2) $f(p) = -\frac{1}{2} n \binom{n}{2}$ if and only if p is Hadamard.

STEP 3. *The code associated with f.*

Let $K_n := L_f^{\perp}$ denote the dual of the binary code L_f associated with f, as defined in Section 3. Explicitly, we consider the map

$$\begin{array}{rccc} \phi_n: & \mathbf{F}_2^{\binom{n}{2}^2} & \to & \mathbf{F}_2^{n^2} \\ & E(k, l; r, s) & \mapsto & e_{k,r} + e_{l,r} + e_{k,s} + e_{l,s} , \end{array}$$

where $\{E(k, l; r, s)\}_{1 \leqslant k < l \leqslant n, 1 \leqslant r < s \leqslant n}$ and $\{e_{i,j}\}_{1 \leqslant i, j \leqslant n}$ denote the standard bases of the left and right spaces, respectively; by construction then, $K_n = \operatorname{Ker}(\phi_n)$.

As a direct consequence of Theorem 4 and of the above-mentioned properties of f, we obtain the

THEOREM 6. *Let* K_n *(n even) be the code of length* $\binom{n}{2}^2$ *defined as the kernel of the above map* $\phi_n : \mathbf{F}_2^{\binom{n}{2}^2} \to \mathbf{F}_2^{n^2}$. *Let* $P_n(T)$ *denote the weight enumerator of* K_n. *Then the number* $h(n)$ *of Hadamard matrices of order* n *is given by*

$$h(n) = \frac{1}{2^{\beta(n)}\,\alpha(n)!} \cdot P_n^{(\alpha(n))}(-1)\,,$$

where

1. $\alpha(n) = n^2(n-1)(n-2)/8$;
2. $\beta(n) = n^3(n-1)/8 - n^2$;
3. $P_n^{(\alpha(n))}(-1)$ *denotes the* $\alpha(n)$*-th derivative of* $P_n(T)$, *evaluated at* -1.

Proof. In the formula of Theorem 4, replace:

- N, the length of the code, by $\binom{n}{2}^2$;
- v, a lower bound for the values of f, by $-\frac{1}{2}n\binom{n}{2}$; and
- n, the number of variables in f, by n^2. □

Thus, the determination of the weight enumerator of K_n is an important problem. We will give below, without proof, the number of codewords of weight 3, 4 and 5 of K_n. (Of course, there are no words of weight 1 or 2 in K_n.) But the problem can be generalized a little bit, as follows. Consider the map

$$\begin{aligned} \phi_{m,n} : \quad \mathbf{F}_2^{\binom{m}{2}\binom{n}{2}} &\to \mathbf{F}_2^{mn} \\ E(k,l;r,s) &\mapsto e_{k,r} + e_{l,r} + e_{k,s} + e_{l,s}\,, \end{aligned}$$

where now, the indices $k < l$ range from 1 to m instead of 1 to n. We denote by $K_{m,n}$ the kernel of $\phi_{m,n}$.

Let $\Gamma = \{1, \ldots, m\} \times \{1, \ldots, n\}$. We can think of the vector basis $e_{i,j}$ as the point on row i and column j in the grid Γ, and of $E(k,l;r,s)$ as the rectangle determined by rows k, l and columns r, s in Γ. The image of $E(k,l;r,s)$ under $\phi_{m,n}$, then, is the formal sum of its four corners.

Thus, an element of weight w in $K_{m,n}$ can be pictured as a set of w rectangles in the grid Γ, such that every point in the grid coincides with an *even* number of corners of the rectangles in the set.

For example, all elements of weight 3 in $K_{m,n}$ can be represented (up to proper size and location) by the following picture:



or its vertical analogue. This picture represents a codeword of the form

$$E(k, l; r_1, r_2) + E(k, l; r_1, r_3) + E(k, l; r_2, r_3) .$$

Thus, the number of codewords of weight 3 in $K_{m,n}$ is equal to

$$w_3(K_{m,n}) = \binom{m}{2}\binom{n}{3} + \binom{m}{3}\binom{n}{2} .$$

Similarly, one can show that

$$w_4(K_{m,n}) = 3\binom{m}{2}\binom{n}{4} + 9\binom{m}{3}\binom{n}{3} + 3\binom{m}{4}\binom{n}{2} ;$$

$$w_5(K_{m,n}) = 12\binom{m}{2}\binom{n}{5} + 72\binom{m}{3}\binom{n}{4} + 72\binom{m}{4}\binom{n}{3} + 12\binom{m}{2}\binom{n}{5} + 9\binom{m}{3}\binom{n}{3} .$$

As a last remark, note that an upper bound for the weights in the associated code L_f is given by $\frac{1}{8}n^3(n-1)$, and that this bound is actually attained for some n if and only if there exists a Hadamard matrix of order n. This follows from, say, Corollary 3.

6. The number of proper 4-colorings of a graph

Let $G = (V, E)$ be a simple graph (no loops, no multiple edges) with vertex set V and edge set E. We will identify V with $\{1, \ldots, n\}$, and denote the cardinality of E by e.

A 4-*coloring* of G is the assignment to every vertex of one among four fixed colors; such a coloring is *proper* if the colors assigned to the end vertices of any edge are distinct. For a survey on the 4-colorings of planar graphs, see [SK].

We will count the number of proper 4-colorings of G, in terms of the weight enumerator of a certain code of length $3e$.

Step 1. *The defining equations for proper 4-colorings.*

As our palette of colors, we will choose the 4-set $\{1, -1\}^2$. The space of all 4-colorings of G can thus be identified with $\{1, -1\}^{2n}$, for example as follows: