

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ENUMERATIVE COMBINATORICS AND CODING THEORY
Autor: Eliahou, Shalom
Kapitel: 6. The number of proper 4-colorings of a graph
DOI: <https://doi.org/10.5169/seals-61109>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)


Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

For example, all elements of weight 3 in $K_{m,n}$ can be represented (up to proper size and location) by the following picture:



or its vertical analogue. This picture represents a codeword of the form

$$E(k, l; r_1, r_2) + E(k, l; r_1, r_3) + E(k, l; r_2, r_3).$$

Thus, the number of codewords of weight 3 in $K_{m,n}$ is equal to

$$w_3(K_{m,n}) = \binom{m}{2} \binom{n}{3} + \binom{m}{3} \binom{n}{2}.$$

Similarly, one can show that

$$w_4(K_{m,n}) = 3 \binom{m}{2} \binom{n}{4} + 9 \binom{m}{3} \binom{n}{3} + 3 \binom{m}{4} \binom{n}{2};$$

$$w_5(K_{m,n}) = 12 \binom{m}{2} \binom{n}{5} + 72 \binom{m}{3} \binom{n}{4} + 72 \binom{m}{4} \binom{n}{3} + 12 \binom{m}{2} \binom{n}{5} + 9 \binom{m}{3} \binom{n}{3}.$$

As a last remark, note that an upper bound for the weights in the associated code L_f is given by $\frac{1}{8}n^3(n-1)$, and that this bound is actually attained for some n if and only if there exists a Hadamard matrix of order n . This follows from, say, Corollary 3.

6. THE NUMBER OF PROPER 4-COLORINGS OF A GRAPH

Let $G = (V, E)$ be a simple graph (no loops, no multiple edges) with vertex set V and edge set E . We will identify V with $\{1, \dots, n\}$, and denote the cardinality of E by e .

A *4-coloring* of G is the assignment to every vertex of one among four fixed colors; such a coloring is *proper* if the colors assigned to the end vertices of any edge are distinct. For a survey on the 4-colorings of planar graphs, see [SK].

We will count the number of proper 4-colorings of G , in terms of the weight enumerator of a certain code of length $3e$.

STEP 1. *The defining equations for proper 4-colorings.*

As our palette of colors, we will choose the 4-set $\{1, -1\}^2$. The space of all 4-colorings of G can thus be identified with $\{1, -1\}^{2n}$, for example as follows:

Convention. If $p = (p_1, \dots, p_{2n}) \in \{1, -1\}^{2n}$, then the pair (p_i, p_{i+n}) represents the color assigned to vertex i , for $i = 1, \dots, n$.

Consider now $2n$ variables x_1, \dots, x_{2n} , and define

$$g_{i,j} := (1 + x_i x_j) (1 + x_{i+n} x_{j+n})$$

for $1 \leq i, j \leq n$.

If p is a 4-coloring of G , then $g_{i,j}(p) = 0$ if and only if the colors assigned to vertices i and j are distinct.

Thus, a 4-coloring p of G is proper if and only if

$$g_{i,j}(p) = 0 \quad \text{for all } (i, j) \in E.$$

STEP 2. Reduction to a single equation.

Let

$$g := \sum_{(i,j) \in E} g_{i,j}^2.$$

By construction, g satisfies the following properties:

- (1) $g(p) \geq 0$ for all 4-colorings p ;
- (2) $g(p) = 0$ if and only if p is proper.

Developing the expression for g , we obtain:

$$\begin{aligned} g &= \sum_E g_{i,j}^2 \\ &= \sum_E (1 + x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n})^2 \\ &= 4 \sum_E (1 + x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n}) \\ &= 4e + 4f, \end{aligned}$$

where

$$f := \sum_E (x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n}).$$

(Here again, the computation was performed modulo $x_i^2 = 1$ for all i .)

Obviously, f satisfies the following properties:

- (1) $f(p) \geq -e$ for all 4-colorings p ;
- (2) $f(p) = -e$ if and only if p is proper.

STEP 3. *The code associated with f .*

Let $K_G := L_f^\perp$ be the dual of the code L_f associated with f . To describe it, we consider the map

$$\begin{aligned}\phi_G : \mathbf{F}_2^{3e} &\rightarrow \mathbf{F}_2^{2n} \\ E_{i,j} &\mapsto e_i + e_j \\ E_{i+n, j+n} &\mapsto e_{i+n} + e_{j+n} \\ E_{i+2n, j+2n} &\mapsto e_i + e_j + e_{i+n} + e_{j+n} \quad ((i,j) \in E)\end{aligned}$$

where $\{E_{i,j}, E_{i+n, j+n}, E_{i+2n, j+2n}\}_{(i,j) \in E}$ and $\{e_i\}_{i \in V}$ denote the standard bases of the left and right spaces, respectively. By construction, $K_G = \text{Ker}(\phi_G)$.

Here again, as a direct consequence of Theorem 4 and of the stated properties of f , we have:

THEOREM 7. *Let K_G be the code of length $3e$ defined above, and let $P_G(T)$ denote its weight enumerator. Then, the number $\chi_G(4)$ of proper 4-colorings of G , is given by*

$$\chi_G(4) = \frac{1}{4^{e-n} e!} P_G^{(e)}(-1),$$

where $P_G^{(e)}(-1)$ denotes the value at -1 of the e -th derivative of $P_G(T)$.

Proof. Apply the formula of Theorem 4 by replacing

- N , the length of the code, by $3e$;
- v , a lower bound for the range of f , by $-e$;
- n , the number of variables in f , by $2n$. □

Note that there are some obvious elements of weight 3 in K_G , namely

$$E_{i,j} + E_{i+n, j+n} + E_{i+2n, j+2n} \quad ((i,j) \in E).$$

More interestingly, every cycle of length r in G gives rise to at least 3 elements of weight r in K_G . Indeed, if (i_1, \dots, i_r) is a cycle, then

$$E_{i_1+kn, i_2+kn} + E_{i_2+kn, i_3+kn} + \cdots + E_{i_r+kn, i_1+kn} \in K_G,$$

for $k = 0, 1, 2$.

One can go a little bit further. A somewhat technical computation shows that $P_G(T)$ can be decomposed as follows:

$$P_G(T) = (1 + T^3)^e P_{\tilde{K}} \left(\frac{T}{1 - T + T^2} \right),$$

where \tilde{K} is the \mathbf{F}_4 -code defined by the exact sequence

$$0 \rightarrow \tilde{K} \rightarrow \mathbf{F}_4^e \rightarrow \mathbf{F}_4^n ,$$

with weight enumerator $P_{\tilde{K}}(T)$. (The map on the right sends the basis vector corresponding to an edge, to the formal sum of its two endvertices.)

With the above formula, we find that $P_G^{(e)}(-1) = 3^e e! P_{\tilde{K}}\left(-\frac{1}{3}\right)$. Plugging this into Theorem 7, we obtain

$$\chi_G(4) = \frac{3^e}{4^{e-n}} P_{\tilde{K}}\left(-\frac{1}{3}\right) .$$

In particular, G is 4-colorable if and only if $P_{\tilde{K}}\left(-\frac{1}{3}\right) \neq 0$. See also [E, Theorem 5.7], for a different formulation and proof of this formula.

REFERENCES

- [B] BARKER, R. H. Group synchronizing of binary digital systems. In *Communication Theory*, W. Jackson, Ed., London: Butterworth, 1953, 273-287.
- [E] ELIAHOU, S. An algebraic criterion for a graph to be four-colourable. *Aportaciones Matemáticas, Notas de Investigación* 6 (1992), 3-27.
- [EKS] ELIAHOU, S., M. KERVAIRE and B. SAFFARI. On Golay polynomial pairs. *Adv. Appl. Math.* 12 (1991), 235-292.
- [G] GOLAY, M. J. E. Static multislit spectrometry and its application to the panoramic display of infrared spectra. *J. Opt. Soc. Amer.* 41 (1951), 468-472.
- [H] HADAMARD, J. Résolution d'une question relative aux déterminants. *Bull. Sci. Math. (2)* 17 (1893), 240-246.
- [M] MACWILLIAMS, F. J. A theorem on the distribution of weights in a systematic code. *Bell Syst. Tech. J.* 42 (1963), 79-94.
- [MS] MACWILLIAMS, F. J. and N. J. A. SLOANE. *The Theory of Error-Correcting Codes*. North-Holland, 1977.
- [SK] SAATY, T. L. and P. C. KAINEN. *The Four-Color Problem, Assaults and Conquest*. McGraw-Hill, New York, 1977.
- [SY] SEBERRY, J. and M. YAMADA. Hadamard matrices, Sequences, and Block Designs. In *Contemporary Design Theory: A Collection of Surveys*, J. H. Dinitz and D. R. Stinson, Eds., Wiley-Interscience, New York, 1992, 431-560.

(Reçu le 16 novembre 1993)

Shalom Eliahou

Section de Mathématiques
Université de Genève
C.P. 240
1211 Genève 24, Switzerland
e-mail: shalom@sc2a.unige.ch