

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE THEOREM OF KERÉKJÁRTÓ ON PERIODIC HOMEOMORPHISMS OF THE DISC AND THE SPHERE
Autor: Constantin, Adrian / Kolev, Boris
Kapitel: 4. Periodic homeomorphisms of the sphere
DOI: <https://doi.org/10.5169/seals-61111>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. PERIODIC HOMEOMORPHISMS OF THE SPHERE

The main result of this section is

THEOREM 4.1. *Let $f: S^2 \rightarrow S^2$ be a periodic homeomorphism. Then there exists $r \in O(3)$ and a homeomorphism $h: S^2 \rightarrow S^2$ such that $f = hrh^{-1}$.*

Proof of 4.1. We will divide the proof of Theorem 4.1 into two cases according to whether or not f has at least one fixed point.

Suppose first that f has a fixed point. Using 2.5, we deduce the existence of an invariant simple closed curve c which divides S^2 into two invariant discs D_1 and D_2 .

If f is orientation preserving and $f \neq Id$, then f has no fixed point on c (cf. 3.2). Therefore, by Brouwer's fixed point theorem we know then that f has at least two fixed points; after a conjugacy, we can suppose that f fixes the two poles N and S of S^2 . Using the results of last section, we are able to find n arcs joining N and S such that their union is an invariant set under f . As in Section 3, we can then construct a conjugacy between f and a rotation by angle $2k\pi/n$ around the South-North axis.

If f is orientation-reversing, then f has two fixed points on c . In each of the invariant disc D^1 and D^2 , the fixed point set of f consists of a simple arc which joins the two fixed points of f on c . The union of these two arcs is a simple closed curve which coincides with the fixed point set of f on S^2 . It is then easy to construct a conjugacy between f and the reflection about the equator.

Let now suppose that f has no fixed point on S^2 . Up to conjugacy, we can assume that the second iterate of f , f^2 is a periodic rotation around the North-South axis. In particular the points N and S are exchanged by f . For $t \in (-1, 1)$, let C_t be the circle obtained by cutting the sphere by the plane $z = t$, D_t the disc bordered by C_t on S^2 which contains N and:

$$t_0 = \inf \{ t \in (-1, 1) ; D_t \cap f(D_t) = \emptyset \} .$$

We write $D = D_{t_0}$ and $C = C_{t_0}$ for convenience. Then D meets $f(D)$ on its boundary and only on its boundary (see Figure 4). Let $P_0 \in C \cap f(C)$ and P_1, P_2, \dots, P_{n-1} , the orbit of P_0 under f . The points P_0, P_2, \dots, P_n and P_1, P_3, \dots, P_{n-1} are distinct because f^2 is a rotation of period $n/2$.

Suppose that there exists $i \in \{1, 3, \dots, n-1\}$ such that P_0 and $P_i = f^i(P_0)$ coincide. Then P_0, S and N are fixed by f^{2i} so $f^{2i} = Id$. Therefore $2i = n$.

Let b_0 be the arc of great circle that joins N to P_0 in D and $b_{n/2}$ its image under $f^{n/2}$. Then $b = b_0 \cup b_{n/2}$ is a simple arc joining N and S and not meeting its first $(n/2) - 1$ iterates under f away from N and S . These arcs divide the sphere into $n/2$ sectors and we can build a conjugacy between f and the composition of a rotation of period $n/2$ around the North-South axis with a reflexion about the equator.

Suppose now that the points P_0, P_1, \dots, P_{n-1} are distinct. Let b_0 an arc of great circle joining N and P_0 in D and b'_0 an arc joining S to P_0 in $f(D)$ disjoint from $f(b_0)$ and from its first $n - 1$ iterates (which is possible since f^2 is a rotation). The union of these two arcs is again a simple arc joining N and S which does not meet its first $n - 1$ iterates under f away from N and S . The union of this arc and its iterates divides the sphere S^2 into n disjoint sectors. In that case, f is topologically equivalent to the composition of a rotation of period n around the North-South axis with a reflexion about the equator. \square




FIGURE 4

COROLLARY 4.2. *Let $f: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be a periodic homeomorphism. Then f is topologically conjugate to a finite order rotation around the origin or to the reflexion about the x -axis.*

Proof of 4.2. We can extend f to a homeomorphism of the Sphere S^2 by identifying the plane \mathbf{R}^2 with the complement of the North pole using the stereographic projection. Looking at the proof of 4.1, f is either equivalent to a rotation around the North-South pole or to a reflexion about a great circle which we can assume to pass through the north pole N . It is not difficult to

show that the conjugacy can be chosen to fix also the North pole N . This equivalence induces, therefore, a topological equivalence between f and a rotation or a reflexion about the x -axis. \square

Remark. The investigation of periodic homeomorphisms on surfaces of positive genus has been studied extensively. We cannot give here a complete bibliography on the subject. We would just like to cite original works of Kerékjártó [4] and Nielsen [13] which lead to the conclusion that a periodic homeomorphism of a Riemannian surface of positive genus is conjugate to a conformal isometry.

Acknowledgement. The authors express their gratitude to Jérôme Fehrenbach, Lucien Guillou and Toby Hall for the discussions that helped to improve this paper.

REFERENCES

- [1] BROUWER, L.E.J. Über die periodischen Transformationen der Kugel. *Math. Ann.* 80 (1919), 39-41.
- [2] CAIRNS, S.S. An elementary proof of the Jordan-Schoenflies theorem. *Proc. AMS* 2 (1951), 860-867.
- [3] KERÉKJÁRTÓ de, B. Über die periodischen Transformationen der Kreisscheibe und der Kugelfläche. *Math. Ann.* 80 (1919-1920), 36-38.
- [4] ——— Über reguläre Abbildungen von Flächen auf sich. *Acta scient. math. Szeged* 7 (1934), 65-75.
- [5] DEVANEY, R. *An Introduction to Chaotic Dynamical Systems*. Benjamin-Comings 1986.
- [6] EILENBERG, S. Sur les transformations périodiques de la surface de la sphère. *Fund. Math.* 22 (1934), 28-44.
- [7] EPSTEIN, D.B.A. Pointwise periodic homeomorphisms. *Proc. London Math. Soc.* 42 (3) (1981), 415-460.
- [8] HOCKING, J.G. and G.S. YOUNG. *Topology*. Dover 1988.
- [9] MAEHARA, R. The Jordan curve theorem via the Brouwer fixed point theorem. *Amer. Math. Monthly* 91 (1984), 641-643.
- [10] MASKIT, B. *Kleinian Groups*. A Series of Comprehensive Studies in Mathematics, Springer Verlag 1988.
- [11] MILNOR, J. *Dynamics in one complex variable*. Preprint, Stony Brook, New York 1990.
- [12] NEWMANN, M.H.A. *Elements of the Topology of Plane Sets of Points*. Cambridge Univ. Press (2nd edition) 1951.