

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 40 (1994)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNITS OF CLASSICAL ORDERS: A SURVEY
Autor: Kleinert, Ernst
Kapitel: 2. Elementary Properties
DOI: <https://doi.org/10.5169/seals-61112>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. ELEMENTARY PROPERTIES

(1) Let N denote the regular norm $A^\times \rightarrow \mathbf{Q}^\times$. It is easy to see that $\Gamma = \{x \in \Lambda \mid N^2(x) = 1\}$.

If we specify a \mathbf{Z} -basis of Λ , this becomes a polynomial equation in the coefficients of x with respect to this basis, and the elements of Γ correspond precisely to the integral solutions. This shows that Γ is an *arithmetic group*, and thereby makes available all the general results on this class of groups. (A reference ideally suited to the present theme is Serre's survey article [Se4]; we also mention [Pl].) In fact, a good deal of the present paper will be concerned with specifying the general results to the case of unit groups.

(2) Let $\Lambda \subset \Lambda'$ be orders in A with unit groups Γ, Γ' . Then $\Gamma = \Gamma' \cap \Lambda$, and $|\Gamma' : \Gamma|$ is finite.

Proof. For any $x \in \Gamma$ we have $x^{-1} \in \mathbf{Z}[x]$ since x is a zero of a monic integral polynomial with constant term ± 1 . This proves the first statement. For the second, assume that, for $x, y \in \Gamma'$, we have

$$x - y = mz, \quad \text{where } m = |\Lambda' : \Lambda|, z \in \Lambda'.$$

Then

$$xy^{-1} = 1 + mzy^{-1} \in \Gamma' \cap \Lambda = \Gamma.$$

This shows that

$$|\Gamma' : \Gamma| \leq |\Lambda' : m\Lambda| = (\dim A)^m.$$

(2) allows us to reduce all questions concerning virtual properties of Γ to arbitrary orders in simple algebras. (A group is said to have a property virtually if a subgroup of finite index has that property). Finite presentability is such a property: if $\Gamma_0 \subset \Gamma$, $|\Gamma : \Gamma_0|$ finite, has a finite presentation, then so has, by Reidemeister-Schreier, the intersection of its conjugates, which is normal; now use the fact that the class of finitely presented groups is closed under extensions ([J], p. 187, Th. 1).

(3) Γ is *virtually torsion free*.

Proof. It is easy to see that there is an upper bound, and consequently a lowest common multiple N for the orders of torsion elements $x \in A^\times$; all such $x \neq 1$ satisfy

$$x^{N-1} + x^{N-2} + \dots + x + 1 = 0.$$

For $n \in \mathbf{N}$ let

$$\Gamma(n) = \text{kernel of } (\Gamma \rightarrow (\Lambda/n\Lambda)^\times)$$

the congruence group mod n ; this is a normal subgroup of finite index. Obviously $\Gamma(n)$ is torsion free for $n > N$. With more effort, one can do much better: the regular representation injects $\Gamma(n)$ into the congruence group mod n in $GL_m(\mathbf{Z})$, $m = \dim A$, and Minkowski has shown that this is torsion free for $n > 2$ [Mi].

(4) Γ contains only finitely many isomorphism classes of finite subgroups.

Proof. If $\Gamma_0 < \Gamma$ is torsion free and normal of finite index, then every finite subgroup of Γ is isomorphic to a subgroup of Γ/Γ_0 .

Later, we will show more: Γ contains only finitely many conjugacy classes of finite subgroups.

(5) Γ is residually finite, that is, for every $x \in \Gamma, x \neq 1$, there is a normal subgroup Γ_0 of finite index such that $x \notin \Gamma_0$.

Of course, almost all $\Gamma(n)$ will do. It follows that Γ is hopfian, that is, not isomorphic to a proper factor group (see [MKS], p. 116).

(6) Finally, let us mention here the following result due to Zassenhaus [Z2] (although it is not entirely elementary): Γ contains a solvable subgroup of finite index if and only if the Wedderburn components of A are number fields or definite quaternions over \mathbf{Q} .

Sketch of proof: the problem is readily reduced to simple A . The “If” part is then trivial.

Conversely, if matrices are involved, one knows that Γ has infinitely many subfactor groups of the form $SL_n(F)$, where F is a finite field. The same is therefore true of any subgroup of finite index. In the skew field case, the argument is more intricate; we refer to [Z2].

3. FINITE GENERATION: CLASSICAL REDUCTION THEORY

The most basic fact about Γ is that it is finitely generated; this is even valid for arbitrary arithmetic groups, as has been proved by A. Borel and Harish-Chandra in the fundamental paper [BHC]. Here I shall describe the classical approach, carried out by Siegel [S1], who completed earlier work of