Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 40 (1994)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNITS OF CLASSICAL ORDERS: A SURVEY

Autor: Kleinert, Ernst

Kapitel: 8. Congruence subgroups and normal subgroups

DOI: https://doi.org/10.5169/seals-61112

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Theorem 5. Suppose that A = D is not a totally definite quaternion algebra. Then there are isomorphisms

$$H^k(\Gamma, M) \cong H^{k+g}(\Gamma, M)$$

for all Γ -modules and M and k > r(D). Moreover, r(D) + 1 is the smallest dimension from which on $H^*(\Gamma, -)$ is n-periodic.

The last statement follows from the facts that (1) Γ is a virtual duality group and hence $H^{r(D)}(\Gamma, \mathbb{Z}\Gamma)$ is torsion free ([Br], VIII 11.2) whereas (2) for any group Γ with $vcd\Gamma = k$, $H^m(\Gamma)$ is torsion for m > k ([Se 3], p. 101).

It should be possible to refine the period g in the theorem by one more directly derived from the periods of the finite subgroups of Γ as in the number field case where the period equals 2 if there are nontrivial torsion units of norm 1 (in which case n must be even!).

REMARK. We have seen (in the general case, Γ torsionfree) that

$$H^*(\Gamma, -) = H^*(X(\Gamma), -).$$

Taking real coefficients (with trivial Γ -action) the latter groups are, by de Rham's theorem, given by differential forms on $X(\Gamma)$; these in turn correspond to Γ -automorphic forms on X. In this way, the real cohomology of Γ becomes part of the theory of automorphic forms.

8. Congruence subgroups and normal subgroups

Recall that we have defined

$$\Gamma(m) = \text{kernel of } (\Gamma \to (\Lambda/m\Lambda)^{\times}),$$

the congruence subgroup of level m of Γ . Obviously $\Gamma(m)$ has finite index in Γ . The following question is classical: does every subgroup of finite index of Γ contain a congruence group?

Let us say that Γ satisfies (CP) if this is so. Let $\Lambda \subset \Lambda'$. If Γ' satisfies (CP), so does Γ . To prove the converse, it suffices to show that every $\Gamma(n)$ contains a $\Gamma'(m)$. This will be so if Γ contains a $\Gamma'(m)$. But there is $m \in \mathbb{N}$ with $m\Lambda' \subset \Lambda$, and this implies $\Gamma'(m) \subset \Lambda \cap \Gamma' = \Gamma$. Thus, property (CP) depends only on A.

For A = K a number field, (CP) has essentially been proved by Chevalley [Ch]. Let $H < R^{\times}$ be of finite index, and $H_0 < R^{\times}$ any congruence subgroup. Then $H_0^k \subset H$ for some $k \in \mathbb{N}$; so it suffices to show

that any power of a congruence group contains a congruence group. This follows at once from ([Ch], Th. 1). By an argument presented earlier, this allows us to reduce the problem to $S\Gamma$. If D=K, the answer is given by results of Bass-Milnor-Serre [BMS] and Serre [Se 5]. (Of course, one defines congruence groups with respect to ideals of R. For (CP) to hold, this makes no difference since every ideal of R contains an nR, $n \in \mathbb{Z}$.)

THEOREM 6. Assume D = K.

- (a) [BMS] If $n \ge 3$, $SL_n(R)$ has property (CP) if and only if K has a real embedding.
- (b) [Se 5] $SL_2(R)$ has property (CP) if and only if r(K) > 0.

The only K with r(K) = 0 are $K = \mathbf{Q}$ and K = imaginary quadratic. The failure of (CP) for $SL_2(\mathbf{Z})$ is classic (see [Ne], p. 149). A discussion of the Bianchi case is given in [F, p. 200 ff.].

The case where $K \neq D$ and n > 1 was treated by Bak and Rehmann [BR]. They give an S-arithmetical result, from which we extract:

THEOREM 7. Suppose $K \neq D$ and $n \geqslant 2$; if n = 2 suppose $K \neq \mathbf{Q}$ and that D is not a definite quaternion algebra. Then (CP) holds.

The remaining cases seem still to be unsettled. Note that in the definite quaternion case the problem becomes trivial since $S\Gamma$ is finite.

The congruence groups are special since normal subgroups having finite index. It also makes good sense to ask whether every noncentral normal subgroup of $S\Gamma$ has finite index. This too may be asked more generally for discrete subgroups of Lie groups, and definite results have been obtained by Margulis. We specialize these to the case in question. As above, we can give no details of the proofs, which are extremely involved, and refer the reader to Margulis' monumental volume [M]. We make the following assumptions, excluding A = K:

(H)
$$\begin{cases} \text{(i) if } n = 1, \text{ then } D \text{ splits at all infinite primes of } K; \\ \text{(ii) if } K = \mathbf{Q}, \text{ then } k \geqslant 3, \text{ and } A \neq M_2(D), D \text{ definite quaternion.} \end{cases}$$

This means that the norm-1-group SG has no anisotropic factor and has rank ≥ 2 . It is clear that — in the terminology of [M] — $S\Gamma$ is an irreducible lattice of SG. Applying Theorems (2) and (4) from the introduction of [M], we obtain

THEOREM 8. Assume (H). Then every noncentral normal subgroup of $S\Gamma$ has finite index.

COROLLARY. $S\Gamma/[S\Gamma, S\Gamma]$ is finite.

Again, it is classic that the theorem fails for $SL_2(\mathbf{Z})$; for instance, if $n \ge 0$, the verbal subgroup generated by all 6n-th powers has infinite index ([Ne], p. 143). The corollary holds for $SL_2(\mathbf{Z})$ but fails for torsion free subgroups (which are free). Theorem 7 however carries over to subgroups of finite index.

Two more topics from the general theory of arithmetic groups, which could be specialized to unit groups, are subgroup rigidity and strong approximation. But having promised to keep as near as possible to the unit groups "themselves", we omit this.

9. The Bass unit theorem

In his paper [Ba2] Bass has proved (among other things) a far reaching generalization of Dirichlet's unit theorem which — together with the results of sections 3 and 7 — is certainly one of the strongest general results we have about Γ . The core of the proof is a deep stability theorem from K-theory; we will indicate how it implies the theorem but will say little about its proof. We begin with the relevant definitions. For any ring A, define

$$K_1(A) = \lim_{\longrightarrow} GL_n(A)/[GL_n(A), GL_n(A)],$$

where the direct limit is taken with respect to the embeddings

$$GL_n(A) \to GL_{n+1}(A), \quad x \to \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}.$$

One may also write

$$K_1(A) = \lim_{\longrightarrow} GL_n(A)/\tilde{E}_n(A)$$
,

where $\tilde{E}_n(A)$ is the normal subgroup generated by the elementary matrices; this is Whitehead's lemma. Further, with $K_0(A)$ denoting the Grothendieck group of finitely generated projective A-modules, we put

$$R_0(A) = \mathbf{R} \otimes K_0(A), R_1(A) = \mathbf{R} \otimes K_1(A).$$

Now we turn to algebras and allow A to be semisimple. Let $\Lambda \subset A$ be an order. Any A_R -module V (of finite dimension) gives rise to a homomorphism

$$\Lambda \to \operatorname{End}_{\mathbf{R}} V$$
, hence by functoriality $K_1(\Lambda) \to K_1(\operatorname{End}_{\mathbf{R}} V)$.