

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE COHOMOLOGY OF COMPACT LIE GROUPS
Autor: Reeder, Mark

Kurzfassung

DOI: <https://doi.org/10.5169/seals-61824>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON THE COHOMOLOGY OF COMPACT LIE GROUPS

by Mark REEDER

ABSTRACT. We give a new computation of the cohomology of a Lie group that some mathematicians may find to be shorter and more elementary than previous approaches. The main new ingredient is a result of L. Solomon on differential forms invariant under a finite reflection group. The cohomology is shown to have a bi-grading which has several interpretations.

1. INTRODUCTION

Let G be a compact connected Lie group, and let T be a maximal torus in G . We denote the corresponding Lie algebras by \mathfrak{g} and \mathfrak{t} . Let W be the Weyl group of T in G . Then W acts on \mathfrak{t} as a group generated by reflections about the kernels of the roots of \mathfrak{t} in $\mathfrak{g} \otimes \mathbf{C}$. It has been known since the first half of this century that the cohomology ring $H(G)$, with real coefficients, is an exterior algebra with generators in degrees $2m_1 + 1, \dots, 2m_l + 1$, where $m_1 + 1, \dots, m_l + 1$ are the degrees of homogeneous generators of the ring of W -invariant polynomial functions on \mathfrak{t} . In particular, the Poincaré polynomial of G is $(1 + t^{2m_1+1}) \cdots (1 + t^{2m_l+1})$, and G has the cohomology of a product of odd-dimensional spheres.

Despite its age and familiarity, it is not easy to find a proof of this theorem in the literature. There are many beginnings and sketches in the textbooks, but the difficult part, namely the remarkable connection between degrees of invariant polynomials and Betti numbers, usually goes unproven. One reason is that the standard proofs (for example, [Bo2], [Ch], [L]) require substantial algebraic preliminaries on Hopf algebras, spectral sequences, and differential algebras. (See [Bo1] and [Sam] for historical surveys.)

We offer here a new but less sophisticated computation of the cohomology of a Lie group, avoiding the above algebraic techniques. Instead we use