

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 41 (1995)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE COHOMOLOGY OF COMPACT LIE GROUPS  
**Autor:** Reeder, Mark  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-61824>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 10.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## ON THE COHOMOLOGY OF COMPACT LIE GROUPS

by Mark REEDER

**ABSTRACT.** We give a new computation of the cohomology of a Lie group that some mathematicians may find to be shorter and more elementary than previous approaches. The main new ingredient is a result of L. Solomon on differential forms invariant under a finite reflection group. The cohomology is shown to have a bi-grading which has several interpretations.

### 1. INTRODUCTION

Let  $G$  be a compact connected Lie group, and let  $T$  be a maximal torus in  $G$ . We denote the corresponding Lie algebras by  $\mathfrak{g}$  and  $\mathfrak{t}$ . Let  $W$  be the Weyl group of  $T$  in  $G$ . Then  $W$  acts on  $\mathfrak{t}$  as a group generated by reflections about the kernels of the roots of  $\mathfrak{t}$  in  $\mathfrak{g} \otimes \mathbf{C}$ . It has been known since the first half of this century that the cohomology ring  $H(G)$ , with real coefficients, is an exterior algebra with generators in degrees  $2m_1 + 1, \dots, 2m_l + 1$ , where  $m_1 + 1, \dots, m_l + 1$  are the degrees of homogeneous generators of the ring of  $W$ -invariant polynomial functions on  $\mathfrak{t}$ . In particular, the Poincaré polynomial of  $G$  is  $(1 + t^{2m_1+1}) \cdots (1 + t^{2m_l+1})$ , and  $G$  has the cohomology of a product of odd-dimensional spheres.

Despite its age and familiarity, it is not easy to find a proof of this theorem in the literature. There are many beginnings and sketches in the textbooks, but the difficult part, namely the remarkable connection between degrees of invariant polynomials and Betti numbers, usually goes unproven. One reason is that the standard proofs (for example, [Bo2], [Ch], [L]) require substantial algebraic preliminaries on Hopf algebras, spectral sequences, and differential algebras. (See [Bo1] and [Sam] for historical surveys.)

We offer here a new but less sophisticated computation of the cohomology of a Lie group, avoiding the above algebraic techniques. Instead we use

standard Lie theory and more invariant theory than is customary. We have tried to give a fairly complete treatment, in which it is seen that only a few ideas are used repeatedly. The most serious omission of proof is that of Chevalley's theorem on invariant polynomials, but that has many accessible references. Hopefully, enough background has been included to make the whole story coherent to someone with a basic knowledge of Lie groups and differential topology.

A sketch of our computation of  $H(G)$  goes as follows. Consider the manifold  $M$  consisting of pairs  $(g, T')$ , where  $g \in G$  and  $T'$  is a maximal torus in  $G$  which contains  $g$ . There is a natural map  $\psi: M \rightarrow G$ , known already to Weyl, given by conjugation. We make the apparently new observation that  $\psi$  induces an isomorphism of real cohomology rings  $H(M) \simeq H(G)$ . This uses only standard facts gleaned from the differential of  $\psi$ . One could also invoke the spectral sequence of the fibration  $G \rightarrow G/T$ . This spectral sequence was shown by Leray to degenerate at  $E^3$ . It in fact has a spectral subsequence (the  $W$ -invariants) which already degenerates at  $E^2$  and still computes  $H(G)$  (see (6.4) below).

We still have to compute the cohomology of  $M$ . It is easy to see that  $H(M) = [H(G/T) \otimes H(T)]^W$ . The ring  $H(T)$  is naturally isomorphic to the exterior algebra of  $\mathfrak{t}^*$ , and  $H(G/T)$  is isomorphic to the space  $\mathcal{H}$  of  $W$ -harmonic polynomials on  $\mathfrak{t}$ , according to a famous theorem of Borel. For completeness, we give a proof of this in the same elementary, if less efficient spirit. As with all proofs, the essential thing is to show that the odd cohomology of  $G/T$  vanishes. We do this with a direct generalization of the Morse-theoretic computation of the cohomology of the two-sphere.

So now we are down to invariant theory, and must compute  $[\mathcal{H} \otimes \Lambda \mathfrak{t}^*]^W$ . This follows immediately from Solomon's determination of the  $W$ -invariant differential forms on  $\mathfrak{t}$  with polynomial coefficients, which in turn depends on Chevalley's well-known description of  $W$ -invariant polynomials. This gives us the desired connection between degrees of  $W$ -invariants and Betti numbers of  $G$ . Solomon's result also leads to pretty formulas for the multiplicities of the  $W$ -modules  $\Lambda^q \mathfrak{t}^*$  in spaces of harmonic polynomials (see (3.8)), as well as a generalization of a classical result on the Jacobian of the basic invariants (see (3.9)).

The paper is organized as follows: First the structure of  $G$  and its adjoint representation is recalled, then comes invariant theory, followed by the proof of Borel's theorem, finishing with the computation of  $H(G)$  and some remarks on its natural bigrading. Throughout, cohomology has real coefficients.