

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 41 (1995)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE COHOMOLOGY OF COMPACT LIE GROUPS  
**Autor:** Reeder, Mark  
**Kapitel:** 2. Basic material  
**DOI:** <https://doi.org/10.5169/seals-61824>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 04.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 2. BASIC MATERIAL

For more details in this section see [A], for example.

(2.1) Recall that  $G$  is a compact connected Lie group with maximal torus  $T$ , having respective Lie algebras  $\mathfrak{g}$  and  $\mathfrak{t}$ . The Weyl group is the finite group  $W = N/T$ , where  $N$  is the normalizer in  $G$  of  $T$ . Since  $G$  is compact, there is an  $Ad(G)$ -invariant inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{g}$ , obtained by averaging any inner product over  $G$ . Let  $\mathfrak{m}$  be the orthogonal complement of  $\mathfrak{t}$  in  $\mathfrak{g}$  with respect to this inner product, so

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{m} \quad (\text{orthogonal}) .$$

The infinitesimal version of invariance of the inner product is the identity

$$\langle [X, Y], Z \rangle + \langle Y, [X, Z] \rangle = 0 ,$$

for  $X, Y, Z \in \mathfrak{g}$ .

(2.2) The exponential map  $\exp: \mathfrak{g} \rightarrow G$  is surjective, since  $G$  is compact. This is one of the deeper theorems in a first course on Lie groups. We actually only need this surjectivity for  $\exp: \mathfrak{t} \rightarrow T$ , which is clear.

The Lie algebra  $\mathfrak{t}$  is abelian (the bracket is zero); in fact  $\mathfrak{t}$  is a maximal abelian subalgebra of  $\mathfrak{g}$ . In particular, no nonzero vector in  $\mathfrak{m}$  has zero bracket with all of  $\mathfrak{t}$ . Likewise,  $Ad(T)$  has no nonzero invariant vectors in  $\mathfrak{m}$ .

Now a torus is a topologically cyclic group. That means there exists a *generic element*  $t_0 \in T$  whose powers form a dense subgroup of  $T$ . It follows that the single operator  $Ad(t_0)$  can have no invariants in  $\mathfrak{m}$ . Likewise in the group  $G$ , it can be shown that a maximal torus is its own centralizer, so the centralizer in  $G$  of  $t_0$  is just  $T$ . There is a similar notion in the Lie algebra. A *regular element* of  $\mathfrak{t}$  is one whose  $Ad(G)$ -centralizer is exactly  $Ad(T)$ . To find one, take any  $H_0 \in \mathfrak{t}$  such that  $\exp H_0 = t_0$ .

(2.3) The group  $G$  acts on  $\mathfrak{g}$  via  $Ad$ , and this induces an action of  $W$  on  $\mathfrak{t}$ . No element of  $W$  acts trivially, and the image of  $W$  in  $GL(\mathfrak{t})$  is generated by reflections about certain hyperplanes defined as follows.

Since the nontrivial irreducible representations of a torus are given by two dimensional rotations, we have an orthogonal decomposition  $\mathfrak{m} = \mathfrak{m}_1 \oplus \cdots \oplus \mathfrak{m}_\nu$ , where each  $\mathfrak{m}_k$  is two dimensional and there is a finite set of nonzero linear functionals  $\Delta^+ = \{\alpha_1, \dots, \alpha_\nu\} \subset \mathfrak{t}^*$ , called *positive roots* such that for  $H \in \mathfrak{t}$ , the eigenvalues of  $Ad \exp H$  on  $\mathfrak{m}_i$  are  $\exp(\pm \sqrt{-1} \alpha_i(H))$ . We determine the signs as follows. Fix a regular

element  $H_0 \in \mathfrak{t}$ . We may and shall choose the positive roots so that they take strictly positive values on  $H_0$ . The action of  $W$  on  $\mathfrak{t}$  is generated by reflections about the kernels of the positive roots.

Since each  $\mathfrak{m}_i$  is also preserved by  $ad(\mathfrak{t})$ , we can choose an orthonormal basis  $\{X_i, X_{v+i}\}$  of  $\mathfrak{m}_i$  such that, for  $H \in \mathfrak{t}$ , the matrix of  $ad(H)|_{\mathfrak{m}_i}$  with respect to this basis is

$$\begin{pmatrix} 0 & \alpha(H) \\ -\alpha(H) & 0 \end{pmatrix}.$$

Note that the  $ad$ -invariance of the inner product  $\langle \cdot, \cdot \rangle$  implies, for all  $1 \leq i \leq v$ , all  $1 \leq j \leq 2v$  and all  $H \in \mathfrak{t}$  that

$$\langle H, [X_i, X_j] \rangle = \langle [H, X_i], X_j \rangle = -\alpha_i(H) \langle X_{i+v}, X_j \rangle.$$

By orthonormality, this last pairing can only be nontrivial if  $j = i + v$ . Hence if  $j \neq i + v$ , we have  $[X_i, X_j] \in \mathfrak{m}$ . The same thing happens if  $i > v$  and  $j \neq i - v$ .

On the other hand, for  $1 \leq i \leq v$ , set  $H_i = [X_i, X_{v+i}]$ . This is  $Ad(T)$ -invariant, so  $H_i \in \mathfrak{t}$ , and  $ad(H_i)\mathfrak{m}_i \subseteq \mathfrak{m}_i$ . It follows that the span of  $X_i, X_{i+v}, H_i$  is a Lie subalgebra  $\mathfrak{g}_i$  of  $\mathfrak{g}$ . It is always isomorphic to  $\mathfrak{su}(2)$ .

### 3. INVARIANT THEORY

All proofs missing from this section may be found in the textbook [H], the expository article [F], or [Bk].

(3.1) Let

$$\mathcal{S} = \bigoplus_{p=0}^{\infty} \mathcal{S}^p \quad \text{and} \quad \Lambda = \bigoplus_{q=0}^l \Lambda^q \quad (l = \dim \mathfrak{t})$$

be the symmetric and exterior algebras on  $\mathfrak{t}^*$ , respectively. The adjoint action of  $W$  on  $\mathfrak{t}$  induces representations of  $W$  on  $\mathcal{S}$  and  $\Lambda$  by degree-preserving algebra automorphisms. For example, the action of  $W$  on  $\Lambda^l$  is multiplication by the *sign character*

$$\varepsilon: W \rightarrow \{\pm 1\} \quad \text{given by} \quad \varepsilon(w) = \det Ad(w)_{\mathfrak{t}}.$$

Note that  $\varepsilon(w)$  is the parity of the number of reflections needed to express  $Ad(w)_{\mathfrak{t}}$ .